arXiv:1210.2368v1 [math.AT] 8 Oct 2012

Toric Topology

Victor M. Buchstaber

Taras E. Panov

Steklov Mathematical Institute, Russian Academy of Sciences, Gubkina Street 8, 119991 Moscow, Russia
E-mail address: `buchstab@mi.ras.ru`

Department of Mathematics and Mechanics, Moscow State University, Leninskie Gory, 119991 Moscow, Russia
E-mail address: `tpanov@mech.math.msu.su`

Abstract. The theory of torus actions on topological spaces has a long history and constitutes an important field within algebraic topology. Over the last 20 years a new field emerged on the borders of equivariant topology, algebraic and symplectic geometry, combinatorics and commutative algebra. Under the name *"toric topology"* this field has quickly attracted the attention of specialists from different areas, and is currently under active development.

Contents

Introduction 1
  Chapter dependence guide 1

Chapter 1. Geometry and combinatorics of polytopes 3
  1.1. Convex polytopes 3
  1.2. Gale diagrams 11
  1.3. Face vectors and Dehn–Sommerville relations 11
  1.4. Characterising the face vectors of polytopes 17

 Polytopes: additional topics 23
  1.5. Nestohedra and graph-associahedra 23
  1.6. Flagtopes and truncated cubes 36
  1.7. Differential algebra of combinatorial polytopes 42
  1.8. Families of polytopes and differential equations 47

Chapter 2. Combinatorial structures 53
  2.1. Polyhedral fans 54
  2.2. Simplicial complexes 57
  2.3. Barycentric subdivision and flag complexes 62
  2.4. Alexander duality 65
  2.5. Classes of triangulated spheres 68
  2.6. Triangulated manifolds 73
  2.7. Stellar subdivisions and bistellar moves 76
  2.8. Simplicial posets and simplicial cell complexes 79
  2.9. Cubical complexes 81

Chapter 3. Combinatorial algebra of face rings 89
  3.1. Face rings of simplicial complexes 90
  3.2. Tor-algebras and Betti numbers 95
  3.3. Cohen–Macaulay complexes 103
  3.4. Gorenstein complexes and Dehn–Sommerville relations 107
  3.5. Face rings of simplicial posets 110

 Face rings: additional topics 117
  3.6. Cohen–Macaulay simplicial posets 117
  3.7. Gorenstein simplicial posets 121
  3.8. Generalised Dehn–Sommerville relations 123

Chapter 4. Moment-angle complexes 125
  4.1. Definitions and functorial properties 127
  4.2. Basic homotopical properties 135

4.3. Cell decomposition 138
4.4. Cohomology ring 138
4.5. Bigraded Betti numbers 144
4.6. Coordinate subspace arrangements 151

Moment-angle complexes: additional topics 157
4.7. Free and almost free torus actions on $\mathcal{Z}_{\mathcal{K}}$ 157
4.8. Massey products in the cohomology of moment-angle complexes 164
4.9. Moment-angle complexes from simplicial posets 167

Chapter 5. Toric varieties and manifolds 173
5.1. Classical construction from rational fans 173
5.2. Projective toric varieties and polytopes 177
5.3. Cohomology of toric manifolds 179
5.4. Algebraic quotient construction 181
5.5. Hamiltonian actions and symplectic reduction 188
5.6. Quotient space by the torus action 192

Chapter 6. Geometric structures on moment-angle manifolds 195
6.1. Intersections of quadrics 195
6.2. Moment-angle manifolds from polytopes 195
6.3. LVM-manifolds 195
6.4. Moment-angle manifolds from complete simplicial fans 195
6.5. Complex geometry of moment-angle manifolds 195
6.6. Dolbeault cohomology and Hodge numbers 195
6.7. Hamiltonian-minimal Lagrangian submanifolds 195
6.8. Moment-angle complexes as Kempf–Ness sets 195

Chapter 7. Half-dimensional torus actions 197
7.1. Basic cohomological theory 197
7.2. Torus manifolds 197
7.3. Quasitoric manifolds 197
7.4. Topological toric manifolds 197
7.5. Bounded flag manifolds 197
7.6. Bott towers 197
7.7. Weight graphs 197

Chapter 8. Homotopy theory of moment-angle complexes 199
8.1. Colimit and homotopy colimit decompositions 199
8.2. Rational homotopy theory 199
8.3. Explicit homotopy types 199
8.4. Stable decompositions of polyhedral products 199
8.5. Loop spaces 199

Chapter 9. Complex cobordism in toric topology 201
9.1. Hirzebruch genera of combinatorial quasitoric data 201
9.2. Toric generators of cobordism ring 201
9.3. Quasitoric representatives in complex cobordism classes 201
9.4. Toric genera 201

Appendix A. Commutative and homological algebra 203

- A.1. Algebras and modules 203
- A.2. Homological theory of graded rings and modules 206
- A.3. Regular sequences and Cohen–Macaulay algebras 212
- A.4. Formality and Massey products 218

Appendix B. Algebraic topology 223
- B.1. Homotopy and homology 223
- B.2. Elements of rational homotopy theory 232
- B.3. Group actions and equivariant cohomology 234
- B.4. Eilenberg–Moore spectral sequences 237

Appendix C. Categorical constructions 239

Appendix D. Regular T-manifolds and their orbit structures 241
- D.1. Locally standard actions, manifolds with faces, and manifolds with corners 241

Appendix E. Stably complex structures and complex bordism 245
- E.1. Bordism of manifolds 245
- E.2. Thom spaces, bordism and cobordism 246
- E.3. Oriented and complex bordism 249
- E.4. Structure results 254
- E.5. Multiplicative generators 255
- E.6. Invariant stably complex structures 258

Appendix F. Formal group laws and Hirzebruch genera 261
- F.1. Elements of the theory of formal group laws 261
- F.2. Formal group law of geometric cobordisms 262
- F.3. Hirzebruch genera 265

Bibliography 269

Introduction

Chapter dependence guide



Acknowledgements. We are most grateful to Anton Ayzenberg, Nikolai Dolbilin, Nickolai Erochovets, Alexander Gaifullin, Peter Landweber, Ivan Limonchenko, Vadim Volodin for the insight gained from discussions on the subject of our book and their most helpful suggestions on improving the text.

Both authors acknowledge the support of the Russian Foundation for Basic Research, grants no. 11-01-00694 and 12-01-92104-ЯФ, grant НШ-4995-2012.1 from the President of Russia, and grant no. 11.G34.31.0053 from the Russian Government. The first author was also supported by a grant from Dmitri Zimin's 'Dynasty' foundation and grant МД-2253.2011.1 from the President of Russia.

CHAPTER 1

Geometry and combinatorics of polytopes

This chapter is an introductory survey of the geometric and combinatorial theory of convex polytopes, with the emphasis on those of its aspects related to the topological applications later in the book. We do not assume any specific knowledge of the reader here. Algebraic definitions (graded rings and algebras) required in the last section of this chapter are contained in the beginning of Section A.1 of the Appendix.

Convex polytopes have been studied since ancient times. Nowadays both combinatorial and geometrical aspects of polytopes are presented in numerous textbooks and monographs. Among them are the classical monograph [**107**] by Grünbaum and Ziegler's more recent lectures [**228**]. Face vectors and other combinatorial questions are discussed in books by McMullen–Shephard [**158**], Brønsted [**32**], and the survey article [**136**] by Klee and Kleinschmidt; while Yemelichev–Kovalev–Kravtsov [**226**] focus on applications to linear programming and optimisation. All these sources may be recommended for the subsequent study of the theory of polytopes, and contain a host of further references.

1.1. Convex polytopes

Definitions and basic constructions. There are two constructively different ways to define a convex polytope in n-dimensional Euclidean affine space $\mathbb{R}^n$ with the scalar product $\langle\ ,\ \rangle$:

Definition 1.1.1. A *convex polytope* is the convex hull $\operatorname{conv}(\boldsymbol{v}_1,\dots,\boldsymbol{v}_q)$ of a finite set of points $\boldsymbol{v}_1,\dots,\boldsymbol{v}_q\in\mathbb{R}^n$.

Definition 1.1.2. A *convex polyhedron* P is an intersection of finitely many half-spaces in some $\mathbb{R}^n$:

$$P=\bigl\{\boldsymbol{x}\in\mathbb{R}^n\colon \langle\boldsymbol{a}_i,\boldsymbol{x}\rangle+b_i\geqslant 0\quad\text{for } i=1,\dots,m\bigr\}, \tag{1.1}$$

where $\boldsymbol{a}_i\in\mathbb{R}^n$ and $b_i\in\mathbb{R}$. A *convex polytope* is a bounded convex polyhedron.

All polytopes in this book will be convex. The two definitions above produce the same geometrical object, i.e. a subset of $\mathbb{R}^n$ is the convex hull of a finite point set if and only if it is a bounded intersection of finitely many half-spaces. This classical fact is proved in many textbooks on polytopes and convex geometry, and it lies at the heart of many applications of polytope theory to linear programming and optimisation, see e.g. [**228**, Theorem 1.1].

The *dimension* of a polytope is the dimension of its affine hull. We usually abbreviate a 'polytope of dimension n' to *n-polytope.* A *supporting hyperplane* of P is an affine hyperplane H which has common points with P and for which the polytope is contained in one of the two closed half-spaces determined by the hyperplane. The intersection $P\cap H$ with a supporting hyperplane is called a *face* of the polytope. We also regard the polytope P itself as a face; other faces are called *proper*. The *boundary* ∂P is the union of all proper faces of P; the complement $\operatorname{int} P = P \setminus \partial P$ is the (relative) *interior* of P. Each face of an n-polytope is itself a polytope of dimension $\leqslant n$. 0-dimensional faces are called *vertices*, 1-dimensional faces are *edges*, and faces of codimension one are *facets*.

Two polytopes $P \subset \mathbb{R}^{n_1}$ and $Q \subset \mathbb{R}^{n_2}$ of the same dimension are said to be *affinely equivalent* (or *affinely isomorphic*) if there is an affine map $\mathbb{R}^{n_1} \to \mathbb{R}^{n_2}$ establishing a bijection between the points of the two polytopes. Two polytopes are *combinatorially equivalent* if there is a bijection between their faces preserving the inclusion relation. Note that two affinely isomorphic polytopes are combinatorially equivalent, but the opposite is not true.

The faces of all dimensions in a given P form a partially ordered set (a *poset*) with respect to inclusion (so that $G \leqslant H$ if the face G is contained in H). This poset is called the *face poset* of P. We therefore obtain that two polytopes are combinatorially equivalent if and only if their face posets are isomorphic.

Definition 1.1.3. A *combinatorial polytope* is a class of combinatorially equivalent polytopes.

Many topological constructions later in this book will depend only on the combinatorial equivalence class of a polytope. Nevertheless, it is always helpful, and sometimes necessary, to keep in mind a particular geometric representative P rather than thinking in terms of abstract posets. Depending on the context, we shall denote by P, Q, etc., geometric polytopes or their combinatorial equivalent classes (combinatorial polytopes). Whenever we consider with both geometric and combinatorial polytopes, we shall use the notation $P \approx Q$ for combinatorial equivalence.

We shall assume our polytopes in $\mathbb{R}^n$ to be of full dimension n, unless otherwise stated. Under this assumption there is no distinction between $\operatorname{int} P$ and the interior of $P \subset \mathbb{R}^n$ in the topological sense.

Let P be given by (1.1). We shall further assume, unless otherwise stated, that there are no redundant inequalities $\langle \boldsymbol{a}_i, \boldsymbol{x} \rangle + b_i \geqslant 0$ in this representation. That is, no inequality can be removed from (1.1) without changing the polytope P. In this case P has exactly m facets, which are given by

$$F_i = \{\boldsymbol{x} \in P \colon \langle \boldsymbol{a}_i, \boldsymbol{x} \rangle + b_i = 0\}$$

for $1 \leqslant i \leqslant m$. The vector $\boldsymbol{a}_i$ is orthogonal to the facet F_i and points towards the interior of the polytope.

Example 1.1.4 (simplex and cube). An n-dimensional *simplex* Δ^n is the convex hull of $n+1$ points in $\mathbb{R}^n$ that do not lie on a common affine hyperplane. All faces of an n-simplex are simplices of dimension $\leqslant n$. Any two n-simplices are affinely equivalent. Let $\boldsymbol{e}_1, \ldots, \boldsymbol{e}_n$ be the standard basis in $\mathbb{R}^n$. The n-simplex $\operatorname{conv}(\boldsymbol{0}, \boldsymbol{e}_1, \ldots, \boldsymbol{e}_n)$ is called *standard*. Equivalently, the standard n-simplex is specified by the $n+1$ inequalities

$$x_i \geqslant 0 \quad \text{for } i = 1, \ldots, n, \quad \text{and} \quad -x_1 - \cdots - x_n + 1 \geqslant 0. \tag{1.2}$$

The *regular* n-simplex is the convex hull of the endpoints of vectors $\boldsymbol{e}_1, \ldots, \boldsymbol{e}_{n+1}$ in $\mathbb{R}^{n+1}$.

The *standard n-cube* is given by

$$\mathbb{I}^n = [0,1]^n = \{(x_1, \ldots, x_n) \in \mathbb{R}^n \colon 0 \leqslant x_i \leqslant 1 \quad \text{for } i = 1, \ldots, n\}. \tag{1.3}$$

Equivalently, the standard n-cube is the convex hull of 2^n points $(\varepsilon_1, \ldots, \varepsilon_n) \in \mathbb{R}^n$, where $\varepsilon_i = 0$ or 1 for $1 \leqslant i \leqslant n$. Whenever we work with combinatorial polytopes, we shall refer to any polytope combinatorially equivalent to $\mathbb{I}^n$ as a *cube*, and denote it by I^n.

The cube $\mathbb{I}^n$ has $2n$ facets. We denote by F_k^0 the facet specified by the equation $x_k = 0$, and by F_k^1 that specified by the equation $x_k = 1$, for $1 \leqslant k \leqslant n$.

The following construction shows that any n-polytope with m facets is affinely equivalent to the intersection of the *positive cone* (or *orthant*)

$$\mathbb{R}^m_{\geqslant} = \bigl\{(y_1, \ldots, y_m) \in \mathbb{R}^m \colon y_i \geqslant 0 \quad \text{for } i = 1, \ldots, m\bigr\} \subset \mathbb{R}^m \tag{1.4}$$

with a certain n-dimensional affine plane.

Construction 1.1.5. Let $P \subset \mathbb{R}^n$ be an n-polytope given by (1.1). Form an $m \times n$-matrix A_P whose rows are the vectors $\boldsymbol{a}_i$ written in the standard basis of $\mathbb{R}^n$, i.e. $(A_P)_{ij} = (\boldsymbol{a}_i)_j$. Note that A_P is of rank n. Likewise, let $\boldsymbol{b}_P = (b_1, \ldots, b_m)^t \in \mathbb{R}^m$ be the column vector of b_i's. Now we can write (1.1) as

$$P = \bigl\{\boldsymbol{x} \in \mathbb{R}^n \colon (A_P\boldsymbol{x} + \boldsymbol{b}_P)_i \geqslant 0 \quad \text{for } i = 1, \ldots, m\bigr\}, \tag{1.5}$$

where $\boldsymbol{x} = (x_1, \ldots, x_n)^t$ is the column of coordinates. Consider the affine map

$$i_P \colon \mathbb{R}^n \to \mathbb{R}^m, \quad i_P(\boldsymbol{x}) = A_P\boldsymbol{x} + \boldsymbol{b}_P. \tag{1.6}$$

It is monomorphic onto a certain n-dimensional plane in $\mathbb{R}^m$, and $i_P(P)$ is the intersection of this plane with the positive cone $\mathbb{R}^m_{\geqslant}$. Let C be an $(m-n) \times m$-matrix whose rows form a basis of linear dependencies between the vectors $\boldsymbol{a}_i$. That is, C is of rank $(m-n)$ and satisfies $CA_P = 0$. Then

$$i_P(P) = \bigl\{\boldsymbol{y} \in \mathbb{R}^m \colon C\boldsymbol{y} = C\boldsymbol{b}_P,\ y_i \geqslant 0 \quad \text{for } i = 1, \ldots, m\bigr\}. \tag{1.7}$$

By definition, the polytopes P and $i_P(P)$ are affinely equivalent.

Example 1.1.6. Consider the standard n-simplex $\Delta^n \subset \mathbb{R}^n$ specified by (1.2). It has $m = n+1$ facets and may be given by (1.1) with $\boldsymbol{a}_1 = (1, 0, \ldots, 0)^t$, …, $\boldsymbol{a}_n = (0, \ldots, 0, 1)^t$, $\boldsymbol{a}_{n+1} = (-1, \ldots, -1)^t$, $b_1 = \cdots = b_n = 0$, $b_{n+1} = 1$. We may take $C = (1, \ldots, 1)$ in Construction 1.1.5. Then $C\boldsymbol{y} = y_1 + \cdots + y_m$, $C\boldsymbol{b}_P = 1$, and we have

$$i_{\Delta^n}(\Delta^n) = \bigl\{\boldsymbol{y} \in \mathbb{R}^{n+1} \colon y_1 + \cdots + y_{n+1} = 1,\ y_i \geqslant 0 \quad \text{for } i = 1, \ldots, n\bigr\}.$$

This is the regular n-simplex in $\mathbb{R}^{n+1}$.

Simple and simplicial polytopes. Polarity. The notion of a *generic* polytope depends on the choice of definition. Below we describe the two possibilities.

A set of $q > n$ points in $\mathbb{R}^n$ is in *general position* if no $n+1$ of them lie on a common affine hyperplane. Now, assuming Definition 1.1.1, we may say that a polytope is generic if it is the convex hull of a set of generally positioned points. This implies that all proper faces of the polytope are simplices, i.e. every facet has the minimal number of vertices (namely, n). Polytopes with this property are called *simplicial*.

On the other hand, a set of $m > n$ hyperplanes $\langle \boldsymbol{a}_i, \boldsymbol{x} \rangle + b_i = 0$, $\boldsymbol{a}_i \in \mathbb{R}^n$, $b_i \in \mathbb{R}$, $1 \leqslant i \leqslant m$, is in *general position* if no point belongs to more than n hyperplanes. From the viewpoint of Definition 1.1.2, a polytope P is generic if its bounding hyperplanes are in general position. This implies that there are exactly n facets meeting at each vertex of P. Such polytopes are called *simple*. Each face of a simple polytope is again a simple polytope. Every vertex of a simple polytope has a neighbourhood affinely equivalent to a neighbourhood of $\mathbf{0}$ in $\mathbb{R}^n_{\geqslant}$. It follows that every vertex is contained in exactly n edges, and each subset of k edges with a common vertex spans a k-face.

For any polytope $P \subset \mathbb{R}^n$ define its *polar set* $P^* \subset \mathbb{R}^n$ as

$$P^* = \bigl\{\boldsymbol{u} \in \mathbb{R}^n \colon \langle \boldsymbol{u}, \boldsymbol{x}\rangle \geqslant -1 \ \text{ for all } \ \boldsymbol{x} \in P\bigr\}. \tag{1.8}$$

Remark. If we do not assume the ambient space to be Euclidean, then the polar set naturally lies in the dual space. Also, our choice of the inequality $\geqslant -1$ in the definition of polarity is not the most standard one (in convex geometry $\leqslant 1$ is usually used instead), but it is better suited for applications in toric geometry. Obviously, these two ways of defining the polar set are taken into each other by the central symmetry. Note also that $\mathbf{0} \in \operatorname{int} P^*$.

The following fact is well known in convex geometry (see [**32**, Theorem 9.1] or [**228**, Theorem 2.11]).

Theorem 1.1.7.

(a) *if a polytope Q is given as a convex hull, $Q = \operatorname{conv}(\boldsymbol{a}_1, \dots, \boldsymbol{a}_m)$, then Q^* is given by inequalities* (1.1) *with $b_i = 1$ for $1 \leqslant i \leqslant m$; in particular, Q^* is a convex polyhedron, but not necessarily bounded;*

(b) *if a polytope P is given by inequalities* (1.1) *with $b_i = 1$, then $P^* = \operatorname{conv}(\mathbf{0}, \boldsymbol{a}_1, \dots, \boldsymbol{a}_m)$;*

(c) *if $\mathbf{0} \in P$ then $(P^*)^* = P$;*

(d) *if $\mathbf{0} \in \operatorname{int} P$, then P^* is bounded, i.e., it is also a polytope;*

(e) *if P^* is a polytope, then its face poset is the opposite to the face poset of P; in particular, the combinatorial type of P^* is determined by that of P.*

Example 1.1.8. The difference between the situations $\mathbf{0} \in P$ and $\mathbf{0} \in \operatorname{int} P$ may be illustrated by the following example. Let $Q = \operatorname{conv}(\mathbf{0}, \boldsymbol{e}_1, \boldsymbol{e}_2)$ be the standard 2-simplex in $\mathbb{R}^2$. Then by Theorem 1.1.7 (a) Q^* is specified by three inequalities

$$\langle \mathbf{0}, \boldsymbol{x}\rangle + 1 \geqslant 0, \quad \langle \boldsymbol{e}_1, \boldsymbol{x}\rangle + 1 \geqslant 0, \quad \langle \boldsymbol{e}_2, \boldsymbol{x}\rangle + 1 \geqslant 0,$$

of which the first is satisfied for all $\boldsymbol{x}$, so we obtain an unbounded polyhedron. Its dual is $\operatorname{conv}(\mathbf{0}, \boldsymbol{e}_1, \boldsymbol{e}_2)$ by Theorem 1.1.7 (b), giving back the standard 2-simplex.

The condition $0 \in \operatorname{int} P$ can be achieved by a parallel translation, without changing the affine or combinatorial type of P. Having this in mind, we refer to P^* as the *polar polytope* of P. It follows from Theorem 1.1.7 (e) that if P is simple then P^* is simplicial, and vice versa. Any polygon is both simple and simplicial.

Proposition 1.1.9. *In dimensions $n \geqslant 3$ a simplex is the only polytope which is both simple and simplicial.*

Proof. Let be P be a polytope which is both simple and simplicial. Choose a vertex $v \in P$. Since P is simple, v is connected by edges to exactly n other vertices, say $v_1, \dots, v_n$. We claim that there are no other vertices in P. To prove this it is enough to show that all vertices $v_1, \dots, v_n$ are pairwise connected by edges, as then every vertex from $v, v_1, \dots, v_n$ will be connected to the remaining n vertices. Indeed, take a pair v_i, v_j. Since P is simple and v is connected to both v_i and v_j, all these three vertices belong to a 2-face. Since P is simplicial and $n \geqslant 3$, this face is a 2-simplex, in which v_i and v_j are connected by an edge. We conclude that P has $n+1$ vertices, so it is an n-simplex. $\square$

The proof above also shows that if all 2-faces of a simple polytope are triangular, then P is a simplex. A similar property is valid for a cube: if all 2-faces of a simple polytope P quadrangular, then P is a cube (an exercise).

EXAMPLE 1.1.10. The polar of a simplex is again a simplex. To describe the polar of a cube, we consider the cube $[-1,1]^n$ (the standard cube (1.3) is not good as 0 is not in its interior). Then the polar set is the convex hull of the endpoints of $2n$ vectors $\pm\boldsymbol{e}_k,\ 1\leqslant k\leqslant n$. It is called the *cross-polytope*. The 3-dimensional cross-polytope is the octahedron.

A combinatorial polytope P is called *self-dual* if P^* is combinatorially equivalent to P. There are many examples of self-dual non-simple polytopes; an infinite family of them is given by k-gonal pyramids for $k\geqslant 4$. Here is a more interesting *regular* example:

EXAMPLE 1.1.11 (24-cell). Let Q be the 4-polytope obtained by taking the convex hull of the following 24 points in $\mathbb{R}^4$: endpoints of 8 vectors $\pm\boldsymbol{e}_i,\ 1\leqslant i\leqslant 4$, and 16 points of the form $(\pm\frac12,\pm\frac12,\pm\frac12,\pm\frac12)$. By Theorem 1.1.7 (a), the polar Q^* is given by the following 24 inequalities:

$$\pm x_i+1\geqslant 0 \ \text{ for } i=1,\dots,4, \quad \text{and } \ \tfrac12(\pm x_1\pm x_2\pm x_3\pm x_4)+1\geqslant 0. \tag{1.9}$$

Each of these inequalities turns into equality in exactly one of the specified 24 points, so it defines a supporting hyperplane whose intersection with Q is only one point. This implies that Q has exactly 24 vertices. The vertices of Q^* may be determined by applying the 'elimination process' to (1.9) (see [**228**, §1.2]), and as a result we obtain 24 points of the form $\pm\boldsymbol{e}_i\pm\boldsymbol{e}_j$ for $1\leqslant i<j\leqslant 4$. Each supporting hyperplane defined by (1.9) contains exactly 6 vertices of Q^*, which form an octahedron. So both Q and Q^* have 24-vertices and 24 octahedral facets. In fact, both Q and Q^* provide examples of a *regular* 4-*polytope* called a 24-*cell*. It is the only regular self-dual polytope different from a simplex. For more details on the 24-cell and other regular polytopes see [**67**].

Basic methods of producing new polytopes from given ones.

CONSTRUCTION 1.1.12 (Product). The *product* $P_1\times P_2$ of two simple polytopes P_1 and P_2 is again a simple polytope. The dual operation on simplicial polytopes can be described as follows. Let $S_1\subset\mathbb{R}^{n_1}$ and $S_2\subset\mathbb{R}^{n_2}$ be two simplicial polytopes. Assume that both S_1 and S_2 contain 0 in their interiors. Now define

$$S_1\circ S_2=\operatorname{conv}\bigl(S_1\times 0\cup 0\times S_2\bigr)\subset\mathbb{R}^{n_1+n_2}.$$

Then $S_1\circ S_2$ is again a simplicial polytope. For any two simple polytopes P_1, P_2 containing 0 in their interiors the following identity holds:

$$P_1^*\circ P_2^*=(P_1\times P_2)^*.$$

Both operations $\times$ and $\circ$ are also defined on combinatorial polytopes; in this case the above formula holds without any restrictions.

CONSTRUCTION 1.1.13 (Hyperplane cuts and face truncations). Assume given a simple polytope (1.1) and a hyperplane $H=\{\boldsymbol{x}\in\mathbb{R}^n\colon\langle\boldsymbol{a},\boldsymbol{x}\rangle+b=0\}$ that does not contain any vertex of P. Then the intersections $P\cap H_\geqslant$ and $P\cap H_\leqslant$ of P with either of the halfspaces determined by H are simple polytopes; we refer to them as *hyperplane cuts* of P. To see that $P\cap H_\geqslant$ and $P\cap H_\leqslant$ are simple we note that their new vertices are transverse intersections of H with the edges of P. Since P

is simple, each of those edges is contained in $n-1$ facets, so each new vertex of $P \cap H_{\geqslant}$ or $P \cap H_{\leqslant}$ is contained in exactly n facets.

If H separates all vertices of a certain i-face $G \subset P$ from the other vertices of P and $G \subset H_{\geqslant}$, then $P \cap H_{\geqslant}$ is combinatorially equivalent to $G \times \Delta^{n-i}$ (an exercise), and we say that the polytope $P \cap H_{\leqslant}$ is obtained from P by a *face truncation*.

In particular, if G is a vertex, the result is a *vertex truncation* of P. When the choice of the truncated vertex is clear or irrelevant we use the notation $\mathrm{vt}(P)$. We also use the notation $\mathrm{vt}^k(P)$ for a polytope obtained from P by a k-fold iteration of the vertex truncation.

We may describe the face poset of the simple polytope $\widetilde{P}$ obtained from P by truncation of a face $G \subset P$ as follows. Let $F_1, \ldots, F_m$ be the facets of P, and assume that $G = F_{i_1} \cap \cdots \cap F_{i_k}$. The polytope $\widetilde{P}$ has m facets corresponding to $F_1, \ldots, F_m$ (and obtained from them by truncation), which we denote by the same letters for simplicity, and a new facet $F = P \cap H$. Then we have

$$\begin{aligned}
&F_{j_1} \cap \cdots \cap F_{j_\ell} \neq \varnothing \text{ in } \widetilde{P}\\
&\qquad\qquad \Longleftrightarrow F_{j_1} \cap \cdots \cap F_{j_\ell} \neq \varnothing \text{ in } P, \text{ and } F_{j_1} \cap \cdots \cap F_{j_\ell} \not\subset G,\\
&F \cap F_{j_1} \cap \cdots \cap F_{j_\ell} \neq \varnothing \text{ in } \widetilde{P}\\
&\qquad\qquad \Longleftrightarrow G \cap F_{j_1} \cap \cdots \cap F_{j_\ell} \neq \varnothing \text{ in } P, \text{ and } F_{j_1} \cap \cdots \cap F_{j_\ell} \not\subset G.
\end{aligned}$$

Note that $F_{j_1} \cap \cdots \cap F_{j_\ell} \not\subset G$ if and only if $\{i_1, \ldots, i_k\} \not\subset \{j_1, \ldots, j_\ell\}$.

The two previous construction worked for geometric polytopes. Here is an example of a construction which is more suitable for combinatorial ones.

CONSTRUCTION 1.1.14 (Connected sum of polytopes). Suppose we are given two simple polytopes P and Q, both of dimension n, with distinguished vertices v and w respectively. An informal way to obtain the *connected sum* $P \#_{v,w} Q$ of P at v and Q at w is as follows. We cut off v from P and w from Q; then, after a projective transformation, we can glue the rest of P to the rest of Q along the new simplex facets to obtain $P \#_{v,w} Q$. A more formal definition is given below, following [**52**, §6].

First, we introduce an n-dimensional polyhedron Γ, which will be used as a template for the construction; it arises by considering the standard $(n-1)$-simplex Δ^{n-1} in the subspace $\{\boldsymbol{x}\colon x_1 = 0\} \subset \mathbb{R}^n$, and taking its cartesian product with the first coordinate axis. The facets G_r of Γ therefore have the form $\mathbb{R} \times D_r$, where D_r, $1 \le r \le n$, are the facets of Δ^{n-1}. Both Γ and the G_r are divided into positive and negative halves, determined by the sign of the coordinate x_1.

We order the facets of P meeting in v as $E_1, \ldots, E_n$, and the facets of Q meeting in w as $F_1, \ldots, F_n$. Denote the complementary sets of facets by $\mathcal{C}_v$ and $\mathcal{C}_w$; those in $\mathcal{C}_v$ avoid v, and those in $\mathcal{C}_w$ avoid w.

We now choose projective transformations φ_P and φ_Q of $\mathbb{R}^n$, whose purpose is to map v and w to the infinity of the x_1 axis. We insist that φ_P embeds P in Γ so as to satisfy two conditions; firstly, that the hyperplane defining E_r is identified with the hyperplane defining G_r, for each $1 \le r \le n$, and secondly, that the images of the hyperplanes defining $\mathcal{C}_v$ meet Γ in its negative half. Similarly, φ_Q identifies the hyperplane defining F_r with that defining G_r, for each $1 \le r \le n$, but the images of the hyperplanes defining $\mathcal{C}_w$ meet Γ in its positive half. We define the *connected sum* $P \#_{v,w} Q$ of P at v and Q at w to be the simple n-polytope determined by the images of the hyperplanes defining $\mathcal{C}_v$ and $\mathcal{C}_w$ and hyperplanes defining G_r,

$1 \leqslant r \leqslant n$. It is defined only up to combinatorial equivalence; moreover, different choices for either of v and w, or either of the orderings for E_r and F_r, are likely to affect the combinatorial type. When the choices are clear, or their effect on the result irrelevant, we use the abbreviation $P \mathbin{\#} Q$.

The related construction of connected sum $P \mathbin{\#} S$ of a simple polytope P and a simplicial polytope S is described in [**228**, Example 8.41].

Example 1.1.15.
1. If P is an m_1-gon and Q is an m_2-gon then $P \mathbin{\#} Q$ is an $(m_1 + m_2 - 2)$-gon.
2. If P is an n-simplex, then $P \mathbin{\#} Q \approx \mathrm{vt}(P)$.
3. If both P and Q are n-simplices, then $P \mathbin{\#} Q \approx \mathrm{vt}(\Delta^n) \approx \Delta^{n-1} \times \Delta^1$. The combinatorial type of $\mathrm{vt}(\Delta^n)$ does not depend on the choice of the truncated vertex. All the vertices of the resulting polytope $\Delta^{n-1} \times \Delta^1$ are equivalent, therefore, the combinatorial type of $\mathrm{vt}^2(\Delta^n) \approx \Delta^n \mathbin{\#} \Delta^n \mathbin{\#} \Delta^n$ is still independent of the choices. The choice of the truncated vertex becomes significant from the next step, i.e. for $\mathrm{vt}^3(\Delta^n)$, see Exercise 1.1.25.

Neighbourly polytopes.

Definition 1.1.16. A polytope is called *k-neighbourly* if any set of its k or fewer vertices spans a face. According to Exercise 1.1.26, the only n-polytope which is more than $\left[\frac{n}{2}\right]$-neighbourly is a simplex. An n-polytope which is $\left[\frac{n}{2}\right]$-neighbourly is called simply *neighbourly.*

Example 1.1.17 (neighbourly 4-polytope). Let $P = \Delta^2 \times \Delta^2$, the product of two triangles. Then P is simple, and it is easy to see that any two facets of P share a common 2-face. Therefore, any two vertices of P^* are connected by an edge, so P^* is a neighbourly simplicial 4-polytope.

More generally, if P_1^* is k_1-neighbourly and P_2^* is k_2-neighbourly, then $(P_1 \times P_2)^*$ is $\min(k_1, k_2)$-neighbourly. It follows that $(\Delta^n \times \Delta^n)^*$ and $(\Delta^n \times \Delta^{n+1})^*$ are neighbourly $2n$- and $(2n+1)$-polytopes respectively. The next example gives a neighbourly polytope with an arbitrary number of vertices.

Example 1.1.18 (cyclic polytopes). The *moment curve* in $\mathbb{R}^n$ is given by

$$\boldsymbol{x}\colon \mathbb{R} \longrightarrow \mathbb{R}^n, \qquad t \mapsto \boldsymbol{x}(t) = (t, t^2, \ldots, t^n) \in \mathbb{R}^n.$$

For any $m > n$ define the *cyclic polytope* $C^n(t_1, \ldots, t_m)$ as the convex hull of m distinct points $\boldsymbol{x}(t_i)$, $t_1 < t_2 < \ldots < t_m$, on the moment curve.

Theorem 1.1.19.

(a) *$C^n(t_1, \ldots, t_m)$ is a simplicial n-polytope;*
(b) *$C^n(t_1, \ldots, t_m)$ has exactly m vertices $\boldsymbol{x}(t_i)$, $1 \leqslant i \leqslant m$;*
(c) *the combinatorial type of $C^n(t_1, \ldots, t_m)$ does not depend on the specific choice of parameters $t_1, \ldots, t_m$;*
(d) *$C^n(t_1, \ldots, t_m)$ is a neighbourly polytope.*

Proof. This proof is taken from [**228**, Theorem 0.7]. Recall the well-known Vandermonde determinant identity

$$\det\begin{pmatrix} 1 & 1 & \cdots & 1 \\ \boldsymbol{x}(t_0) & \boldsymbol{x}(t_1) & \cdots & \boldsymbol{x}(t_n) \end{pmatrix} = \begin{vmatrix} 1 & 1 & \cdots & 1 \\ t_0 & t_1 & \cdots & t_n \\ \vdots & \vdots & \ddots & \vdots \\ t_0^{n-1} & t_1^{n-1} & \cdots & t_n^{n-1} \\ t_0^n & t_1^n & \cdots & t_n^n \end{vmatrix} = \prod_{0\leqslant i<j\leqslant n} (t_j - t_i).$$

This implies that no $n+1$ points on the moment curve belong to a common affine hyperplane, proving (a). Denote $[m] = \{1, \ldots, m\}$. Properties (b) and (c) follow from the following statement: an n-element subset $\omega \subset [m]$ corresponds to the vertex set of a facet of $C^n(t_1, \ldots, t_m)$ if and only if the following 'Gale's evenness condition' is satisfied:

> *If elements $i < j$ are not in ω, then the number of elements $k \in \omega$ between i and j is even.*

To prove this we write $\omega = \{i_1, \ldots, i_n\}$ and consider the hyperplane H_ω through the corresponding points $\boldsymbol{x}(t_{i_s})$, $1 \leqslant s \leqslant n$, on the moment curve. We have

$$H_\omega = \{\boldsymbol{x} \in \mathbb{R}^n \colon F_\omega(\boldsymbol{x}) = 0\},$$

where

$$F_\omega(\boldsymbol{x}) = \det\begin{pmatrix} 1 & 1 & \ldots & 1 \\ \boldsymbol{x} & \boldsymbol{x}(t_{i_1}) & \ldots & \boldsymbol{x}(t_{i_n}) \end{pmatrix}.$$

(The latter is exactly the linear function vanishing on the prescribed points.) Now let the point $\boldsymbol{x}(t)$ move on the moment curve. Then $F_\omega(\boldsymbol{x}(t))$ is a polynomial in t of degree n. It has n different roots $t_{i_1}, \ldots, t_{i_n}$, and changes sign at each of them. Now ω corresponds to the vertex set of a facet if and only if $F_\omega(\boldsymbol{x}(t_i))$ has the same sign for all the points $\boldsymbol{x}(t_i)$ with $i \notin \omega$; that is, if $F_\omega(\boldsymbol{x}(t))$ has an even number of sign changes between $t = t_i$ and $t = t_j$, for $i < j$ and $i, j \notin \omega$. This proves Gale's condition, and statements (b) and (c).

It remains to prove (d). We need to check that any subset $\tau = \{i_1, \ldots, i_k\} \subset [m]$ of cardinality $k \leqslant \left[\frac{n}{2}\right]$ corresponds to the vertex set of a face. Choose some $\varepsilon > 0$ so that $t_i < t_i + \varepsilon < t_{i+1}$ for all $i < m$, and some $N > t_m + \varepsilon$. Define a linear function $F_\tau(\boldsymbol{x})$ as

$$\det\bigl(\boldsymbol{x}, \boldsymbol{x}(t_{i_1}), \boldsymbol{x}(t_{i_1} + \varepsilon), \ldots, \boldsymbol{x}(t_{i_k}), \boldsymbol{x}(t_{i_k} + \varepsilon), \boldsymbol{x}(N+1), \ldots, \boldsymbol{x}(N+n-2k)\bigr).$$

It vanishes on $\boldsymbol{x}(t_i)$ with $i \in \tau$. Now $F_\tau(\boldsymbol{x}(t))$ is a polynomial in t of degree n, and it has n different roots

$$t_{i_1}, t_{i_1} + \varepsilon, \ldots, t_{i_k}, t_{i_k} + \varepsilon, N+1, \ldots, N+n-2k.$$

If $i, j \notin \tau$, then there is an even number of roots between $t = t_i$ and $t = t_j$, because a root $t = t_l$ always come in a pair with the root $t = t_l + \varepsilon$. Thus, the linear function $F_\tau(\boldsymbol{x})$ has the same sign on all the points $\boldsymbol{x}(t_i)$ with $i \notin \tau$. This linear function defines a supporting hyperplane, so τ corresponds to the vertex set of a face. □

We shall denote the combinatorial cyclic n-polytope with m vertices by $C^n(m)$.

The vertices and edges of a polytope P determine a graph, which is called the *graph of the polytope* and denoted $\Gamma(P)$. This graph is *simple*, that is, it has no loops and multiple edges. The following theorem is due to Blind and Mani, see also [**228**, §3.4] for a simpler proof given by Kalai.

Theorem 1.1.20. *The combinatorial type of a simple polytope P is determined by its graph $\Gamma(P)$. In other words, two simple polytopes are combinatorially equivalent if their graphs are isomorphic.*

This theorem fails for general polytopes: the graph of a neighbourly polytope is isomorphic to that of a simplex with the same number of vertices. In general, simplicial n-polytopes are determined by their $[\frac{n}{2}]$-skeleta. General n-polytopes are determined by their $(n-2)$-skeleta. See [**228**, §3.4] for more history and references.

Exercises.

1.1.21. Show that if P and Q are n-polytopes, and the face poset of P is a subposet of the face poset of Q, then P and Q are combinatorially equivalent.

1.1.22. Show that a simple n-polytope all of whose 2-faces are quadrangular is combinatorially equivalent to an n-cube.

1.1.23. Show that any hyperplane cut of Δ^n is combinatorially equivalent to a product of two simplices. Conclude that any combinatorial simple n-polytope with $n+2$ facets is combinatorially equivalent (in fact, projectively equivalent) to a product of two simplices.

1.1.24. Let P be a simple polytope. Show that if a hyperplane H separates all vertices of a certain i-face $G \subset P$ from the other vertices of P and $G \subset H_{\geqslant}$, then $P \cap H_{\geqslant} \approx G \times \Delta^{n-i}$.

1.1.25. How many combinatorially different polytopes may be obtained as $\mathrm{vt}^3(\Delta^n)$?

1.1.26. Show that if a polytope is k-neighbourly, then every $(2k-1)$-face is a simplex. Conclude that if an n-polytope is $([\frac{n}{2}]+1)$-neighbourly, then it is a simplex. Conclude also that a neighbourly $2k$-polytope is simplicial. Are there neighbourly non-simplicial polytopes of odd dimension?

1.1.27. Are the polytopes $(\Delta^n \times \Delta^n)^*$ and $(\Delta^n \times \Delta^{n+1})^*$ combinatorially equivalent to cyclic polytopes?

1.2. Gale diagrams

1.3. Face vectors and Dehn–Sommerville relations

The notion of the f-vector (or face vector) is a central concept in the combinatorial theory of polytopes. It has been studied since the time of Euler.

Definition 1.3.1. Let P be a convex n-polytope. Denote by f_i the number of i-dimensional faces of P. The integer sequence $\boldsymbol{f}(P) = (f_0, f_1, \ldots, f_n)$ is known as the *f-vector* (or the *face vector*) of P. Note that $f_n = 1$. The homogeneous *F-polynomial* of P is defined by

$$F(P)(s,t) = s^n + f_{n-1}s^{n-1}t + \cdots + f_1 st^{n-1} + f_0 t^n.$$

The *h-vector* $\boldsymbol{h}(P) = (h_0, h_1, \ldots, h_n)$ and the *H-polynomial* of P are defined by

$$\begin{aligned} h_0 s^n + h_1 s^{n-1}t + \cdots + h_n t^n &= (s-t)^n + f_{n-1}(s-t)^{n-1}t + \cdots + f_0 t^n, \\ H(P)(s,t) = h_0 s^n + h_1 s^{n-1}t + \cdots + h_{n-1}st^{n-1} + h_n t^n &= F(P)(s-t,t). \end{aligned} \tag{1.10}$$

The *g-vector* of a simple polytope P is the vector $\boldsymbol{g}(P) = (g_0, g_1, \ldots, g_{[\frac{n}{2}]})$, where $g_0 = 1$ and $g_i = h_i - h_{i-1}$ for $i = 1, \ldots, [n/2]$.

EXAMPLE 1.3.2. We have

$$F(\Delta^n) = s^n + \binom{n+1}{1} s^{n-1}t + \binom{n+1}{2} s^{n-2}t^2 + \cdots + t^n = \frac{(s+t)^{n+1} - t^{n+1}}{s},$$

$$H(\Delta^n) = s^n + s^{n-1}t + s^{n-2}t^2 + \cdots + t^n = \frac{s^{n+1} - t^{n+1}}{s-t}.$$

Obviously, the f-vector is a *combinatorial invariant* of a polytope, that is, it depends only on the face poset. This invariant is far from being complete, even for simple polytopes:

EXAMPLE 1.3.3. Two different combinatorial simple polytopes may have same f-vectors. For instance, let P be the 3-cube and Q a simple 3-polytope with 2 triangular, 2 quadrangular and 2 pentagonal facets, see Figure 1.1. (Note that Q is obtained by truncating two vertices of a tetrahedron, it is also dual to the cyclic polytope $C^3(6)$ from Definition 1.1.18.) Then $\boldsymbol{f}(P) = \boldsymbol{f}(Q) = (8, 12, 6)$.



FIGURE 1.1. Two combinatorially non-equivalent simple polytopes with the same f-vectors.

The f-vector and the h-vector contain equivalent combinatorial information, and determine each other by means of linear relations, namely

$$h_k = \sum_{i=0}^{k} (-1)^{k-i} \binom{n-i}{n-k} f_{n-i}, \quad f_k = \sum_{q=k}^{n} \binom{q}{k} h_{n-q}, \quad \text{for } 0 \leqslant k \leqslant n. \tag{1.11}$$

In particular, $h_0 = 1$ and $h_n = f_0 - f_1 + \cdots + (-1)^n f_n$. By the *Euler formula*,

$$f_0 - f_1 + \cdots + (-1)^n f_n = 1, \tag{1.12}$$

which is equivalent to $h_n = h_0$. This is the first evidence of the fact that many combinatorial relations for the face numbers have much simpler form when written in terms of the h-vector. Another example of this phenomenon is given by the following generalisation of the Euler formula for simple or simplicial polytopes.

THEOREM 1.3.4 (Dehn–Sommerville relations). *The h-vector of any simple n-polytope is symmetric, that is,*

$$H(s,t) = H(t,s), \quad \textit{or} \quad h_i = h_{n-i} \quad \textit{for} \ \ 0 \leqslant i \leqslant n.$$

The Dehn–Sommerville relations can be proved in many different ways. We present a proof from [**32**], which can be viewed as a combinatorial version of Morse-theoretic arguments. An alternative proof will be given in Section 1.7.

Proof of Theorem 1.3.4. Let $P \subset \mathbb{R}^n$ be a simple polytope. Choose a generic linear function $\varphi\colon \mathbb{R}^n \to \mathbb{R}$ which distinguishes the vertices of P. Write $\varphi(\boldsymbol{x}) = \langle \boldsymbol{\nu}, \boldsymbol{x}\rangle$ for some vector $\boldsymbol{\nu}$ in $\mathbb{R}^n$. The assumption on φ implies that $\boldsymbol{\nu}$ is parallel to no edge of P. We can view φ as a height function on P and turn the 1-skeleton of P into a directed graph by orienting each edge in such a way that φ increases along it, see Figure 1.2. For each vertex v of P define the index $\mathrm{ind}_\nu(v)$ as the number of incident edges that point towards v. Denote the number of vertices of index i by $I_\nu(i)$. We claim that $I_\nu(i) = h_{n-i}$. Indeed, each face of P has a unique top vertex (the maximum of the height function φ restricted to the face) and a unique bottom vertex (the minimum of φ). Let G be a k-face of P, and v_G its top vertex. Since P is simple, there are exactly k edges of G meeting at v_G, whence $\mathrm{ind}(v_G) \geqslant k$. On the other hand, each vertex of index $q \geqslant k$ is the top vertex for exactly $\binom{q}{k}$ faces of dimension k. It follows that the number of k-faces of P can be calculated as



Figure 1.2. Orienting the 1-skeleton of P.

$$f_k = \sum_{q \geqslant k} \binom{q}{k} I_\nu(q).$$

Now the second identity from (1.11) shows that $I_\nu(q) = h_{n-q}$, as claimed. In particular, the number $I_\nu(q)$ does not depend on $\boldsymbol{\nu}$. At the same time, we have $\mathrm{ind}_\nu(v) = n - \mathrm{ind}_{-\nu}(v)$ for any vertex v, which implies that

$$h_{n-q} = I_\nu(q) = I_{-\nu}(n-q) = h_q. \qquad \square$$

Remark. The above proof also shows that the numbers $h_k = I_\nu(n-k)$ are nonnegative, which is not evident from (1.11). On the other hand, the nonnegativity of the h-vector translates into certain conditions on the f-vector. This will be important in the subsequent study of f-vectors for combinatorial objects more general than simple polytopes.

Theorem 1.3.5. *The f-vector of a simple n-polytope satisfies*

$$f_k = \sum_{i=0}^{k} (-1)^i \binom{n-i}{n-k} f_i, \quad \text{for } 0 \leqslant k \leqslant n. \tag{1.13}$$

Proof. By the Dehn–Sommerville relations,

$$F(s-t,t)=H(s,t)=H(t,s)=F(t-s,s).$$

By substituting $u=s-t$ we obtain $F(u,t)=F(-u,t+u)$, or

$$\begin{aligned}&u^n+f_{n-1}u^{n-1}t+\dots+f_1ut^{n-1}+f_0u_n\\&\qquad=(-u)^n+f_{n-1}(-u)^{n-1}(t+u)+\dots+f_1(-u)(t+u)^{n-1}+f_0(t+u)^n.\end{aligned}$$

Calculating the coefficient of u^kt^{n-k} in both sides above yields (1.13). □

It follows from Theorem 1.1.7 (e) that the f-vector of the polar n-polytope P^* satisfies

$$f_i(P^*)=f_{n-1-i}(P),\quad \text{for } 0\leqslant i\leqslant n-1.$$

Then it follows from (1.13) that the f-vector of a simplicial polytope satisfies the relations $f_{n-1-k}=\sum_{i=0}^{k}(-1)^i\binom{n-i}{n-k}f_{n-1-i}$ or, equivalently,

$$f_{q-1}=\sum_{j=q}^{n}(-1)^{n-j}\binom{j}{q}f_{j-1},\quad \text{for } 1\leqslant q\leqslant n+1.$$

Proposition 1.3.6. *The F- and H-polynomial are multiplicative, i.e.*

$$F(P_1\times P_2)=F(P_1)F(P_2),\quad H(P_1\times P_2)=H(P_1)H(P_2). \tag{1.14}$$

for any convex polytopes P_1 and P_2.

Proof. Let $\dim P_1=n_1$ and $\dim P_2=n_2$. Each k-face of $P_1\times P_2$ is the product of an i-face of P_1 and a $(k-i)$-face of P_2 for some i, whence

$$f_k(P_1\times P_2)=\sum_{i=0}^{n_1}f_i(P_1)f_{k-i}(P_2),\quad \text{for } 0\leqslant k\leqslant n_1+n_2. \tag{1.15}$$

This implies the first identity, and the second follows from (1.10). □

Example 1.3.7. We have $F(I^n)=(F(\Delta^1))^n$ and $H(I^n)=(H(\Delta^1))^n$, i.e.

$$F(I^n)=(s+2t)^n,\quad H(I^n)=(s+t)^n.$$

We can also express the f-vector and the h-vector of the connected sum $P\mathbin{\#}Q$ in terms of those of P and Q (the proof is left as an exercise):

Proposition 1.3.8. *Let P and Q be simple n-polytopes. Then*

$$\begin{aligned}&f_0(P\mathbin{\#}Q)=f_0(P)+f_0(Q)-2;\quad h_0(P\mathbin{\#}Q)=h_n(P\mathbin{\#}Q)=1;\\&f_i(P\mathbin{\#}Q)=f_i(P)+f_i(Q)-\binom{n}{i},\quad \textit{for } 1\leqslant i\leqslant n;\\&h_i(P\mathbin{\#}Q)=h_i(P)+h_i(Q),\quad \textit{for } 1\leqslant i\leqslant n-1.\end{aligned}$$

Using the Dehn–Sommerville relations we can show that a simplicial polytope cannot be ‘too neighbourly’ (see Definition 1.1.16) if it is not a simplex.

Proposition 1.3.9. *Let S be a q-neighbourly simplicial n-polytope, and let $S\neq\Delta^n$. Then $q\leqslant\left[\frac{n}{2}\right]$.*

Proof. Let S^* be the polar simple polytope. By Theorem 1.1.7 (e), $f_{n-i}(S^*) = f_{i-1}(S) = \binom{m}{i}$ for $1 \leqslant i \leqslant q$, where m is the number of vertices of S. From (1.11) we get

$$h_k(S^*) = \sum_{i=0}^{k}(-1)^{k-i}\binom{n-i}{k-i}\binom{m}{i} = \binom{m-n+k-1}{k}, \quad \text{for } k \leqslant q, \tag{1.16}$$

The second equality is obtained by calculating the coefficient of t^k on both sides of

$$\frac{1}{(1+t)^{n-k+1}}(1+t)^m = (1+t)^{m-n+k-1}.$$

If $S \neq \Delta^n$, then $m > n+1$, which together with (1.16) gives $h_0(S^*) < h_1(S^*) < \cdots < h_q(S^*)$. It then follows from the Dehn–Sommerville relations that $q \leqslant \left[\frac{n}{2}\right]$. $\square$

Since the H-polynomial of a simple n-polytope P satisfies the identity $H(P)(s,t) = H(P)(t,s)$, we can express it in terms of elementary symmetric functions as follows:

$$H(P)(s,t) = \sum_{i=0}^{[n/2]} \gamma_i (s+t)^{n-2i}(st)^i. \tag{1.17}$$

The identity $h_0 = h_n = 1$ implies that $\gamma_0 = 1$.

Definition 1.3.10. The integer sequence $\gamma(P) = (\gamma_0, \ldots, \gamma_{[n/2]})$ is called the *γ-vector* of P. We refer to

$$\gamma(P)(\tau) = \gamma_0 + \gamma_1\tau + \cdots + \gamma_{[n/2]}\tau^{[n/2]}$$

as the *γ-polynomial* of P.

Example 1.3.11. We have $\gamma(I^1)(\tau) = 1$. If P^2_m is an m-gon, then $\gamma(P^2_m)(\tau) = 1 + (m-4)\tau$.

The components of the γ-vector can be expressed via the components of any of the f-, h- or g-vector by means of linear relations, and vice versa. The explicit transition formulae between the g- and γ-vectors are given by the next lemma.

Lemma 1.3.12. *Let P be a simple n-polytope. Then*

$$g_i = (n-2i+1)\sum_{j=0}^{i}\frac{1}{n-i-j+1}\binom{n-2j}{i-j}\gamma_j;$$

$$\gamma_i = (-1)^i\sum_{j=0}^{i}(-1)^j\binom{n-i-j}{i-j}g_j, \qquad 0 \leqslant i \leqslant \left[\tfrac{n}{2}\right].$$

Proof. Using the formulae of Examples 1.3.2 and 1.3.7 we calculate

$$H(P) = \sum_{j=0}^{[n/2]}\gamma_j(st)^j H(I^{n-2j}),$$

$$H(I^q) = \sum_{j=0}^{q}\binom{q}{j}s^j t^{q-j} = \sum_{k=0}^{[q/2]}c_{k,q}(st)^k H(\Delta^{q-2k}),$$

where $c_{0,q}=1$ and $c_{k,q}=\binom{q}{k}-\binom{q}{k-1}=\frac{q-2k+1}{q-k+1}\binom{q}{k}>0$ for $k=1,\ldots,[\frac{q}{2}]$. Hence

$$H(P)(s,t)=\sum_{i=0}^{[n/2]}\Bigl(\sum_{j=0}^{i}c_{i-j,n-2j}\gamma_j\Bigr)(st)^iH(\Delta^{n-2i}).$$

On the other hand,

$$H(P)(s,t)=\sum_{i=0}^{[n/2]}g_i(st)^iH(\Delta^{n-2i}).$$

Comparing the last two formulae we obtain $g_i=\sum_{j=0}^{i}c_{i-j,n-2j}\gamma_j$, which is equivalent to the first required formula. The second is left as an exercise. $\square$

Proposition 1.3.13. *Let P,Q be two simple n-polytopes, and consider the following four conditions:*

(a) $f_i(P)\geqslant f_i(Q)$ *for* $i=0,1,\ldots,n$*;*
(b) $h_i(P)\geqslant h_i(Q)$ *for* $i=0,1,\ldots,n$*;*
(c) $g_i(P)\geqslant g_i(Q)$ *for* $i=0,1,\ldots,\bigl[\frac{n}{2}\bigr]$*;*
(d) $\gamma_i(P)\geqslant \gamma_i(Q)$ *for* $i=0,1,\ldots,\bigl[\frac{n}{2}\bigr]$.

Then (d)⇒(c)⇒(b)⇒(a).

Proof. The first of equations (1.11) implies that the components of $\boldsymbol{f}(P)$ are expressed via the components of $\boldsymbol{h}(P)$ with positive coefficients, which proves the implication (b)⇒(a). We have $h_k=\sum_{i=0}^{k}g_i$ for $0\leqslant k\leqslant\bigl[\frac{n}{2}\bigr]$, which gives the implication (c)⇒(b). The implication (d)⇒(c) follows similarly from the first formula of Lemma 1.3.12. $\square$

The components of the f-vector of any polytope are nonnegative. The nonnegativity of the components of the h-vector of a simple polytope follows from their geometric interpretation obtained in the proof of Theorem 1.3.4; the h-vector of a non-simple polytope may have negative components (e.g. for the octahedron). The nonnegativity of the g-vector of a simple n-polytope (that is, the inequalities $g_i(P)\geqslant g_i(\Delta^n)$) is a much more subtle property; it follows from the g-theorem discussed in the next section. The γ-vector of a simple polytope may have negative components (e.g. for $P=\Delta^2$); its nonnegativity for special classes of simple polytopes will be discussed in Section 1.6. The nonnegativity of the γ-vector can be expressed by the inequalities $\gamma_i(P)\geqslant\gamma_i(I^n)$, see Exercise 1.3.17.

Exercises.

1.3.14. Show that any 3-polytope has a face with $\leqslant 5$ vertices.

1.3.15. Prove Proposition 1.3.8.

1.3.16. Prove the second transition formula of Lemma 1.3.12.

1.3.17. Show that the γ-polynomial is multiplicative, that is,

$$\gamma(P\times Q)(\tau)=\gamma(P)(\tau)\cdot\gamma(Q)(\tau)$$

In particular, $\gamma(I^n)(\tau)=1$, i.e. $\gamma(I^n)=(1,0,\ldots,0)$.

1.4. Characterising the face vectors of polytopes

The face numbers are the simplest combinatorial invariants of polytopes, and they arise in many hard problems of combinatorial geometry. One of the most natural and basic questions is to describe all possible face numbers, or, more precisely, determine which integer vectors arise as the f-vectors of polytopes. In the general case this questions is probably intractable (see the end of this section), but a particularly nice answer exists in the case of simple (or, equivalently, simplicial) polytopes. Obviously, the Dehn–Sommerville relations provide a necessary condition. As far as only linear equations are concerned, there are no further restrictions:

Proposition 1.4.1 (Klee [**135**]). *The Dehn–Sommerville relations are the most general linear equations satisfied by the face numbers of all simple polytopes.*

Proof. Once again, the use of the h-vector simplifies the proof significantly. Given a simple polytope P, we set

$$h(P)(t) = H(P)(1,t) = h_0(P) + h_1(P)t + \cdots + h_n(P)t^n.$$

It is enough to prove that the affine hull of the h-vectors $(h_0, h_1, \ldots, h_n)$ of simple n-polytopes is an $\left[\frac{n}{2}\right]$-dimensional plane. This can be done by presenting $\left[\frac{n}{2}\right]+1$ simple polytopes with affinely independent h-vectors. Take $Q_k = \Delta^k \times \Delta^{n-k}$ for $0 \leqslant k \leqslant \left[\frac{n}{2}\right]$. Since $h(\Delta^k)(t) = 1 + t + \cdots + t^k$, formula (1.14) gives

$$h(Q_k)(t) = \frac{1-t^{k+1}}{1-t} \cdot \frac{1-t^{n-k+1}}{1-t}.$$

It follows that the lowest degree term in the polynomial $h(Q_{k+1})(t) - h(Q_k)(t)$ is t^{k+1}, for $0 \leqslant k \leqslant \left[\frac{n}{2}\right]-1$. Therefore, the vectors $\boldsymbol{h}(Q_k)$ are affinely independent for these values of k. □

Example 1.4.2. Let P be a simple polytope. Since every vertex is contained in exactly n edges and each edge connects two vertices, we have a linear relation

$$2f_1 = nf_0. \tag{1.18}$$

By Proposition 1.4.1 this must be a consequence of the Dehn–Sommerville relations (in fact, it is equation (1.13) for $k = 1$.)

Equation (1.18) together with the Euler identity (1.12) shows that the f-vector of a simple 3-polytope P^3 is completely determined by the number of facets, namely,

$$\boldsymbol{f}(P^3) = (2f_2 - 4, 3f_2 - 6, f_2, 1).$$

Similarly, the f-vector of a simplicial 3-polytope S^3 is determined by the number of vertices, namely,

$$\boldsymbol{f}(S^3) = (f_0, 3f_0 - 6, 2f_0 - 4, 1).$$

Remark. Euler's formula (1.12) is the only linear relation satisfied by the face vectors of general polytopes. This can be proved similarly to Proposition 1.4.1, by specifying sufficiently many polytopes with affinely independent face vectors.

Apart from the linear equations, the f-vectors of polytopes satisfy certain inequalities. Here are some of the simplest of them.

Example 1.4.3. There are the following obvious lower bounds for the number of vertices and the number of facets of an n-polytope:

$$f_0 \geqslant n+1, \quad f_{n-1} \geqslant n+1.$$

Since every pair of vertices is joined by at most one edge, and every pair of facets intersect at most one face of codimension 2, we have the upper bounds

$$f_1 \leqslant \binom{f_0}{2}, \quad f_{n-2} \leqslant \binom{f_{n-1}}{2}.$$

If the polytope is simplicial, then there is also the following lower bound for f_1:

$$f_1 \geqslant nf_0 - \binom{n+1}{2}.$$

It is much more difficult to prove though, even for 4-polytopes. For simplicial 3-polytopes the inequality above turns into identity.

Historically, the most important inequality-type results preceding the general characterisation of f-vectors were the *Upper Bound Theorem (UBT)* and the *Lower Bound Theorem (LBT)*. They give respectively an upper and a lower bound for the number of faces in a simplicial polytope with the given number of vertices.

THEOREM 1.4.4 (UBT for simplicial polytopes). *Among all simplicial n-polytopes S with m vertices the cyclic polytope $C^n(m)$ (Example 1.1.18) has the maximal number of i-faces for $1 \leqslant i \leqslant n$. That is, if $f_0(S) = m$, then*

$$f_i(S) \leqslant f_i\bigl(C^n(m)\bigr) \quad \text{for } i = 1, \ldots, n.$$

Equality is achieved for all i if and only if S is a neighbourly polytope.

The UBT was conjectured by Motzkin and proved by McMullen [**154**] in 1970. Since $C^n(m)$ is neighbourly, we have

$$f_i\bigl(C^n(m)\bigr) = \binom{m}{i+1} \quad \text{for } 0 \leqslant i \leqslant \left[\tfrac{n}{2}\right] - 1.$$

Due to the Dehn–Sommerville relations, this determines the full f-vector of $C^n(m)$. The exact values are given by the following lemma.

LEMMA 1.4.5. *The number of i-faces of the cyclic polytope $C^n(m)$ (or any neighbourly n-polytope with m vertices) is given by*

$$f_i = \sum_{q=0}^{\left[\frac{n}{2}\right]} \binom{q}{n-1-i}\binom{m-n+q-1}{q} + \sum_{p=0}^{\left[\frac{n-1}{2}\right]} \binom{n-p}{n-1-i}\binom{m-n+p-1}{p}$$

for $0 \leqslant i \leqslant n$, where we set $\binom{p}{q} = 0$ for $p < q$ or $q < 0$.

PROOF. We set $C = C^n(m)$. Using the second identity from (1.11), the identity $\left[\frac{n}{2}\right] + 1 = n - \left[\frac{n-1}{2}\right]$, the Dehn–Sommerville relations for C^*, and (1.16), we calculate

$$\begin{aligned} f_i(C) = f_{n-1-i}(C^*) &= \sum_{q=0}^{n} \tbinom{q}{n-1-i} h_{n-q}(C^*) \\ &= \sum_{q=0}^{[n/2]} \tbinom{q}{n-1-i} h_q(C^*) + \sum_{q=[n/2]+1}^{n} \tbinom{q}{n-1-i} h_{n-q}(C^*) \\ &= \sum_{q=0}^{[n/2]} \tbinom{q}{n-1-i}\tbinom{m-n+q-1}{q} + \sum_{p=0}^{[(n-1)/2]} \tbinom{n-p}{n-1-i}\tbinom{m-n+p-1}{p}. \end{aligned}$$

□

Lemma 1.4.6. *Assume that the inequalities*

$$h_i(P) \leqslant \binom{m-n+i-1}{i}, \qquad i = 0, \dots, \left[\tfrac{n}{2}\right]$$

hold for the h*-vector of a simple polytope* P *with* m *facets. Then the dual simplicial polytope* P^* *satisfies the UBT inequalities* $f_i(P^*) \leqslant f_i\bigl(C^n(m)\bigr)$ *for* $i = 1, \dots, n$*.*

Proof. Since $h_i\bigl((C^n(m))^*\bigr) = \binom{m-n+i-1}{i}$ for $i = 0, \dots, \left[\tfrac{n}{2}\right]$, the statement follows from Proposition 1.3.13. Alternatively, replace the last '=' in the calculation from the proof of Lemma 1.4.5 by '$\leqslant$'. □

This was one of the key observations in McMullen's original proof [**154**] of the UBT for simplicial polytopes (the whole proof can also be found in [**32**, §18] and [**228**, §8.4]). R. Stanley gave an algebraic argument establishing the inequalities from Lemma 1.4.6, and therefore the UBT, in a much more general setting of *triangulated spheres*. We shall discuss Stanley's approach and conclude the proof of the UBT in Section 3.3.

Remark. The UBT holds for all convex polytopes. That is, the cyclic polytope $C^n(m)$ has the maximal number of i-faces from all convex n-polytopes with m vertices. The argument for this builds on the following observation of Klee and McMullen, which we reproduce from [**228**, Lemma 8.24].

Lemma 1.4.7. *By a small perturbation of vertices of an* n*-polytope* P *one can achieve that the resulting polytope* P' *is simplicial, and*

$$f_i(P) \leqslant f_i(P') \quad \textit{for } i = 1, \dots, n-1.$$

Definition 1.4.8. A simplicial n-polytope S is called *stacked* if there is a sequence $S_0, S_1, \dots, S_k = S$ of n-polytopes such that S_0 is an n-simplex and S_{i+1} is obtained from S_i by adding a pyramid over a facet of S_i (the vertex of the added pyramid is chosen close enough to its base, so that the whole construction remains convex and simplicial). The polar simple polytopes are those obtained from a simplex by iterating the vertex cut operation of Example 1.1.15.2. These are sometimes called *truncation polytopes*.

The f-vector of a stacked polytope is easy to calculate (see Exercise 1.4.17).

Theorem 1.4.9 (LBT for simplicial polytopes). *Among all simplicial* n*-polytopes* S *with* m *vertices a stacked polytope has the minimal number of* i*-faces for* $2 \leqslant i \leqslant n-1$*. That is, if* $f_0(S) = m$*, then*

$$\begin{aligned} f_i(S) &\geqslant \tbinom{n}{i}m - \tbinom{n+1}{i+1}i \quad \textit{for } i = 1, \dots, n-2; \\ f_{n-1}(S) &\geqslant (n-1)m - (n+1)(n-2). \end{aligned}$$

For $n \neq 3$ *equality is achieved for all* i *if and only if* S *is a stacked polytope.*

Remark. For $n = 3$ the LBT inequalities $f_1 \geqslant 3m - 6$ and $f_2 \geqslant 2m - 4$ turn into equalities for all simplicial polytopes.

An inductive argument by McMullen, Perles and Walkup [**159**] reduced the LBT to the case $i = 1$, namely, to the inequality $f_1 \geqslant nm - \binom{n+1}{2}$. It was finally proved by Barnette [**15**], [**16**]. Barnette's proof of the LBT, with some simplifications, can also be found in [**32**]. The fact that equality is achieved only for stacked polytopes (if $n \neq 3$) was proved by Billera and Lee [**23**].

Remark. Unlike the UBT, little is know about generalisations of the LBT to non-simplicial convex polytopes. Some results in this direction were obtained in [**130**] along with generalisations of the LBT to triangulated spheres and manifolds, which we also discuss later in this book.

An easy calculation shows that the inequalities $f_{n-1} \geqslant n+1$ and and $f_{n-2} \geqslant nf_{n-1} - \binom{n+1}{2}$ for simple polytopes from Example 1.4.3 can be written in terms of the h-vector as follows: $h_0 \leqslant h_1 \leqslant h_2$. Having reduced the whole LBT to the inequality $h_1 \leqslant h_2$, McMullen and Walkup [**159**] conjectured that the components of the h-vector 'grow up to the middle', that is the inequalities

$$h_0 \leqslant h_1 \leqslant \cdots \leqslant h_{\left[\frac{n}{2}\right]} \tag{1.19}$$

hold for a simple n-polytope. It has since become known as the *Generalised Lower Bound Conjecture (GLBC)*.

McMullen also suggested a generalisation to the UBT, whose formulation requires an algebraic digression.

Definition 1.4.10. For any two positive integers a, i there exists a unique *binomial i-expansion* of a of the form

$$a = \binom{a_i}{i} + \binom{a_{i-1}}{i-1} + \cdots + \binom{a_j}{j},$$

where $a_i > a_{i-1} > \cdots > a_j \geqslant j \geqslant 1$.

The binomial i-expansion of a can be constructed by choosing a_i as the unique number satisfying $\binom{a_i}{i} \leqslant a < \binom{a_i+1}{i}$, then choosing a_{i-1}, and so on. One needs only to check that $a_i > a_{i-1}$, which is straightforward.

Now define the ith *pseudopower* of a as

$$a^{\langle i\rangle} = \binom{a_i+1}{i+1} + \binom{a_{i-1}+1}{i} + \cdots + \binom{a_j+1}{j+1}, \quad 0^{\langle i\rangle} = 0.$$

Example 1.4.11.

1. For $a > 0$, $a^{\langle 1\rangle} = \binom{a+1}{2}$.

2. If $i \geqslant a$ then the binomial expansion has the form

$$a = \binom{i}{i} + \binom{i-1}{i-1} + \cdots + \binom{i-a+1}{i-a+1} = 1 + \cdots + 1,$$

and therefore $a^{\langle i\rangle} = a$.

3. Let $a = 28$, $i = 4$. Then

$$28 = \binom{6}{4} + \binom{5}{3} + \binom{3}{2}, \quad 28^{\langle 4\rangle} = \binom{7}{5} + \binom{6}{4} + \binom{4}{3} = 40.$$

The importance of the binomial expansion and pseudopowers comes from the following fundamental result of combinatorial commutative algebra.

Theorem 1.4.12 (Macaulay, Stanley). *The following two conditions are equivalent for a sequence of integers $(k_0, k_1, k_2, \ldots)$:*

(a) $k_0 = 1$ *and* $0 \leqslant k_{i+1} \leqslant k_i^{\langle i\rangle}$ *for* $i \geqslant 1$*;*

(b) *there exists a connected commutative graded algebra* $A = A^0 \oplus A^1 \oplus A^2 \oplus \cdots$ *over a field* $\mathbf{k}$ *such that* A *is generated by its degree-one elements and* $\dim_{\mathbf{k}} A^i = k_i$ *for* $i \geqslant 0$.

Macaulay's original theorem [**144**] says that (b) above is equivalent to the existence of a *multicomplex* whose h-vector is given by $(k_0, k_1, k_2, \ldots)$. The original proof is long and complicated. The reformulation of Macaulay's condition in terms of pseudopowers, i.e. condition (a), is due to Stanley [**208**, Theorem 2.2]. Simpler proofs of Theorem 1.4.12 can be found in [**62**] and [**35**, §4.2].

Definition 1.4.13. A sequence of integers satisfying either of the conditions of Theorem 1.4.12 is called an *M-sequence*. Finite M-sequences are *M-vectors*.

Now observe that the number $\binom{m-n+i-1}{i}$ on the right hand side of the inequality of Lemma 1.4.6 equals the number of degree i monomials in $h_1 = m-n$ generators. It follows that the inequalities of Lemma 1.4.6, and therefore the UBT, hold if the h-vector is an M-vector.

In 1970 McMullen [**155**] combined all known and conjectured information about the f-vectors, including the Dehn–Sommerville relations and the generalisations to the LBT and UBT discussed above, into a (conjectured) complete characterisation. McMullen's conjecture is now proved, and remains up to the present time perhaps the most impressive achievement of the combinatorial theory of face numbers:

Theorem 1.4.14 (g-theorem). *An integer vector* $(f_0, f_1, \ldots, f_n)$ *is the* f*-vector of a simple* n*-polytope if and only if the corresponding sequence* $(h_0, \ldots, h_n)$ *determined by* (1.10) *satisfies the following three conditions:*

(a) $h_i = h_{n-i}$ *for* $i = 0, 1, \ldots, n$ *(the Dehn–Sommerville relations);*
(b) $h_0 \leqslant h_1 \leqslant \cdots \leqslant h_{\left[\frac{n}{2}\right]}$;
(c) $h_0 = 1$, $h_{i+1} - h_i \leqslant (h_i - h_{i-1})^{\langle i \rangle}$ *for* $i = 1, \ldots, \left[\frac{n}{2}\right] - 1$.

Remark. Condition (b) says that the components of the h-vector 'grow up to the middle', while (c) gives a restriction on the rate of this growth. Both (b) and (c) can be reformulated by saying that the g-vector $(g_0, \ldots, g_{[n/2]})$ (see Definition 1.3.1) is an M-vector; this explains the name 'g-theorem'. The fact that the g-vector is an M-vector implies that the h-vector is also an M-vector (see Exercise 1.4.18).

Both necessity and sufficiency parts of the g-theorem were proved almost simultaneously (around 1980), although by radically different methods.

The sufficiency part was proved by Billera and Lee [**22**], [**23**]. The proof is quite elementary and relies upon a remarkable combinatorial-geometrical construction combining cyclic polytopes (achieving the upper bound for the number of faces) with the operation of 'adding a pyramid' (used to produce polytopes achieving the lower bound). As a result, a simplicial polytope can be produced with any prescribed g-vector between the minimal and the maximal ones. Another important consequence of the results of [**159**] and [**23**] is that the GLBC inequalities (1.19) are the most general linear inequalities satisfied by the f-vectors of simplicial polytopes.

On the other hand, Stanley's proof [**204**] of the necessity part of g-theorem (i.e. that the g-vector of a simple polytope is an M-vector) used deep results from algebraic geometry, in particular, the *Hard Lefschetz theorem* for the cohomology of *toric varieties*. We shall give this argument in Section 5.3. After the appearance of Stanley's paper combinatorists had been looking for a more elementary combinatorial proof of his theorem, until in 1993 a first such proof was found by McMullen [**156**]. It builds up on the notion of the *polytope algebra*, which may be thought of as a combinatorial model for the cohomology algebras of toric varieties. Despite being elementary, it was a complicated proof. Later McMullen simplified his approach in [**157**]. Yet another elementary proof of the g-theorem was given by Timorin [**218**]. It relies on an interpretation of McMullen's polytope algebra as the algebra of differential operators with constant coefficients vanishing on the *volume polynomial* of the polytope.

By duality, the UBT and the LBT provide upper and lower bounds for the number of faces of a simple polytope with the given number of facets. Similarly, the g-theorem also provides a characterisation for the f-vectors of simplicial polytopes. During the last three decades some work was done in extending the g-theorem to objects more general than simplicial (or simple) polytopes, although the most important conjecture here remains open since 1971 (see Section 2.5). There are basically two diverging routes for generalisations of the g-theorem: towards non-polytopal objects (like triangulations of spheres or manifolds), and towards general convex polytopes which are neither simple nor simplicial. The former requires machinery from combinatorial topology and commutative algebra; we shall discuss the corresponding generalisations in more detail in the next chapters. The generalisations of the g-theorem to non-simplicial convex polytopes are mostly beyond the scope of this book; they require algebraic geometry techniques such as *intersection homology*, which we only briefly discuss in Section 5.3.

Exercises.

1.4.15. Prove the following upper bound for the number of k-faces in a simple n-polytope P with f_0 vertices:

$$f_k \leqslant \frac{1}{k+1}\binom{n}{k} f_0, \quad \text{for } k = 1, \ldots, n,$$

where the equality is achieved only for $P = \Delta^n$. Observe that this inequality gives a better upper bound for simple polytopes that the UBT.

1.4.16. Prove Lemma 1.4.7.

1.4.17. Show that the numbers of faces of a stacked n-polytope S with $m = f_0$ vertices are given by

$$\begin{aligned} f_i(S) &= \tbinom{n}{i} m - \tbinom{n+1}{i+1} i \quad \text{for } 1 \leqslant i \leqslant n-2; \\ f_{n-1}(S) &= (n-1)m - (n+1)(n-2). \end{aligned}$$

1.4.18. Let $\boldsymbol{h} = (h_0, h_1, \ldots, h_n)$ be an integer vector with $h_0 = 1$ and $h_i = h_{n-i}$ for $0 \leqslant i \leqslant n$, and let $\boldsymbol{g} = (g_0, g_1, \ldots, g_{[n/2]})$ where $g_0 = 1$, $g_i = h_i - h_{i-1}$ for $i > 0$. Show that if $\boldsymbol{g}$ is an M-vector, then $\boldsymbol{h}$ is also an M-vector.

1.4.19. Prove directly that parts (a) and (b) of the g-theorem imply the LBT, while parts (a) and (c) imply the UBT.

Polytopes: additional topics

1.5. Nestohedra and graph-associahedra

Several constructions of series of simple polytopes with remarkable properties appeared in the beginning of the 1990s under the common name of 'generalised associahedra'. (The original associahedron, or *Stasheff polytope*, was first introduced in homotopy theory [**210**].) Nowadays generalised associahedra find numerous applications in algebraic geometry [**98**], the theory of knot and link invariants [**29**], representation theory and cluster algebras [**92**], and the theory of operads and geometric 'field theories' originating from quantum physics [**211**].

Without attempting to overview all aspects of generalised associahedra, we describe one particular construction which uses the Minkowski sum and the combinatorial concept of a *building set*. Our exposition is much influenced by the original works of Feichtner–Sturmfels [**88**] and Postnikov [**187**]. The resulting family of polytopes is known as *nestohedra*. Although it does not include all possible generalisations of associahedra, this family is wide enough to contain all classical series, and its construction is elementary enough so that it requires no specific knowledge.

The *Minkowski sum* is a classical geometric construction allowing one to produce new polytopes from known ones, just like the product, hyperplane cut and connected sum described in Section 1.1. However, a Minkowski sum of simple polytopes usually fails to be simple. Interesting examples of polytopes can be obtained by taking Minkowski sums of regular simplices. Simplices in such a Minkowski sum are indexed by a collection $\mathcal{S}$ of subsets in a finite set. It was shown in [**88**] and [**187**] that a Minkowski sum of regular simplices is a simple polytope if $\mathcal{S}$ satisfies certain combinatorial condition, identifying it as a *building set*. The resulting family of simple polytopes was called *nestohedra* in [**188**] because of its connection to *nested sets* considered by De Concini and Procesi [**71**] in the context of subspace arrangements. An example of a building set is provided by the collection of subsets of vertices which span connected subgraphs in a given graph. The corresponding nestohedra are called *graph-associahedra*; they were introduced and studied in the work of Carr and Devadoss [**57**], and independently in the work of Toledano Laredo [**219**] under the name *De Concini–Procesi associahedra*. These include the classical series of *permutahedra* and *associahedra*.

Minkowski sums of simplices. Recall that the *Minkowski sum* of two subsets $A, B \subset \mathbb{R}^n$ is defined as

$$A + B = \{\boldsymbol{x} + \boldsymbol{y} \colon \boldsymbol{x} \in A,\ \boldsymbol{y} \in B\}.$$

Proposition 1.5.1. *The Minkowski sum of two polytopes is a polytope. Moreover, if* $P = \operatorname{conv}(\boldsymbol{v}_1, \dots, \boldsymbol{v}_k)$ *and* $Q = \operatorname{conv}(\boldsymbol{w}_1, \dots, \boldsymbol{w}_l)$*, then*

$$P + Q = \operatorname{conv}(\boldsymbol{v}_1 + \boldsymbol{w}_1, \dots, \boldsymbol{v}_i + \boldsymbol{w}_j, \dots, \boldsymbol{v}_k + \boldsymbol{w}_l).$$

PROOF. Follows directly from the definition of Minkowski sum. ☐

For every subset $S \subset [n+1] = \{1, \dots, n+1\}$ consider the regular simplex

$$\Delta_S = \operatorname{conv}(\boldsymbol{e}_i \colon i \in S) \subset \mathbb{R}^{n+1}.$$

Let $\mathcal{F}$ be a collection of nonempty subsets of $[n+1]$. We assume that $\mathcal{F}$ contains all singletons $\{i\},\ 1 \leqslant i \leqslant n+1$. As usual, denote by $|\mathcal{F}|$ the number of elements in $\mathcal{F}$. Given a subset $N \subset [n+1]$, denote by $\mathcal{F}|_N$ the *restriction* of $\mathcal{F}$ to N, i.e.

$$\mathcal{F}|_N = \{S \in \mathcal{F} \colon S \subset N\}$$

of subsets in N. A collection $\mathcal{F}$ is *connected* if $[n+1]$ cannot be represented as a disjoint union of nonempty subsets N_1 and N_2 such that for every $S \in \mathcal{F}$ either $S \subset N_1$ or $S \subset N_2$. Obviously, every collection $\mathcal{F}$ splits into the disjoint union of its *connected components*.

REMARK. In some further considerations we shall allow $\mathcal{F}$ to contain some subsets of $[n+1]$ with multiplicities.

Now consider the convex polytope

$$P_{\mathcal{F}} = \sum_{S \in \mathcal{F}} \Delta_S \subset \mathbb{R}^{n+1}. \tag{1.20}$$

The following statement gives some of its basic properties.

PROPOSITION 1.5.2. *Let $\mathcal{F} = \mathcal{F}_1 \sqcup \dots \sqcup \mathcal{F}_q$ be the decomposition into connected components. Then*

(a) $P_{\mathcal{F}} = P_{\mathcal{F}_1} \times \dots \times P_{\mathcal{F}_q}$;
(b) $\dim P_{\mathcal{F}} = n + 1 - q$.

PROOF. Statement (a) follows from the fact that the polytopes $P_{\mathcal{F}_i}$ are contained in complementary subspaces for $1 \leqslant i \leqslant q$, so their Minkowski sum is the product. Because of (a), it is enough to verify (b) for connected $\mathcal{F}$ only. Then we need to prove that $\dim P_{\mathcal{F}} = n$. Observe that $P_{\mathcal{F}}$ is contained in the hyperplane $H_{\mathcal{F}} = \bigl\{\boldsymbol{x} \in \mathbb{R}^{n+1} \colon \sum_{i=1}^{n+1} x_i = |\mathcal{F}|\bigr\}$, and therefore $\dim P_{\mathcal{F}} \leqslant n$. If $\mathcal{F}$ has a unique maximal element, then this element is $[n+1]$, because $\mathcal{F}$ is connected. Hence, $P_{\mathcal{F}}$ has an n-simplex as a Minkowski summand, and therefore $\dim P_{\mathcal{F}} = n$. We shall only need this case in the further considerations, so we skip the rest of the proof and leave it as an exercise. ☐

We now describe two extreme examples of polytopes $P_{\mathcal{F}}$, corresponding to the minimal and the maximal connected collections.

EXAMPLE 1.5.3 (Simplex). Let $\mathcal{S}$ be the collection consisting of all singletons and the whole set $[n+1]$. Then $P_{\mathcal{S}}$ is the regular n-simplex $\Delta_{[n+1]}$ shifted by the vector $\boldsymbol{e}_1 + \dots + \boldsymbol{e}_{n+1}$.

Permutahedron. Let $\mathcal{C}$ be the *complete* collection, consisting of all subsets in $[n+1]$. The polytope $P_{\mathcal{C}}$ is n-dimensional by Proposition 1.5.2.

THEOREM 1.5.4. *$P_{\mathcal{C}}$ can be described as the intersection of the hyperplane*

$$H_{\mathcal{C}} = \Bigl\{\boldsymbol{x} \in \mathbb{R}^{n+1} \colon \sum_{i=1}^{n+1} x_i = 2^{n+1} - 1\Bigr\}$$

with the halfspaces

$$H_{S,\geqslant} = \big\{\boldsymbol{x} \in \mathbb{R}^{n+1} \colon \sum_{i\in S} x_i \geqslant 2^{|S|} - 1\big\}$$

for all proper subsets $S \subsetneq [n+1]$*. Moreover, every halfspace above is irredundant, i.e. determines a facet* F_S *of* $P_{\mathcal{C}}$*, so there are* $|\mathcal{C}| = 2^{n+1} - 2$ *facets in total.*

PROOF. By definition, every point $\boldsymbol{x} = (x_1, \dots, x_{n+1}) \in P_{\mathcal{C}}$ can be written as $\boldsymbol{x} = \sum_{S\in\mathcal{C}} \boldsymbol{x}^S$ where $\boldsymbol{x}^S = (x_1^S, \dots, x_{n+1}^S) \in \Delta_S$. Then

$$\sum_{i=1}^{n+1} x_i = \sum_{S\in\mathcal{C}} \sum_{i=1}^{n+1} x_i^S = \sum_{S\in\mathcal{C}} 1 = |\mathcal{C}| = 2^{n+1} - 1,$$

which implies that $P_{\mathcal{C}} \subset H_{\mathcal{C}}$. Similarly,

$$\sum_{i\in S} x_i = \sum_{T\in\mathcal{C}} \sum_{i\in S} x_i^T \geqslant \sum_{T\subset S} \sum_{i\in S} x_i^T = \sum_{T\subset S} 1 = \big|\mathcal{C}|_S\big| = 2^{|S|} - 1, \tag{1.21}$$

so $P_{\mathcal{C}}$ is contained in all subspaces $H_{S,\geqslant}$.

It remains to show that any facet of $P_{\mathcal{C}}$ has the form $P_{\mathcal{C}} \cap H_S$, where H_S is the bounding hyperplane for $H_{S,\geqslant}$. Since $P_{\mathcal{C}}$ is a Minkowski sum of simplices, each of its faces G is a Minkowski sum of faces of these simplices. We therefore may write G as $\sum_{S\in\mathcal{C}} \Delta_{T_S}$ where $T_S \subset S$. By Proposition 1.5.2, if G is a facet (i.e. $\dim G = n-1$), then the collection $\mathcal{T} = \{T_S\}$ of subsets in $[n+1]$ has exactly two connected components. (Note that $\mathcal{T}$ may contain some subsets more than once.) Let $[n+1] = N_1 \sqcup N_2$ and $\mathcal{T} = \mathcal{T}|_{N_1} \sqcup \mathcal{T}|_{N_2}$ be the decomposition into components. Then the hyperplane containing the facet G is defined by each of the two equations

$$\sum_{i\in N_1} x_i = \big|\mathcal{T}|_{N_1}\big| \quad \text{or} \quad \sum_{i\in N_2} x_i = \big|\mathcal{T}|_{N_2}\big|. \tag{1.22}$$

Since every T_S is contained in the corresponding S, we have

$$\big|\mathcal{T}|_{N_1}\big| \geqslant \big|\mathcal{C}|_{N_1}\big| \quad \text{and} \quad \big|\mathcal{T}|_{N_2}\big| \geqslant \big|\mathcal{C}|_{N_2}\big|. \tag{1.23}$$

We claim that at least one of these inequalities turns into equality. Indeed, assume the converse. By (1.21) the minimum of the linear function $\sum_{i\in N_1} x_i$ on $P_{\mathcal{C}}$ is $\big|\mathcal{C}|_{N_1}\big|$, so there is a point $\boldsymbol{x}' \in P_{\mathcal{C}}$ with

$$\sum_{i\in N_1} x_i' = \big|\mathcal{C}|_{N_1}\big| < \big|\mathcal{T}|_{N_1}\big|. \tag{1.24}$$

Similarly, there is a point $\boldsymbol{x}'' \in P_{\mathcal{C}}$ with

$$\sum_{i\in N_2} x_i'' = \big|\mathcal{C}|_{N_2}\big| < \big|\mathcal{T}|_{N_2}\big|.$$

Since $N_1 \sqcup N_2 = [n+1]$, the latter inequality is equivalent to $\sum_{i\in N_1} x_i'' > \big|\mathcal{T}|_{N_1}\big|$. This together with (1.24) implies that there are points of $P_{\mathcal{C}}$ in both open halfspaces determined by the first of the equations (1.22), which contradicts the assumption that G is a facet. So at least one of (1.23) is an equality, which implies that the hyperplane (1.22) containing G has the form H_S (where S is either N_1 or N_2).

It follows that every facet is contained in the hyperplane H_S for some S. On the other hand, every subset S can be taken as N_1 in the construction of the preceding paragraph, which shows that every H_S contains a facet. □

Having identified the facets, we may derive the following description of the whole face poset of $P_{\mathcal{C}}$.

PROPOSITION 1.5.5. *Faces of $P_{\mathcal{C}}$ of dimension k are in one-to-one correspondence with ordered partitions of the set $[n+1]$ into $n+1-k$ nonempty parts. An inclusion of faces $G \subset F$ occurs whenever the ordered partition corresponding to G can be obtained by refining the ordered partition corresponding to F.*

PROOF. This follows from the fact that two facets F_{S_1} and F_{S_2} of $P_{\mathcal{C}}$ have nonempty intersection if and only if $S_1 \subset S_2$ or $S_2 \subset S_1$. We skip the details to avoid repetitive arguments; see Theorem 1.5.13 below for a more general result. $\square$

COROLLARY 1.5.6.

(a) *$P_{\mathcal{C}}$ is a simple polytope;*
(b) *the vertices of $P_{\mathcal{C}}$ are obtained by all permutations of the coordinates of the point $(1, 2, 4, \ldots, 2^n) \in \mathbb{R}^{n+1}$.*

The polytope whose vertices are obtained by permuting the coordinates of a given point is known as the *permutahedron*; it has been studied by convex geometers since the beginning of the 20th century. More precisely, given a point $\boldsymbol{a} = (a_1, \ldots, a_{n+1}) \in \mathbb{R}^{n+1}$ with $a_1 < a_2 < \cdots < a_{n+1}$, define the corresponding permutahedron as

$$Pe^n(\boldsymbol{a}) = \operatorname{conv}\bigl(\sigma(a_1), \ldots, \sigma(a_{n+1}) \colon \sigma \in \Sigma_{n+1}\bigr),$$

where Σ_{n+1} denotes the group of permutations of $n+1$ elements. In particular, $P_{\mathcal{C}} = Pe^n(1, 2, \ldots, 2^n)$. All n-dimensional permutahedra $Pe^n(\boldsymbol{a})$ are combinatorially equivalent; this follows from the following description of their faces:

THEOREM 1.5.7.

(a) *Every facet of $Pe^n(\boldsymbol{a})$ is the intersection of $Pe^n(\boldsymbol{a})$ with the hyperplane*

$$H_S = \bigl\{\boldsymbol{x} \in \mathbb{R}^{n+1} \colon \sum_{i \in S} x_i = a_1 + a_2 + \cdots + a_{|S|}\bigr\}$$

for a proper subset $S \subsetneq [n+1]$.

(b) *The faces of $Pe^n(\boldsymbol{a})$ are described in the same way as in Proposition 1.5.5.*

PROOF. This can be proved by mimicking the proof of Theorem 1.5.4. Another way to proceed is as follows. Every face of $Pe^n(\boldsymbol{a})$ is a set of points where a certain linear function $\varphi_{\boldsymbol{b}} = (\boldsymbol{b}, \cdot\,)$ restricted to the polytope achieves its minimum. We denote this face by $G_{\boldsymbol{b}}$. Then $\boldsymbol{b} = (b_1, \ldots, b_{n+1})$ defines an ordered partition $[n+1] = N_1 \sqcup \cdots \sqcup N_k$ according to the sets of equal coordinates of $\boldsymbol{b}$. Namely, if $b_k = b_l$ then k and l are in the same N_i, while if $b_k < b_l$ then $k \in N_i$ and $l \in N_j$ with $i < j$. It can be shown that (a) $\dim G_{\boldsymbol{b}} = n+1-k$, and (b) the face $G_{\boldsymbol{b}}$ only depends on the ordered partition above and does not depend on the particular values of b_k. In particular, the vertices of $Pe^n(\boldsymbol{a})$ correspond to $\varphi_{\boldsymbol{b}}$ where all coordinates of $\boldsymbol{b}$ are different, while the facets correspond to $\varphi_{\boldsymbol{b}}$ where all coordinates of $\boldsymbol{b}$ are either 0 or 1. We leave the details to the reader. $\square$

We shall denote an n-dimensional combinatorial permutahedron by Pe^n. The classical permutahedron corresponds to $\boldsymbol{a} = (1, 2, \ldots, n+1)$. It can also be obtained as a Minkowski sum (1.20) as follows (we leave the proof as an exercise):

Proposition 1.5.8. *Let $\mathcal{I}$ be the collection of all subsets of cardinality $\leqslant 2$ in $[n+1]$. Then*

$$P_{\mathcal{I}} = Pe^n(1, 2, \ldots, n+1).$$

The polytope $P_{\mathcal{I}}$ is the Minkowski sum of segments of the form $\operatorname{conv}(\boldsymbol{e}_i, \boldsymbol{e}_j)$ for $1 \leqslant i < j \leqslant n+1$, shifted by the vector $\boldsymbol{e}_1 + \cdots + \boldsymbol{e}_{n+1}$. Minkowski sums of segments are known as *zonotopes*; this family of polytopes has many remarkable properties [**228**, §7.3]. However, general zonotopes are rarely simple; the permutahedron is one of the few exceptions.

Building sets and nestohedra. We now return to general Minkowski sums (1.20). The next statement gives a description of $P_{\mathcal{F}}$ in terms of inequalities, and generalises the first part of Theorem 1.5.4.

Proposition 1.5.9 ([**88**, Proposition 3.12]). *$P_{\mathcal{F}}$ can be described as the intersection of the hyperplane*

$$H_{\mathcal{F}} = \Bigl\{\boldsymbol{x} \in \mathbb{R}^{n+1} \colon \sum_{i=1}^{n+1} x_i = |\mathcal{F}|\Bigr\}$$

with the halfspaces

$$H_{T,\geqslant} = \Bigl\{\boldsymbol{x} \in \mathbb{R}^{n+1} \colon \sum_{i\in T} x_i \geqslant \bigl|\mathcal{F}|_T\bigr|\Bigr\}$$

corresponding to all proper subsets $T \subsetneq [n+1]$.

Proof. Since $P_{\mathcal{F}}$ is a Minkowski summand in the permutahedron $P_{\mathcal{C}}$, it is defined by inequalities of the form $\sum_{i\in T} x_i \geqslant b_T$ for some parameters b_T. The minimum value of the linear function $\sum_{i\in T} x_i$ on a simplex Δ_S equals one if $S \subset T$, and equals zero otherwise. Therefore, $b_T = \bigl|\mathcal{F}|_T\bigr|$, as needed. □

It follows that $P_{\mathcal{F}}$ can be obtained by iteratively cutting the n-simplex

$$H_{\mathcal{F}} \cap \{\boldsymbol{x} \colon x_i \geqslant 1 \ \text{ for } i = 1, \ldots, n+1\}$$

by the hyperplanes $H_{T,\geqslant}$ corresponding to subsets T of cardinality $\geqslant 2$. In the case of the permutahedron, each of these cuts is nontrivial, that is, the corresponding hyperplane is not redundant. In general, the description of $P_{\mathcal{F}}$ in Proposition 1.5.9 is redundant. The concept of a building set will allow us to achieve an irredundant description of $P_{\mathcal{F}}$ for certain $\mathcal{F}$ and therefore describe the face posets. The resulting polytopes $P_{\mathcal{F}}$ will be simple; furthermore they will be obtained from a simplex by a sequence of face truncations.

Definition 1.5.10. A collection $\mathcal{B}$ of nonempty subsets of $[n+1]$ is called a *building set* on $[n+1]$ if the following two conditions are satisfied:

(a) $S', S'' \in \mathcal{B}$ with $S' \cap S'' \neq \varnothing$ implies $S' \cup S'' \in \mathcal{B}$;
(b) $\{i\} \in \mathcal{B}$ for all $i \in [n+1]$.

Remark. The terminology comes from a more general notion of a *building set in a finite lattice*, so that the building set above corresponds to the case of the Boolean lattice $2^{[n+1]}$. We do not give the general definition because it requires too much poset terminology. See [**88**, §3] for the details.

Note that a building set $\mathcal{B}$ on $[n+1]$ is *connected* if and only if $[n+1] \in \mathcal{B}$. Given $S \subsetneq [n+1]$ define the *contraction of S from $\mathcal{B}$* as

$$\mathcal{B}/S = \{T \setminus S \colon T \in \mathcal{B},\ T \setminus S \neq \varnothing\} = \{S' \subset [n+1]\setminus S \colon S' \in \mathcal{B} \ \text{ or } \ S' \cup S \in \mathcal{B}\}.$$

The restriction $\mathcal{B}|_S$ and the contraction $\mathcal{B}/S$ are building sets on S and $[n+1] \setminus S$ respectively. Note that $\mathcal{B}|_S$ is connected if and only if $S \in \mathcal{B}$. If $\mathcal{B}$ is connected, then $\mathcal{B}/S$ is also connected for any S.

We now consider polytopes $P_{\mathcal{B}}$ (1.20) corresponding to building sets $\mathcal{B}$. The following specification of Proposition 1.5.9 gives an irredundant description of $P_{\mathcal{B}}$ as an intersection of halfspaces.

PROPOSITION 1.5.11 ([**88**], [**187**]). *We have*

$$P_{\mathcal{B}} = \Bigl\{\boldsymbol{x} \in \mathbb{R}^{n+1} \colon \sum_{i=1}^{n+1} x_i = |\mathcal{B}|, \quad \sum_{i\in S} x_i \geqslant \bigl|\mathcal{B}|_S\bigr| \quad \text{for every } S \in \mathcal{B}\Bigr\}.$$

If $\mathcal{B}$ is connected, then this representation is irredundant, that is, every hyperplane $H_S = \bigl\{\boldsymbol{x} \in \mathbb{R}^{n+1} \colon \sum_{i\in S} x_i = \bigl|\mathcal{B}|_S\bigr|\bigr\}$ with $S \neq [n+1]$ defines a facet F_S of $P_{\mathcal{B}}$ (so that the number of facets of $P_{\mathcal{B}}$ is $|\mathcal{B}| - 1$).

PROOF. The halfspace $H_{T,\geqslant}$ in the presentation of $P_{\mathcal{B}}$ from Proposition 1.5.9 is irredundant if the intersection of $P_{\mathcal{B}}$ with the corresponding hyperplane H_T is a facet. This intersection is a face of $P_{\mathcal{B}}$ given by

$$P_{\mathcal{B}|_T} + P_{\mathcal{B}/T}$$

(since $P_{\mathcal{B}|_T}$ and $P_{\mathcal{B}/T}$ lie in complementary subspaces, their Minkowski sum is actually a product). In order for this face to have codimension one in $P_{\mathcal{B}}$, it is necessary, by Proposition 1.5.2, that the collection $\mathcal{B}|_T$ is connected. This condition is equivalent to $T \in \mathcal{B}$. If $\mathcal{B}$ is connected, then this condition is also sufficient, because then $\mathcal{B}/T$ is also connected, and $\dim(P_{\mathcal{B}|_T} + P_{\mathcal{B}/T}) = n - 1$ by Proposition 1.5.2. $\square$

COROLLARY 1.5.12. *If $\mathcal{B}$ is a connected building set on $[n+1]$, then every facet of $P_{\mathcal{B}}$ can be written as*

$$P_{\mathcal{B}|_T} \times P_{\mathcal{B}/T}$$

for some $T \in \mathcal{B} \setminus [n+1]$.

THEOREM 1.5.13. *The intersection of facets $F_{S_1} \cap \dots \cap F_{S_k}$ is nonempty (and therefore is a face of $P_{\mathcal{B}}$) if and only if the following two conditions are satisfied:*

(a) *for any i, j, $1 \leqslant i < j \leqslant k$, either $S_i \subset S_j$, or $S_j \subset S_i$, or $S_i \cap S_j = \varnothing$;*
(b) *if the sets $S_{i_1}, \dots, S_{i_p}$ are pairwise nonintersecting, then $S_{i_1} \sqcup \dots \sqcup S_{i_p} \notin \mathcal{B}$.*

DEFINITION 1.5.14. A subcollection $\{S_1, \dots, S_k\} \subset \mathcal{B}$ satisfying conditions (a) and (b) of Theorem 1.5.13 is called a *nested set*. Following [**188**], we refer to polytopes $P_{\mathcal{B}}$ (1.20) corresponding to building sets $\mathcal{B}$ as *nestohedra*.

PROOF OF THEOREM 1.5.13. Assume $F_{S_1} \cap \dots \cap F_{S_k} \neq \varnothing$.

If (a) fails, then $S_i \cup S_j \in \mathcal{B}$, and for any $\boldsymbol{x} \in F_{S_i} \cap F_{S_j}$ we have

$$\sum_{k\in S_i} x_k = \bigl|\mathcal{B}|_{S_i}\bigr|, \qquad \sum_{k\in S_j} x_k = \bigl|\mathcal{B}|_{S_j}\bigr|, \qquad \sum_{k\in S_i\cup S_j} x_k \geqslant \bigl|\mathcal{B}|_{S_i\cup S_j}\bigr|.$$

Adding the first two equalities and subtracting the third inequality we obtain

$$\sum_{k\in S_i\cap S_j} x_k \leqslant \bigl|\mathcal{B}|_{S_i}\bigr| + \bigl|\mathcal{B}|_{S_j}\bigr| - \bigl|\mathcal{B}|_{S_i\cup S_j}\bigr| < \bigl|\mathcal{B}|_{S_i}\bigr| + \bigl|\mathcal{B}|_{S_j}\bigr| - \bigl|\mathcal{B}|_{S_i} \cup \mathcal{B}|_{S_j}\bigr| = \bigl|\mathcal{B}|_{S_i\cap S_j}\bigr|$$

where the second inequality is strict because $\mathcal{B}|_{S_i} \cup \mathcal{B}|_{S_j} \subsetneq \mathcal{B}|_{S_i \cup S_j}$. Now the inequality $\sum_{k\in S_i\cap S_j} x_k < \bigl|\mathcal{B}|_{S_i\cap S_j}\bigr|$ contradicts Proposition 1.5.9.

If (b) fails, then $S_{i_1} \sqcup \cdots \sqcup S_{i_p} \in \mathcal{B}$, and for any $\boldsymbol{x} \in F_{S_{i_1}} \cap \cdots \cap F_{S_{i_p}}$ we have

$$\sum_{k\in S_{i_q}} x_k = \bigl|\mathcal{B}|_{S_{i_q}}\bigr| \quad \text{for } 1 \leqslant q \leqslant p, \quad \text{and} \quad \sum_{k\in S_{i_1}\sqcup\cdots\sqcup S_{i_p}} x_k \;\geqslant \bigl|\mathcal{B}|_{S_{i_1}\sqcup\cdots\sqcup S_{i_p}}\bigr|.$$

Subtracting the first p equalities from the last inequality we obtain

$$\bigl|\mathcal{B}|_{S_{i_1}}\bigr| + \cdots + \bigl|\mathcal{B}|_{S_{i_p}}\bigr| \geqslant \bigl|\mathcal{B}|_{S_{i_1}\sqcup\cdots\sqcup S_{i_p}}\bigr|.$$

This leads to a contradiction because $S_{i_1} \sqcup \cdots \sqcup S_{i_p} \in \mathcal{B}$.

Now assume that both (a) and (b) are satisfied. We need to show that $F_{S_1} \cap \cdots \cap F_{S_k} \neq \varnothing$.

We write $\boldsymbol{x} = \sum_{T\in\mathcal{B}} x^T$ and note that the inequality $\sum_{i\in S} x_i \geqslant \bigl|\mathcal{B}|_S\bigr|$ defining the facet F_S turns into equality if and only if $x_i^T = 0$ for every $T \in \mathcal{B}$, $T \not\subset S$ and $i \in S$ (this follows from (1.21)). Hence,

$$\begin{aligned} &(1.25) \quad \boldsymbol{x} = \sum_{T\in\mathcal{B}} x^T \;\in\; F_{S_1} \cap \cdots \cap F_{S_k} \\ &\qquad\iff\quad x_i^T = 0 \quad \text{whenever } T \in \mathcal{B},\ T \not\subset S_j,\ i \in S_j, \quad \text{for } 1 \leqslant j \leqslant k. \end{aligned}$$

We therefore need to find $\boldsymbol{x}$ whose coordinates satisfy the k conditions on the right hand side of (1.25). Given $T \in \mathcal{B}$, the jth condition is not void only if $T \not\subset S_j$ and $T \cap S_j \neq \varnothing$. We may assume without the loss of generality that the first k' conditions in (1.25) are not void, and the rest are void. That is, $T \not\subset S_j$ and $T \cap S_j \neq \varnothing$ for $1 \leqslant j \leqslant k'$, while $T \subset S_j$ or $T \cap S_j = \varnothing$ for $j > k'$. Then we claim that $T \setminus (S_1 \cup \cdots \cup S_{k'}) \neq \varnothing$. Indeed, otherwise choosing among $S_1, \ldots, S_{k'}$ the maximal subsets $S_{i_1}, \ldots, S_{i_p}$ (which are pairwise disjoint by (a)) we obtain $S_{i_1} \sqcup \cdots \sqcup S_{i_p} = T \cup S_{i_1} \cup \cdots \cup S_{i_p} \in \mathcal{B}$, which contradicts (b). Now setting $x_i^T = 1$ for only one $i \in T \setminus (S_1 \cup \cdots \cup S_{k'})$ and $x_i^T = 0$ for the rest, we obtain the required point $\boldsymbol{x}$ in the intersection $F_{S_1} \cap \cdots \cap F_{S_k}$. □

From the description of the face lattice of nestohedra in Theorem 1.5.13 it is easy to deduce their following main property.

THEOREM 1.5.15. *Every nestohedron $P_{\mathcal{B}}$ is a simple polytope.*

PROOF. By Proposition 1.5.2 we may assume that $\mathcal{B}$ is connected. A collection $S_1, \ldots, S_k$ may satisfy both conditions of Theorem 1.5.13 only if $k \leqslant n$. □

EXAMPLE 1.5.16. If $\mathcal{B}$ is a connected building set on a 2-element set, then $P_{\mathcal{B}}$ is an interval I^1. If $\mathcal{B}$ is a connected building set on a 3-element set, then $P_{\mathcal{B}}$ is a polygon, and only m-gons with $3 \leqslant m \leqslant 6$ arise in this way.

More examples will appear in the next subsections.

Proposition 1.5.11 gives a particular way to obtain a nestohedron $P_{\mathcal{B}}$ from a simplex by a sequence of hyperplane cuts. The next result shows that these hyperplane cuts can be organised in such a way that we get a sequence of face truncations (see Construction 1.1.13).

Let $\mathcal{B}_0 \subset \mathcal{B}_1$ be building sets on $[n+1]$, and $S \in \mathcal{B}_1$. We define a *decomposition of S into elements of $\mathcal{B}_0$* as $S = S_1 \sqcup \cdots \sqcup S_k$, where S_j are pairwise nonintersecting elements of $\mathcal{B}_0$ and k is minimal among such disjoint representations of S. It can be easily seen that this decomposition exists and is unique.

LEMMA 1.5.17. *Let $\mathcal{B}_0 \subset \mathcal{B}_1$ be connected building sets on $[n+1]$. Then $P_{\mathcal{B}_1}$ is obtained from $P_{\mathcal{B}_0}$ by a sequence of truncations of faces $G_i = \bigcap_{j=1}^{k_i} F_{S^i_j}$ corresponding to the decompositions $S^i = S^i_1 \sqcup \dots \sqcup S^i_{k_i}$ of elements $S^i \in \mathcal{B}_1 \setminus \mathcal{B}_0$, numbered in any order that is reverse to inclusion (i.e. $S^i \supseteq S^{i'} \Rightarrow i \leqslant i'$).*

PROOF. We use induction on the number $N = |\mathcal{B}_1| - |\mathcal{B}_0|$. For $N = 1$, we have $\mathcal{B}_1 = \mathcal{B}_0 \cup \{S^1\}$. We shall show that $P_{\mathcal{B}_1}$ is obtained from $P_{\mathcal{B}_0}$ by a single truncation of the face $G = F_{S^1_1} \cap \dots \cap F_{S^1_k}$, where $S^1 = S^1_1 \sqcup \dots \sqcup S^1_k$ is the decomposition of S^1 into elements of $\mathcal{B}_0$. Let $\widetilde{P}_{\mathcal{B}_0}$ denote the polytope obtained from $P_{\mathcal{B}_0}$ by truncation of G. Since both $\widetilde{P}_{\mathcal{B}_0}$ and $P_{\mathcal{B}_1}$ are n-dimensional polytopes (here we use the assumption that both $\mathcal{B}_0$ and $\mathcal{B}_1$ are connected), it is enough to verify that the face poset of $P_{\mathcal{B}_1}$ is a subposet of the face poset of $\widetilde{P}_{\mathcal{B}_0}$ (see Exercise 1.1.21).

The facets of $P_{\mathcal{B}_1}$ are F_{S^1} and F_{S_j} with $S_j \in \mathcal{B}_0$. We first consider a nonempty intersection of the form $F_{S_1} \cap \ldots \cap F_{S_\ell}$ in $P_{\mathcal{B}_1}$, i.e. a nested set $\{S_1, \ldots, S_\ell\}$ of $\mathcal{B}_1$, with all $S_j \in \mathcal{B}_0$. Then obviously, $\{S_1, \ldots, S_\ell\}$ is a nested set of $\mathcal{B}_0$, i.e. $F_{S_1} \cap \ldots \cap F_{S_\ell} \neq \varnothing$ in $P_{\mathcal{B}_0}$. Furthermore, since S^1 is the only element of $\mathcal{B}_1 \setminus \mathcal{B}_0$, we have that

$$S_{j_1} \sqcup \dots \sqcup S_{j_p} \neq S^1 = S^1_1 \sqcup \dots \sqcup S^1_k$$

for any $\{j_1, \ldots, j_p\} \subset [\ell]$. The latter condition implies that $\{S^1_1, \ldots, S^1_k\} \not\subset \{S_1, \ldots, S_\ell\}$, i.e. $F_{S_1} \cap \ldots \cap F_{S_\ell} \not\subset G$ in the face poset of $P_{\mathcal{B}_0}$. By the description of the face poset of $\widetilde{P}_{\mathcal{B}_0}$ given in Construction 1.1.13, this implies that $F_{S_1} \cap \ldots \cap F_{S_\ell} \neq \varnothing$ in $\widetilde{P}_{\mathcal{B}_0}$.

Now we consider a nonempty intersection of the form $F_{S^1} \cap F_{S_1} \cap \ldots \cap F_{S_\ell}$ in $P_{\mathcal{B}_1}$, i.e. a nested set $\{S^1, S_1, \ldots, S_\ell\}$ of $\mathcal{B}_1$, with $S_j \in \mathcal{B}_0$ and $S^1 \in \mathcal{B}_1 \setminus \mathcal{B}_0$. We claim that $\{S^1_1, \ldots, S^1_k, S_1, \ldots, S_\ell\}$ is a nested set of $\mathcal{B}_0$, i.e. $G \cap F_{S_1} \cap \ldots \cap F_{S_\ell} \neq \varnothing$ in $P_{\mathcal{B}_0}$. To do this we need to verify (a) and (b) of Theorem 1.5.13.

We need to check condition (a) for pairs of the form S^1_p, S_q; for other pairs it is obvious. That is, we need to check that if $S^1_p \cap S_q \neq \varnothing$, then one of S^1_p, S_q is contained in the other. The condition $S^1_p \cap S_q \neq \varnothing$ implies that $S^1 \cap S_q \neq \varnothing$. Since $\{S^1, S_1, \ldots, S_\ell\}$ is a nested set of $\mathcal{B}_1$, we obtain that $S^1_p \subset S^1 \subset S_q$ or $S_q \subset S^1$. By the minimality of the decomposition $S^1 = S^1_1 \sqcup \dots \sqcup S^1_k$, the inclusion $S_q \subset S^1$ implies that S_q is contained in some S^1_r, which can be only S^1_p, since $S^1_p \cap S_q \neq \varnothing$.

To verify condition (b) of Theorem 1.5.13 for $\{S^1_1, \ldots, S^1_k, S_1, \ldots, S_\ell\}$, we consider a subcollection $\{S^1_{i_1}, \ldots, S^1_{i_p}, S_{j_1}, \ldots, S_{j_q}\}$ consisting of pairwise nonintersecting subsets. We need to check that its union is not in $\mathcal{B}_0$. For obvious reasons, we may assume that $p > 0$ and $q > 0$. Since $\{S^1, S_1, \ldots, S_\ell\}$ is a nested set of $\mathcal{B}_1$, we have that either $S_{j_i} \subset S^1$ or $S_{j_i} \cap S^1 = \varnothing$ for each $i = 1, \ldots, q$. Suppose that $S^1_{i_1} \sqcup \dots \sqcup S^1_{i_p} \sqcup S_{j_1} \sqcup \dots \sqcup S_{j_q} \in \mathcal{B}_0$. Then $S^1 \sqcup S_{j_1} \sqcup \dots \sqcup S_{j_q} \in \mathcal{B}_1$ by the definition of the building set. If any of S_{j_i} is disjoint with S^1, then we get a contradiction with condition (b) for the nested set $\{S^1, S_1, \ldots, S_\ell\}$ of $\mathcal{B}_1$. Therefore, $S_{j_i} \subset S^1$ for $i = 1, \ldots, q$, so that $S^1_{i_1} \sqcup \dots \sqcup S^1_{i_p} \sqcup S_{j_1} \sqcup \dots \sqcup S_{j_q} \subset S^1$. By the argument of the previous paragraph, for each S_{j_i} we have that $S_{j_i} \subset S^1_r$ or $S^1_r \subset S_{j_i}$ for some $r = 1, \ldots, k$. Then it follows from the minimality of the decomposition $S^1 = S^1_1 \sqcup \dots \sqcup S^1_k$ and the definition of a building set that $S^1_{i_1} \sqcup \dots \sqcup S^1_{i_p} \sqcup S_{j_1} \sqcup \dots \sqcup S_{j_q} = S^1$, which contradicts the assumption that $S^1 \notin \mathcal{B}_0$.

Hence, $\{S^1_1, \ldots, S^1_k, S_1, \ldots, S_\ell\}$ is a nested set of $\mathcal{B}_0$. Similarly to the case considered in the previous paragraph, we also obtain that $F_{S_1} \cap \ldots \cap F_{S_\ell} \not\subset G$ in the face poset of $P_{\mathcal{B}_0}$. Once again, by the description of the face poset of $\widetilde{P}_{\mathcal{B}_0}$ given in Construction 1.1.13, this implies that $F_{S^1} \cap F_{S_1} \cap \ldots \cap F_{S_\ell} \neq \varnothing$ in $\widetilde{P}_{\mathcal{B}_0}$.

It follows that the face poset of $P_{\mathcal{B}_1}$ is indeed contained as a subposet in the face poset of $\widetilde{P}_{\mathcal{B}_0}$, and thus $P_{\mathcal{B}_1} = \widetilde{P}_{\mathcal{B}_0}$.

It now remains to finish the induction. Assuming the theorem holds for $M < N$, we shall prove it for $M = N$. Since S^1 is not contained in any other S^i, the collection of sets $\mathcal{B}'_0 = \mathcal{B}_0 \cup \{S^1\}$ is a building set. By the induction assumption, $P_{\mathcal{B}'_0}$ is obtained from $P_{\mathcal{B}_0}$ by truncation of the face corresponding to the decomposition of S^1, and $P_{\mathcal{B}_1}$ is obtained from $P_{\mathcal{B}'_0}$ by a sequence of truncations corresponding to the decompositions of S^i for $i = 2, \ldots, N$. ☐

REMARK. The proof given above only establishes a combinatorial equivalence between $P_{\mathcal{B}_1}$ and $\widetilde{P}_{\mathcal{B}_0}$. Since Proposition 1.5.11 gives a geometric presentation of nestohedra by a sequence of hyperplane cuts, it follows easily that the face truncations of $P_{\mathcal{B}_0}$ giving $P_{\mathcal{B}_1}$ are also geometric.

THEOREM 1.5.18. *Every nestohedron $P_{\mathcal{B}}$ corresponding to a connected building set $\mathcal{B}$ can be obtained from a simplex by a sequence of face truncations.*

PROOF. Assume that $\mathcal{B}$ is a connected building set on $[n+1]$. Then we have $\mathcal{S} \subset \mathcal{B}$, where $\mathcal{S}$ is the connected building set of Example 1.5.3, whose corresponding nestohedron is an n-simplex. Now apply Lemma 1.5.17. ☐

The following construction, suggested by N. Erokhovets, will allow us to show that every nestohedron can be obtained from a connected building set, up to combinatorial equivalence.

CONSTRUCTION 1.5.19 (Substitution of building sets). Let $\mathcal{B}_1, \ldots, \mathcal{B}_{n+1}$ be connected building sets on $[k_1], \ldots, [k_{n+1}]$. Then, for every connected building set $\mathcal{B}$ on $[n+1]$, we define a connected building set $\mathcal{B}(\mathcal{B}_1, \ldots, \mathcal{B}_{n+1})$ on $[k_1] \sqcup \ldots \sqcup [k_{n+1}] = [k_1 + \ldots + k_{n+1}]$, consisting of elements $S^i \in \mathcal{B}_i$ and $\bigsqcup\limits_{i \in S} [k_i]$, where $S \in \mathcal{B}$.

When $\mathcal{B}_1, \ldots, \mathcal{B}_n$ are singletons $\{1\}, \ldots, \{n\}$, we shall write $\mathcal{B}(1, 2, \ldots, n, \mathcal{B}_{n+1})$ instead of $\mathcal{B}\bigl(\{1\}, \{2\}, \ldots, \{n\}, \mathcal{B}_{n+1}\bigr)$.

LEMMA 1.5.20. *Let $\mathcal{B}$, $\mathcal{B}_1, \ldots, \mathcal{B}_{n+1}$ be connected building sets on $[n+1]$, $[k_1], \ldots, [k_{n+1}]$, and let $\mathcal{B}' = \mathcal{B}(\mathcal{B}_1, \ldots, \mathcal{B}_{n+1})$. Then $P_{\mathcal{B}'} \approx P_{\mathcal{B}} \times P_{\mathcal{B}_1} \times \cdots \times P_{\mathcal{B}_{n+1}}$.*

PROOF. Set $\mathcal{B}'' = \mathcal{B} \sqcup \mathcal{B}_1 \sqcup \ldots \sqcup \mathcal{B}_{n+1}$ and define the map $\varphi\colon \mathcal{B}'' \to \mathcal{B}'$ by

$$\varphi(S) = \begin{cases} S & \text{if } S \in \mathcal{B}_i \\ \bigsqcup\limits_{i \in S} [k_i] & \text{if } S \in \mathcal{B}\,. \end{cases}$$

Then φ generates a bijection between $\mathcal{B}'' \setminus \mathcal{B}''_{\max}$ and $\mathcal{B}' \setminus [k_1 + \cdots + k_{n+1}]$, where $\mathcal{B}''_{\max} = \{[n+1], [k_1], \ldots, [k_{n+1}]\}$ denotes the collection of maximal sets of $\mathcal{B}''$. Let $\mathcal{S} \subset \mathcal{B} \setminus [n+1]$ and $\mathcal{S}_i \subset B_i \setminus [k_i]$. Notice that $\varphi(\mathcal{S}) \cup \bigcup_{i=1}^{n+1} \varphi(\mathcal{S}_i)$ is a nested set of $\mathcal{B}'$ if and only if $\mathcal{S}$ is a nested set of $\mathcal{B}$ and $\mathcal{S}_i$ is a nested set of $\mathcal{B}_i$ for all i. It follows that $P_{\mathcal{B}'} \approx P_{\mathcal{B}''} = P_{\mathcal{B}} \times P_{\mathcal{B}_1} \times \cdots \times P_{\mathcal{B}_{n+1}}$. ☐

EXAMPLE 1.5.21. Assume that each of $\mathcal{B}$, $\mathcal{B}_1$, $\mathcal{B}_2$ is the building set $\{\{1\}, \{2\}, \{1, 2\}\}$ corresponding to the segment $\mathbb{I}$. Let us describe the building set $\mathcal{B}(\mathcal{B}_1, \mathcal{B}_2)$. In the building set $\{\{a\}, \{b\}, \{a, b\}\}$, we substitute a by

$\mathcal{B}_1 = \{\{1\},\{2\},\{1,2\}\}$ and b by $\mathcal{B}_2 = \{\{3\},\{4\},\{3,4\}\}$. As a result, we obtain the connected building set $\mathcal{B}'$, consisting of $\{1\},\{2\},\{3\},\{4\},\{1,2\},\{3,4\},[4]$. Its corresponding nestohedron is obtained by truncating two nonadjacent edges of a 3-simplex; it is combinatorially equivalent to a 3-cube.

The facet correspondence φ between the combinatorial cubes $P_{\mathcal{B}} \times P_{\mathcal{B}_1} \times P_{\mathcal{B}_2}$ and $P_{\mathcal{B}'}$ is given by

$$\begin{aligned} &\{1\} \in \mathcal{B}_1 \mapsto \{1\} \in \mathcal{B}', && \{2\} \in \mathcal{B}_1 \mapsto \{2\} \in \mathcal{B}', \\ &\{3\} \in \mathcal{B}_2 \mapsto \{3\} \in \mathcal{B}', && \{4\} \in \mathcal{B}_2 \mapsto \{4\} \in \mathcal{B}', \\ &\{a\} \in \mathcal{B}_2 \mapsto \{1,2\} \in \mathcal{B}', && \{b\} \in \mathcal{B}_2 \mapsto \{3,4\} \in \mathcal{B}'. \end{aligned}$$

EXAMPLE 1.5.22. Let $\mathcal{B} = \{\{1\},\dots,\{n+1\},[n+1]\}$ be the building set corresponding to the simplex Δ^n and let $\mathcal{B}_1,\dots,\mathcal{B}_{n+1}$ be arbitrary connected building sets on $[k_1],\dots,[k_{n+1}]$. Then

$$\mathcal{B}' = \mathcal{B}(\mathcal{B}_1,\dots,\mathcal{B}_{n+1}) = (\mathcal{B}_1 \sqcup \dots \sqcup \mathcal{B}_{n+1}) \cup [k_1 + \dots + k_{n+1}],$$

and $P_{\mathcal{B}'} \approx \Delta^n \times P_{\mathcal{B}_1} \times \dots \times P_{\mathcal{B}_{n+1}}$.

PROPOSITION 1.5.23. *For each nestohedron $P_{\mathcal{B}}$ there exists a connected building set $\mathcal{B}'$ such that $P_{\mathcal{B}} \approx P_{\mathcal{B}'}$.*

PROOF. Indeed, any building set $\mathcal{B}$ can be written as $\mathcal{B}_1 \sqcup \dots \sqcup \mathcal{B}_k$, where $\mathcal{B}_i$ are connected building sets on $[k_i+1]$. Define a building set $\widetilde{\mathcal{B}} = \mathcal{B}_1(1,\dots,k_1,\mathcal{B}_2) \sqcup \mathcal{B}_3 \sqcup \dots \sqcup \mathcal{B}_k$, giving the same combinatorial polytope. We have that $\widetilde{\mathcal{B}}$ is a product (disjoint union) of $k-1$ connected building sets. Then we apply again a substitution to $\widetilde{\mathcal{B}}$, and so on. At the end we obtain a connected building set $\mathcal{B}'$. □

Graph associahedra.

DEFINITION 1.5.24. Let Γ be a graph on the vertex set $[n+1]$ without loops and multiple edges (a *simple* graph). The *graphical building set* $\mathcal{B}_\Gamma$ consists of all nonempty subsets $S \subset [n+1]$ such that the graph $\Gamma|_S$ is connected.

The nestohedron $P_\Gamma = P_{\mathcal{B}_\Gamma}$ corresponding to a graphical building set is called a *graph-associahedron* [**57**].

EXAMPLE 1.5.25 (associahedron). Let Γ be a 'path' with n edges $\{i,i+1\}$ for $1 \leqslant i \leqslant n$. Then $\mathcal{B}_\Gamma$ consists of all 'segments' of the form $[i,j] = \{i,i+1,\dots,j\}$ where $1 \leqslant i \leqslant j \leqslant n+1$. To describe the face poset of the corresponding polytope P_Γ it is convenient to use brackets in a string of $n+2$ letters $a_1a_2\cdots a_{n+2}$. We associate with a segment $[i,j]$ a pair of brackets before a_i and after a_{j+1}. Using Theorem 1.5.13 it is easy to see that the facets corresponding to n different segments intersect at a vertex if and only if the corresponding bracketing of $a_1a_2\cdots a_{n+2}$ with n pairs of brackets is correct. In particular, the vertices of $\mathcal{B}_\Gamma$ correspond to all different ways to obtain a product $a_1a_2\cdots a_{n+2}$ when multiplication is not associative. The number of vertices is therefore equal to $\frac{1}{n+2}\binom{2n+2}{n+1}$, the $(n+1)$th *Catalan number*. Two vertices are adjacent if and only if the bracketing corresponding to one vertex can be obtained from the bracketing corresponding to the other vertex by deleting a pair of brackets and inserting, in a unique way, another pair of brackets different from the deleted one. That is, two vertices are adjacent if they correspond to a single application of the associative law. This explains the name *associahedron* for the polytope $\mathcal{P}_\Gamma$ of this example; we shall denote it As^n.



FIGURE 1.3. 3-dimensional associahedron and the corresponding graph.

Proposition 1.5.11 describes the associahedron as the result of consecutive hyperplane cuts of a simplex, see Figure 1.3 for the case $n = 3$. It can also be obtained by hyperplane cuts from a cube, as described in the next theorem.

THEOREM 1.5.26. *The image of* As^n *under a certain affine transformation* $\mathbb{R}^{n+1} \to \mathbb{R}^n$ *is the intersection of the cube*

$$\big\{\boldsymbol{y} \in \mathbb{R}^n \colon 0 \leqslant y_j \leqslant j(n+1-j) \quad \textit{for} \ \ 1 \leqslant j \leqslant n\big\}$$

with the halfspaces

$$\big\{\boldsymbol{y} \in \mathbb{R}^n \colon y_j - y_k + (j-k)k \geqslant 0\big\}$$

for $1 \leqslant k < j \leqslant n$.

PROOF. Proposition 1.5.11 gives the following presentation:

$$(1.26) \quad As^n = \Big\{\boldsymbol{x} \in \mathbb{R}^{n+1} \colon \sum_{k=1}^{n+1} x_k = \frac{(n+1)(n+2)}{2}, \\ \sum_{k=i}^{j} x_k \geqslant \frac{(j-i+1)(j-i+2)}{2} \quad \text{for } 1 \leqslant i \leqslant j \leqslant n+1\Big\}.$$

Apply the affine transformation $\mathbb{R}^{n+1} \to \mathbb{R}^n$ given by

$$(x_1, \dots, x_{n+1}) \mapsto (z_1, \dots, z_n) \quad \text{where } z_l = \sum_{k=1}^{l} x_k, \quad \text{for } 1 \leqslant l \leqslant n.$$

Now we rewrite the inequalities of (1.26) in the new coordinates $(z_1, \dots, z_n)$. The inequalities with $i = 1$ (corresponding to the facets F_S with $\{1\} \in S$) become

$$(1.27) \qquad z_j \geqslant \frac{j(j+1)}{2}, \quad \text{for } 1 \leqslant j \leqslant n.$$

Inequalities (1.26) with $j = n+1$ (corresponding to F_S with $\{n+1\} \in S$) become

$$\frac{(n+1)(n+2)}{2} - z_{i-1} \geqslant \frac{(n+2-i)(n+3-i)}{2}, \quad \text{for } 2 \leqslant i \leqslant n+1,$$

or equivalently,

$$(1.28) \qquad z_j \leqslant (n+2)j - \frac{j(j+1)}{2}, \quad \text{for } 1 \leqslant j \leqslant n.$$

The remaining inequalities (1.26) take the form

$$z_j - z_{i-1} \geqslant \frac{(j-i+1)(j-i+2)}{2} \quad \text{for } 1 < i \leqslant j < n+1. \tag{1.29}$$

Now the required presentation is obtained from (1.27), (1.28) and (1.29) by applying the shift $y_j = z_j - \frac{j(j+1)}{2}$ and setting $k = i - 1$. □

Example 1.5.27. The case $n = 3$ of Theorem 1.5.26 is shown in Figure 1.4 (right). This polytope is obtained by cutting a 3-cube (more precisely, a parallelepiped) specified by the equations



Figure 1.4. 3-dimensional associahedron cut from a cube.

$$0 \leqslant y_1 \leqslant 3,\ 0 \leqslant y_2 \leqslant 4,\ 0 \leqslant y_3 \leqslant 3$$

by the three hyperplanes

$$y_2 - y_1 + 1 = 0,\ y_3 - y_1 + 2 = 0,\ y_3 - y_2 + 1 = 0.$$

Another way to cut a 3-dimensional combinatorial associahedron from a 3-cube is shown in Figure 1.4 (left); this time we cut three nonadjacent and pairwise orthogonal edges. The two associahedra in Figure 1.4 are not affinely equivalent.

The associahedron As^n first appeared (as a combinatorial object) in the work of Stasheff [**210**] as the space of parameters for the higher associativity of the $(n+2)$-fold product map in an H-space. For more information about the associahedra we refer to [**98**] and [**39**, Lec. II], where the reader may find other geometric and combinatorial realisations of As^n.

Example 1.5.28 (permutahedron revisited). Let Γ be a complete graph; then $\mathcal{B}_\Gamma$ is the complete building set $\mathcal{C}$ and P_Γ is the permutahedron Pe^n, see Figure 1.5 for the case $n = 3$.

Example 1.5.29 (cyclohedron). Let Γ be a 'cycle' consisting of $n+1$ edges $\{i, i+1\}$ for $1 \leqslant i \leqslant n$ and $\{n+1, 1\}$. The corresponding P_Γ is known as the *cyclohedron* Cy^n, or *Bott–Taubes polytope*, see Figure 1.6. It was first introduced in [**29**] in connection with the study of link invariants.

Example 1.5.30 (stellahedron). Let Γ be a 'star' consisting of n edges $\{i, n+1\}$, $1 \leqslant i \leqslant n$, emanating from the point $n+1$. The corresponding P_Γ is known as the *stellahedron* St^n, see Figure 1.7.



FIGURE 1.5. 3-dimensional permutahedron and the corresponding graph.



FIGURE 1.6. 3-dimensional cyclohedron and the corresponding graph.



FIGURE 1.7. 3-dimensional stellahedron and the corresponding graph.

Exercises.

1.5.31. Every collection $\mathcal{F}$ of subsets in $[n+1]$ may be completed in a unique way to a building set by iteratively adding to $\mathcal{F}$ the unions $S_1 \cup S_2$ of intersecting sets ($S_1 \cap S_2 \neq \varnothing$) until the process stops. Denote the resulting building set by $\widehat{\mathcal{F}}$. Show that $\widehat{\mathcal{F}}$ is connected if and only if $\mathcal{F}$ is connected, and that $\dim P_{\widehat{\mathcal{F}}} = \dim P_{\mathcal{F}}$ (hint: for a pair of intersecting sets S_1, S_2 compare the polytopes $\Delta_{S_1} + \Delta_{S_2}$ and $\Delta_{S_1} + \Delta_{S_2} + \Delta_{S_1 \cup S_2}$). Use this fact to complete the proof of Proposition 1.5.2.

1.5.32. Finish the argument in the proof of Proposition 1.5.5.

1.5.33. Complete the details in the proof of Theorem 1.5.7.

1.5.34. Prove Proposition 1.5.8.

1.5.35. Given two building sets $\mathcal{B}_0 \subset \mathcal{B}_1$, show that a decomposition $S = S_1 \sqcup \cdots \sqcup S_k$ of $S \in \mathcal{B}_1$ into elements $S_i \in \mathcal{B}_0$ with minimal k is unique.

1.5.36. A combinatorial polytope obtained from a simplex by a sequence of face truncations is called a *truncated simplex*. Notice that a truncated simplex is a simple polytope. By Theorem 1.5.18, every nestohedron corresponding to a connected nested set is a truncated simplex. Give an example of

(a) a simple polytope which is not a truncated simplex;
(b) a truncated simplex which is not a nestohedron;
(c) a Minkowski sum of simplices $P_{\mathcal{F}}$ given by (1.20) which is not a truncated simplex.

1.5.37. Consider the following connected building set on [4]:

$$\mathcal{B} = \bigl\{\{1\}, \{2\}, \{3\}, \{4\}, \{1,2,3\}, \{2,3,4\}, [4]\bigr\}.$$

Then $P_{\mathcal{B}}$ is combinatorially equivalent to the polytope shown in Fig. 1.1, right.

1.6. Flagtopes and truncated cubes

Definition 1.6.1. A simple polytope P is called a *flagtope* (or *flag polytope*) if every collection of its pairwise intersecting facets has a nonempty intersection. (The origins of this terminology will be explained in Section 2.3.)

Flagtopes and flag simplicial complexes (which are the subject of Section 2.3) feature in the well-known *Charney–Davis conjecture* [**58**] and its generalisation due to Gal [**97**]. According to the Gal conjecture, the components of the γ-vector of a flagtope are nonnegative.

Gal's conjecture is important as it connects the combinatorial study of polytopes and sphere triangulations to differential geometry and topology of manifolds. This conjecture was proved in special cases.

Although not all nestohedra are flagtopes (a simplex is the easiest counterexample), flag nestohedra constitute an important family. In particular, all graph-associahedra (and therefore, the classical series of associahedra and permutahedra) are flagtopes, see Proposition 1.6.6.

As we have seen in Theorem 1.5.26, the associahedron can be obtained by hyperplane cuts from a cube. In this section we give a proof of a much more general and precise result: a nestohedron is a flagtope if and only if it can be obtained from a cube by a sequence of truncations of faces of codimension 2 (we refer to such polytopes as *2-truncated cubes*), see Theorem 1.6.16. This result was proved by Buchstaber and Volodin in [**54**, Theorem 6.6]. On the other hand, it can be easily seen that the Gal conjecture is valid for 2-truncated cubes (see Proposition 1.6.12). This observation (formulated in terms of the dual operation of stellar subdivision at an edge) was present in the work of Charney–Davis and later Gal, and used to support their conjectures. As a corollary we obtain that the Gal conjecture holds for all flag nestohedra.

Example 1.6.2.

1. The cube I^n is a flagtope, but the simplex Δ^n is not if $n > 1$.
2. The product $P \times Q$ of two flagtopes is a flagtope.
3. The connected sum $P \# Q$ of two simple n-polytopes and the vertex truncation $\mathrm{vt}(P)$ are not flagtopes if $n > 1$.

A polytope P is said to be *triangle-free* if it does not contain a triangular 2-face.

Proposition 1.6.3. *A flagtope is triangle-free.*

Proof. Assume that a flagtope P of dimension n contains a triangular 2-face T with vertices v_1, v_2, v_3 and edges e_1, e_2, e_3, where v_i is opposite to e_i. Since P is simple, each e_i is an intersection of $n-1$ facets. Hence, for each $i = 1, 2, 3$ there is a unique facet, say F_i, which contains the edge e_i but not the triangle T. Also, T is an intersection of $n-2$ facets, and we may assume that $T = \bigcap_{i=4}^{n+1} F_i$. Now we observe that $\bigcap_{i=1, i\neq j}^{n+1} F_i = v_j$ for $j = 1, 2, 3$. This implies that the intersection of any pair of facets among $F_1, \ldots, F_{n+1}$ is nonempty. On the other hand, $\bigcap_{i=1}^{n+1} F_i = \varnothing$ because P is simple and n-dimensional. Therefore P cannot be a flagtope. □

The converse to Proposition 1.6.3 does not hold in general (see exercises), but it is valid for polytopes with few facets:

Theorem 1.6.4 ([**26**]). *If P is a triangle-free convex n-polytope then $f_i(P) \geqslant f_i(I^n)$ for $i = 0, \ldots, n$. In particular, such P has at least $2n$ facets. Furthermore if P is simple then*

- (a) *$f_{n-1}(P) = 2n$ implies that $P = I^n$;*
- (b) *$f_{n-1}(P) = 2n+1$ implies that $P = P_5 \times I^{n-2}$ where P_5 is a pentagon;*
- (c) *$f_{n-1}(P) = 2n+2$ implies that $P = P_6 \times I^{n-2}$ or $P = Q \times I^{n-3}$ or $P = P_5 \times P_5 \times I^{n-4}$ where P_6 is a hexagon and Q is a 3-polytope obtained from a pentagonal prism by truncating one of the edges forming a pentagonal facet.*

Corollary 1.6.5. *A triangle-free simple n-polytope with at most $2n+2$ facets is a flagtope.*

Another source of examples of flagtopes is provided by graph-associahedra (see Definition 1.5.24):

Proposition 1.6.6. *Every graph-associahedron P_Γ is a flagtope.*

Proof. Let $F_{S_1}, \ldots, F_{S_k}$ be a set of facets of P_Γ with nonempty pairwise intersections. We need to check that $F_{S_1} \cap \cdots \cap F_{S_k} \neq \varnothing$, i.e. that condition (b) of Theorem 1.5.13 is satisfied (condition (a) holds automatically as it depends only on pairwise intersections). Let $S_{i_1}, \ldots, S_{i_p}$ be pairwise nonintersecting sets among $S_1, \ldots, S_k$; then $S_{i_r} \cup S_{i_s} \notin \mathcal{B}_\Gamma$ for $1 \leqslant r < s \leqslant p$ because $F_{S_{i_r}} \cap F_{S_{i_s}} \neq \varnothing$. Therefore, all subgraphs $\Gamma|_{S_{i_r} \cup S_{i_s}}$ are disconnected, which implies that $\Gamma|_{S_{i_1} \cup \cdots \cup S_{i_p}}$ is also disconnected. Thus, $S_{i_1} \cup \cdots \cup S_{i_p} \notin \mathcal{B}_\Gamma$, and $F_{S_1} \cap \cdots \cap F_{S_k} \neq \varnothing$ by Theorem 1.5.13. □

The main conjecture about the face numbers of flagtopes uses the notion of the γ-vector $\gamma(P) = (\gamma_0, \dots, \gamma_{[n/2]})$ (see Definition 1.3.10):

Conjecture 1.6.7 (Gal [**97**]). *A flagtope P of dimension n satisfies $\gamma_i(P) \geqslant 0$ for $i = 0, \dots, \left[\frac{n}{2}\right]$.*

We shall verify the Gal conjecture for all flag nestohedra. To do this we shall show that any flag nestohedron can be obtained from a cube by consecutive truncations of codimension-2 faces.

Our first task is therefore to describe how the γ-vector changes under face truncations. For this it is convenient to use the H-polynomial given by (1.10), and the γ-polynomial

$$\gamma(P)(\tau) = \gamma_0 + \gamma_1\tau + \dots + \gamma_{[n/2]}\tau^{[n/2]}.$$

Proposition 1.6.8. *Let the polytope Q be obtained from a simple n-polytope P by truncation of a k-dimensional face G. Then*

(a) $H(Q) = H(P)(s,t) + stH(G)H(\Delta^{n-k-2})$,

(b) $\gamma(Q) = \gamma(P) + \tau\gamma(G)\gamma(\Delta^{n-k-2})$.

Proof. The truncation removes G and creates a face $G \times \Delta^{n-k-1}$, so that

$$f_i(Q) = f_i(P) + f_i(G \times \Delta^{n-k-1}) - f_i(G), \quad \text{for } 0 \leqslant i \leqslant n.$$

Hence, $F(Q) = F(P) + tF(G)F(\Delta^{n-k-1}) - t^{n-k}F(G)$, and

$$\begin{aligned} H(Q) &= H(P) + tH(G)H(\Delta^{n-k-1}) - t^{n-k}H(G) \\ &= H(P) + tH(G)\left(\sum_{i=0}^{n-k-1} s^i t^{n-k-1-i} - t^{n-k-1}\right) \\ &= H(P) + stH(G)\left(\sum_{j=0}^{n-k-2} s^j t^{n-k-2-j}\right) = H(P) + stH(G)H(\Delta^{n-k-2}), \end{aligned}$$

which proves (a). Furthermore,

$$\begin{aligned} \sum_{i=0}^{[\frac{n}{2}]} \gamma_i(Q)(st)^i(s+t)^{n-2i} &= H(Q) = \sum_{i=0}^{[\frac{n}{2}]} \gamma_i(P)(st)^i(s+t)^{n-2i} \\ &+ st\left(\sum_{p=0}^{[\frac{k}{2}]} \gamma_p(G)(st)^p(s+t)^{k-2p}\right)\left(\sum_{q=0}^{[\frac{n-k-2}{2}]} \gamma_q(\Delta^{n-k-2})(st)^q(s+t)^{n-k-2-2q}\right) \\ = \sum_{i=0}^{[\frac{n}{2}]} \gamma_i(P)(st)^i(s+t)^{n-2i} &+ \sum_{p=0}^{[\frac{k}{2}]} \sum_{p=0}^{[\frac{n-k-2}{2}]} \gamma_p(G)\gamma_q(\Delta^{n-k-2})(st)^{p+q+1}(s+t)^{n-2(p+q+1)}. \end{aligned}$$

Hence, $\gamma_i(Q) = \gamma_i(P) + \sum_{p+q=i-1} \gamma_p(G)\gamma_q(\Delta^{n-k-2})$, which proves (b). □

Definition 1.6.9. We refer to a truncation of a face of codimension 2 as a 2-*truncation.* A combinatorial polytope obtained from a cube by 2-truncations will be called a 2-*truncated cube.*

The following corollary is the dual of a result from [**97**]:

COROLLARY 1.6.10. *Let the polytope Q be obtained from a simple polytope P by 2-truncation of a face G. Then*

(a) $H(Q) = H(P) + stH(G)$,
(b) $\gamma(Q) = \gamma(P) + \tau\gamma(G)$.

PROPOSITION 1.6.11. *Each face of a 2-truncated cube is a 2-truncated cube.*

PROOF. It is enough to show that if P is a 2-truncated cube, then all the facets of P are 2-truncated cubes. The proof is by induction on the number of truncated faces. Let the polytope Q be obtained from a 2-truncated cube P by 2-truncation of a face G of codimension 2. Then the new facet is $G \times I$, and it is a 2-truncated cube by the induction assumption. Every other facet F' of the polytope Q is either a facet of P, or obtained from a facet F'' of P by 2-truncation of a face $G' \subset F''$. $\square$

PROPOSITION 1.6.12. *Every 2-truncated cube P satisfies $\gamma_i(P) \geq 0$, i.e. the Gal conjecture holds for 2-truncated cubes.*

PROOF. We proceed by induction on the dimension of P, using Proposition 1.6.11 and the formula $\gamma(Q) = \gamma(P) + \tau\gamma(G)$. $\square$

Here is a criterion for a nestohedron to be a flagtope.

PROPOSITION 1.6.13 ([**54**], [**188**]). *Let $\mathcal{B}$ be a building set on $[n+1]$. Then the nestohedron $P_{\mathcal{B}}$ is a flagtope if and only if for every element $S \in \mathcal{B}$ with $|S| > 1$ there exist elements $S', S'' \in \mathcal{B}$ such that $S' \sqcup S'' = S$.*

PROOF. Suppose P_B is a flagtope. Consider an element $S \in B$. Then we may write $S = S_1 \sqcup \cdots \sqcup S_k$, where $S_1, \ldots, S_k \in \mathcal{B} \setminus \{S\}$ and k is minimal among such decompositions of S. Then for any subset $J \subset [k]$ with $1 < |J| < k$ we have that $\bigsqcup_{j\in J} S_j \notin \mathcal{B}$, since otherwise k can be decreased. If $k > 2$ then, by Theorem 1.5.13, the facets $F_{S_1}, \ldots, F_{S_k}$ of $P_{\mathcal{B}}$ intersect pairwise, but have empty common intersection. Therefore, $k = 2$.

Suppose for each element $S \in \mathcal{B}$ with $|S| > 1$, there exist elements $S', S'' \in \mathcal{B}$ such that $S' \sqcup S'' = S$. Let $F_{S_1}, \ldots, F_{S_k}$, $k \geqslant 3$, be a minimal collection of facets that intersect pairwise but have empty common intersection. We shall lead this to a contradiction by finding a nontrivial subcollection of $F_{S_1}, \ldots, F_{S_k}$ with empty common intersection.

Assume there is a set $\widetilde{S} \in \mathcal{B}|_S$ intersecting more than one S_i, but not intersecting every S_i. Then the subcollection of facets F_{S_i} satisfying $S_i \cap \widetilde{S} \neq \varnothing$ will have empty common intersection, since

$$\bigsqcup_{S_i\colon S_i\cap\widetilde{S}\neq\varnothing} S_i \in \mathcal{B}$$

by definition of a building set.

It remains to find $\widetilde{S} \in \mathcal{B}|_S$ intersecting more than one S_i, but not intersecting every S_i. By Theorem 1.5.13, $S_1 \sqcup \cdots \sqcup S_k = S \in \mathcal{B}$. Therefore, we can write $S = S' \sqcup S''$, where $S', S'' \in \mathcal{B}$. Let S^1 be that of the sets S' and S'' which intersects more elements S_i than the other. Then S^1 intersects more than one S_i. If S^1 does not intersect all of S_i, then we are done. Otherwise we write $S^1 = S'^1 \sqcup S''^1$, where $S'^1, S''^1 \in \mathcal{B}$, and choose as S^2 that of the sets S'^1 and S''^1 which intersects more elements S_i than the other. If S^2 intersects all the sets S_i, choose S^3 in the same way, and so on. Since the cardinality of the sets $S^1, S^2, S^3, \ldots$ strictly decreases, at some point this process stops, and we get that one of the sets S'^i, S''^i intersects more than one S_i, but does not intersect every S_i. ☐

PROPOSITION 1.6.14 ([**188**]). *Let $\mathcal{B}$ be a building set on $[n+1]$ such that $P_\mathcal{B}$ is a flagtope. Then there exists a building set $\mathcal{B}_0 \subset \mathcal{B}$ such that $P_{\mathcal{B}_0}$ is a combinatorial cube with* $\dim P_{\mathcal{B}_0} = \dim P_\mathcal{B}$.

PROOF. By Proposition 1.5.2, we need to consider only connected building sets. For $n = 1$, the proposition is true. Assuming that the assertion holds for $m < n$, we shall prove it for $m = n$. By Proposition 1.6.13, we have $[n+1] = S' \sqcup S''$, where $S', S'' \in \mathcal{B}$. By the induction assumption, the building sets $\mathcal{B}|_{S'}$ and $\mathcal{B}|_{S''}$ have subsets $\mathcal{B}'_0$ and $\mathcal{B}''_0$ whose corresponding nestohedra are cubes. The building set $\mathcal{B}_0 = (\mathcal{B}'_0 \sqcup \mathcal{B}''_0) \cup [n+1]$ is the desired one (see Example 1.5.22). ☐

It now follows from Lemma 1.5.17 that a flag nestohedron can be obtained from a cube by a sequence of face truncations. The following lemma shows that there is a sequence consisting of truncations of faces of codimension two only:

LEMMA 1.6.15. *Let $\mathcal{B}_1 \subset \mathcal{B}_2$ be connected building sets on $[n+1]$ whose corresponding nestohedra $P_{\mathcal{B}_1}$ and $P_{\mathcal{B}_2}$ are flagtopes. Then*

(a) *$P_{\mathcal{B}_2}$ is obtained from $P_{\mathcal{B}_1}$ by a sequence of 2-truncations;*
(b) *$\gamma_i(P_{\mathcal{B}_1}) \leqslant \gamma_i(P_{\mathcal{B}_2})$ for $i = 0, 1, \ldots, [n/2]$.*

Furthermore, if $\mathcal{B}_1 \neq \mathcal{B}_2$, then the inequality above is strict for some i.

PROOF. Let S be a minimal (by inclusion) element of $\mathcal{B}_2 \setminus \mathcal{B}_1$. We set $\mathcal{B}' = \widehat{\mathcal{B}_1 \cup \{S\}}$ to be the minimal (by inclusion) building set containing $\mathcal{B}_1 \cup \{S\}$. By Proposition 1.6.13, there exist $S', S'' \in \mathcal{B}_2$ such that $S' \sqcup S'' = S$. It follows from the choice of S that $S', S'' \in \mathcal{B}_1$. It is easy to show that $\mathcal{B}'$ is the collection of sets $\mathcal{B}_1 \cup \{T = T' \sqcup T''\colon T', T'' \in \mathcal{B}_1, S' \subset T', S'' \subset T''\}$. Hence, the decomposition of any element of $\mathcal{B}' \setminus \mathcal{B}_1$ consists of two elements. Therefore, by Lemma 1.5.17, the nestohedron $P_{\mathcal{B}'}$ is obtained from $P_{\mathcal{B}_1}$ by a sequence of 2-truncations.

Since $B_1 \subsetneq \mathcal{B}' \subset \mathcal{B}_2$, we can finish the proof of (a) by induction on the number of elements in $\mathcal{B}_2 \setminus \mathcal{B}_1$. Statement (b) follows from Corollary 1.6.10. ☐

Finally we can prove the main result of this section, giving a characterisation of flag nestohedra:

THEOREM 1.6.16 ([**54**]). *A nestohedron $P_\mathcal{B}$ is a flagtope if and only if it is a 2-truncated cube.*

More precisely, if $P_\mathcal{B}$ is a flagtope, then there exists a sequence of building sets $\mathcal{B}_0 \subset \mathcal{B}_1 \subset \cdots \subset \mathcal{B}_N = \mathcal{B}$, where $P_{\mathcal{B}_0}$ is a combinatorial cube, $\mathcal{B}_i = \mathcal{B}_{i-1} \cup \{S_i\}$, and $P_{\mathcal{B}_i}$ is obtained from $P_{\mathcal{B}_{i-1}}$ by a 2-truncation of the face $F_{S_{j_1}} \cap F_{S_{j_2}} \subset P_{\mathcal{B}_{i-1}}$, where $S_i = S_{j_1} \sqcup S_{j_2}$, and $S_{j_1}, S_{j_2} \in \mathcal{B}_{i-1}$.

PROOF. By Proposition 1.5.23, we need to consider only connected building sets. The ‘only if’ statement follows from Proposition 1.6.14 and Lemma 1.6.15. To prove the ‘if’ statement we need to check that a polytope obtained from a flagtope by a 2-truncation is a flagtope. This is left as an exercise. ☐

Together with Proposition 1.6.12, Theorem 1.6.16 implies

COROLLARY 1.6.17. *The Gal conjecture holds for all flag nestohedra $P_\mathcal{B}$, i.e.* $\gamma_i(P_\mathcal{B}) \geqslant 0$.

EXAMPLE 1.6.18. Let us see how Theorem 1.6.16 works in the case of the 3-dimensional associahedron. The building set corresponding to As^3 is given by

$$\mathcal{B} = \bigl\{\{1\},\{2\},\{3\},\{4\},\{1,2\},\{2,3\},\{3,4\},\{1,2,3\},\{2,3,4\},\{1,2,3,4\}\bigr\}$$

(see Fig. 1.4, right). In order to obtain As^3 from I^3 by 2-truncations, we have to specify a building set $\mathcal{B}_0 \subset \mathcal{B}$, such that $P_{\mathcal{B}_0} \approx I^3$, and to order the elements of $\mathcal{B} \setminus \mathcal{B}_0$ in such a way that adding a new element to the building set corresponds to a 2-truncation.

First let $\mathcal{B}_0$ consist of $\{i\},\{1,2\},\{3,4\},[4]$. The associahedron $P_{\mathcal{B}}$ is then obtained from $P_{\mathcal{B}_0} \approx I^3$ by stepwise truncation of the faces $F_{\{1,2\}} \cap F_{\{3\}}, F_{\{2\}} \cap F_{\{3,4\}}, F_{\{2\}} \cap F_{\{3\}}$ in this order. (Warning: $P_{\mathcal{B}_0}$ is *not* the rectangular parallelepiped, it is a tetrahedron with two opposite edges truncated. However the three 2-truncations of $P_{\mathcal{B}_0}$ described here and the three 2-truncations of the rectangular parallelepiped described in Example 1.5.27 give the same (up to affine equivalence) polytope As^3 shown in Fig. 1.4, right.)

Another option is to let $\mathcal{B}_0$ consist of $\{i\},\{1,2\},\{1,2,3\},[4]$. Then $P_{\mathcal{B}_0}$ is obtained from a tetrahedron by truncating a vertex and then truncating an edge which contained this vertex; we have that $P_{\mathcal{B}_0} \approx I^3$. To obtain the associahedron $P_{\mathcal{B}}$ from $P_{\mathcal{B}_0}$ we first truncate the face $F_{\{2\}} \cap F_{\{3\}}$ of $P_{\mathcal{B}_0} \approx I^3$ and get the new facet $F_{\{2,3\}}$. Then we truncate the faces $F_{\{2,3\}} \cap F_{\{4\}}$ and $F_{\{3\}} \cap F_{\{4\}}$.

Exercises.

1.6.19. In a flagtope, a collection of pairwise intersecting faces (of any dimension) has a nonempty intersection.

1.6.20. A face of a flagtope is a flagtope. This generalises Proposition 1.6.3.

1.6.21. Give an example of a triangle-free simple n-polytope with $2n+3$ facets which is not a flagtope.

1.6.22. Give an example of a non-flag simple polytope P with $\gamma_i(P) \geqslant 0$.

1.6.23. A polytope obtained from a flagtope by a 2-truncation is a flagtope.

1.6.24 ([**54**, Theorem 9.1]). The face vectors of n-dimensional flag nestohedra $P_{\mathcal{B}}$ satisfy

(a) $\gamma_i(I^n) \leqslant \gamma_i(P_{\mathcal{B}}) \leqslant \gamma_i(Pe^n)$ for $i = 0,1,\dots,[n/2]$;
(b) $g_i(I^n) \leqslant g_i(P_{\mathcal{B}}) \leqslant g_i(Pe^n)$ for $i = 0,1,\dots,[n/2]$;
(c) $h_i(I^n) \leqslant h_i(P_{\mathcal{B}}) \leqslant h_i(Pe^n)$ for $i = 0,1,\dots,n$;
(d) $f_i(I^n) \leqslant f_i(P_{\mathcal{B}}) \leqslant f_i(Pe^n)$ for $i = 0,1,\dots,n$.

Furthermore, the lower bounds are achieved only for $P_{\mathcal{B}} = I^n$ and the upper bounds are achieved only for $P_{\mathcal{B}} = Pe^n$. (Hint: to prove (a) use Lemma 1.6.15. The other inequalities follow from Proposition 1.3.13.)

1.6.25 ([**54**, Theorem 9.2]). The face vectors of graph associahedra P_Γ corresponding to connected graphs Γ on $[n+1]$ satisfy

(a) $\gamma_i(As^n) \leqslant \gamma_i(P_\Gamma) \leqslant \gamma_i(Pe^n)$ for $i = 0,1,\dots,[n/2]$;
(b) $g_i(As^n) \leqslant g_i(P_\Gamma) \leqslant g_i(Pe^n)$ for $i = 0,1,\dots,[n/2]$;
(c) $h_i(As^n) \leqslant h_i(P_\Gamma) \leqslant h_i(Pe^n)$ for $i = 0,1,\dots,n$;
(d) $f_i(As^n) \leqslant f_i(P_\Gamma) \leqslant f_i(Pe^n)$ for $i = 0,1,\dots,n$.

Furthermore, the lower bounds are achieved only for $P_{\mathcal{B}} = As^n$ and the upper bounds are achieved only for $P_{\mathcal{B}} = Pe^n$.

1.7. Differential algebra of combinatorial polytopes

Here we develop a differential-algebraic formalism that will allow us to analyse the combinatorics of families of polytopes and their face numbers from the viewpoint of differential equations. All polytopes in this section are combinatorial.

Ring of polytopes. Denote by $\mathfrak{P}^{2n}$ the free abelian group generated by all combinatorial n-polytopes. The group $\mathfrak{P}^{2n}$ is infinitely generated for $n > 1$, and it splits into a direct sum of finitely generated components as follows:

$$\mathfrak{P}^{2n} = \bigoplus_{m \geqslant n+1} \mathfrak{P}^{2n,2(m-n)}$$

where $\mathfrak{P}^{2n,2(m-n)}$ is the group generated by n-polytopes with m facets. (The fact that there are finitely many combinatorial polytopes with a given number of facets is left as an exercise.)

Definition 1.7.1. The product of polytopes turns the direct sum

$$\mathfrak{P} = \bigoplus_{n \geqslant 0} \mathfrak{P}^{2n} = \mathfrak{P}^0 + \bigoplus_{m \geqslant 2} \bigoplus_{n=1}^{m-1} \mathfrak{P}^{2n,2(m-n)}$$

into a bigraded commutative ring, the *ring of polytopes*. The unit is P^0, a point.

Simple polytopes generate a bigraded subring of $\mathfrak{P}$, which we denote by $\mathfrak{S}$.

A polytope is *indecomposable* if it cannot be represented as a product of two other polytopes of positive dimension.

Proposition 1.7.2 ([**41**, Proposition 2.22]). *$\mathfrak{P}$ is a polynomial ring generated by indecomposable combinatorial polytopes.*

Proof. We need to show that any polytope P of positive dimension can be represented as a product of indecomposable polytopes, $P = P_1 \times \cdots \times P_k$, and this representation is unique up to reordering the factors. The existence of such a decomposition is clear. Now assume that $P_1 \times \cdots \times P_k \approx Q_1 \times \cdots \times Q_s$. Fix a vertex $v = v_1 \times \cdots \times v_k = w_1 \times \cdots \times w_s$, where $v_i \in P$ and $w_j \in Q_j$. The faces of the form $v_1 \times \cdots \times P_i \times \cdots \times v_k$ and $w_1 \times \cdots \times Q_j \times \cdots \times w_s$ are maximal indecomposable faces of P containing the vertex v, and therefore they bijectively correspond to each other under the combinatorial equivalence. It follows that $k = s$ and $P_i \approx Q_i$ for $i = 1, \ldots, k$ up to a permutation of factors.

Since $\mathfrak{P}$ is a free abelian group generated by combinatorial polytopes, and each polytope can be uniquely represented by a monomial in indecomposable polytopes, it follows that $\mathfrak{P}$ is a polynomial ring on indecomposable polytopes. $\square$

Given $P \in \mathfrak{P}^{2n}$, denote by $dP \in \mathfrak{P}^{2(n-1)}$ the sum of all facets of P in the ring $\mathfrak{P}$. The following lemma is straightforward.

Lemma 1.7.3. *$d\colon \mathfrak{P} \to \mathfrak{P}$ is a linear operator of degree -2 satisfying the identity*

$$d(P_1P_2) = (dP_1)P_2 + P_1(dP_2).$$

Therefore, $\mathfrak{P}$ is a differential ring, and $\mathfrak{S}$ is its differential subring.

Example 1.7.4. We have

$$dI^n = n(dI)I^{n-1} = 2nI^{n-1}, \quad \text{and} \quad d\Delta^n = (n+1)\Delta^{n-1}.$$

Face-polynomials revisited. By Proposition 1.3.6, the F-polynomial and the H-polynomial define ring homomorphisms

$$F\colon \mathfrak{P} \longrightarrow \mathbb{Z}[s,t], \quad H\colon \mathfrak{P} \longrightarrow \mathbb{Z}[s,t],$$

which send $P \in \mathfrak{P}$ to $F(P)(s,t)$ and $H(P)(s,t)$ respectively.

THEOREM 1.7.5. *For any simple polytope P we have*

$$F(dP) = \frac{\partial}{\partial t} F(P). \tag{1.30}$$

PROOF. Assume that P is a simple n-polytopse with facets $P_1, \ldots, P_m$. Then

$$F(dP) = \sum_{i=1}^{m} F(P_i) = \sum_{i=1}^{m} \sum_{k=0}^{n-1} f_k(P_i) s^k t^{n-1-k}.$$

On the other hand,

$$\frac{\partial}{\partial t} F(P) = \sum_{k=0}^{n-1} (n-k) f_k(P) s^k t^{n-1-k}.$$

Comparing the coefficients in the two sums above we reduce (1.30) to

$$\sum_{i=1}^{m} f_k(P_i) = (n-k) f_k(P). \tag{1.31}$$

Since P is simple, every k-face is contained in exactly $n-k$ facets, so it is counted $n-k$ times on the left hand side of the above identity. □

COROLLARY 1.7.6. *Let P_1 and P_2 be two simple n-polytopes such that $dP_1 = dP_2$ in $\mathfrak{S}$. Then $F(P_1) = F(P_2)$.*

PROOF. Indeed, $F(dP_1) = F(dP_2)$ implies that $\frac{\partial F(P_1)}{\partial t} = \frac{\partial F(P_2)}{\partial t}$. Using that $F(P)(s,0) = s^n$ we obtain $F(P_1) = F(P_2)$. □

PROPOSITION 1.7.7. *Let $\widetilde{F}\colon \mathfrak{S} \to \mathbb{Z}[s,t]$ be a linear map such that*

$$\widetilde{F}(dP) = \frac{\partial}{\partial t} \widetilde{F}(P) \quad \textit{and} \quad \widetilde{F}(P)|_{t=0} = s^n.$$

Then $\widetilde{F}(P) = F(P)$.

PROOF. We have $\widetilde{F}(P^0) = 1 = F(P^0)$. Assume by induction the statement is true in dimensions $\leqslant n-1$, and let P be a simple n-polytope. Then $\widetilde{F}(dP)(s,t) = F(dP)(s,t)$. It follows that $\frac{\partial}{\partial t}\widetilde{F}(P) = \frac{\partial}{\partial t}F(P)$. Therefore, $\widetilde{F}(P)(s,t) = F(P)(s,t) + C(s)$. Setting $t = 0$, we obtain $s^n = s^n + C(s)$, whence $C(s) = 0$. □

Theorem 1.7.5 also allows us to reduce the Dehn–Sommerville relations (1.13) to the Euler formula:

THEOREM 1.7.8. *The following identity holds for any simple n-polytope P:*

$$F(P)(s,t) = F(P)(-s, s+t). \tag{1.32}$$

Proof. We have $F(P^0)(s,t) = 1 = F(P^0)(-s,s+t)$. Assume by induction the identity holds in dimensions $\leqslant n-1$. Then for a given P of dimension n we have $F(dP)(s,t) = F(dP)(-s,s+t)$. Therefore, $\frac{\partial}{\partial t}F(P)(s,t) = \frac{\partial}{\partial t}F(P)(-s,s+t)$ and $F(P)(s,t) - F(P)(-s,s+t) = C(s)$. By Euler's formula (1.12),

$$F(-s,s) = \big(f_0 - f_1 + \cdots + (-1)^n f_n\big)s^n = s^n.$$

Therefore, $C(s) = F(P)(s,0) - F(P)(-s,s) = 0$. □

Example 1.7.9. For a simple 3-polytope P^3 with $m = f_2$ facets we have $f_1 = 3(m-2)$, $f_0 = 2(m-2)$. Assume that all facets are k-gons, so $dP^3 = mP^2_k$. Then by Theorem 1.7.5,

$$m(s^2 + kst + kt^2) = \frac{\partial}{\partial t}\big(s^3 + ms^2t + 3(m-2)st^2 + 2(m-2)t^3\big),$$

which implies $m(6-k) = 12$. It follows that the pair (k,m) may only take values $(3,4)$, $(4,6)$ and (5,12) corresponding to a simplex, cube and dodecahedron.

For general polytopes the difference $\delta = Fd - \frac{\partial}{\partial t}F$ measures the failure of identity (1.30).

A polytope is said to be *k-simple* (for $k \geqslant 0$) if each of its k-faces is an intersection of exactly $n-k$ facets. For example, a simple polytope is 0-simple, while 1-simple polytopes are also known as *simple in edges*. Every n-polytope is $(n-1)$- and $(n-2)$-simple.

A polytope is *k-simplicial* if each of its k-dimensional faces is a simplex. A simplicial n-polytope is $(n-1)$-simplicial, and every polytope is 1-simplicial. By polarity, if an n-polytope P is k-simple, then P^* is $(n-1-k)$-simplicial.

Theorem 1.7.10. *Let $P \in \mathfrak{P}$. Then the following identity holds:*

$$F(dP) = \frac{\partial}{\partial t}F(P) + \delta(P),$$

where $\delta\colon \mathfrak{P} \to \mathbb{Z}[s,t]$ is an F-derivation, i.e. it is linear and satisfies the identity

$$\delta(P_1P_2) = \delta(P_1)F(P_2) + F(P_1)\delta(P_2).$$

Moreover, if P is an n-polytope, then $\delta(P) = \delta_2 s^{n-3}t^2 + \cdots + \delta_{n-1}t^{n-1}$ and $\delta_i \geqslant 0$ for $2 \leqslant i \leqslant n-1$. Also, P is k-simple if and only if $\delta_{n-1-k} = 0$; in this case $\delta_i = 0$ for $2 \leqslant i \leqslant n-1-k$.

Proof. The fact that $\delta = Fd - \frac{\partial}{\partial t}F$ is an F-derivation is verified by a direct computation. By (1.31) the coefficient of $s^k t^{n-1-k}$ in $\delta(P)$ is given by

$$\delta_{n-1-k} = \sum_{i=1}^{m} f_k(P_i) - (n-k)f_k(P).$$

It vanishes for $k = n-1$ and $k = n-2$ (as each codimension-two face of P is contained in exactly two facets). Also, δ_{n-1-k} is non-negative because every k-face is contained in at least $n-k$ facets. Finally, if P is k-simple, then every j-face is contained in exactly $n-j$ facets for $j \geqslant k$, so δ_{n-1-j} vanishes for $j \geqslant k$. □

Example 1.7.11. Let P be a simple n-polytope, and P^* the polar simplicial polytope. Since the face poset of P^* is the opposite to the face poset of P,

$$F(P^*)(s,t) = s^n + t\,\frac{F(P)(t,s) - t^n}{s} = s^n + \sum_{k=0}^{n-1} f_k(P)s^{n-1-k}t^{k+1}\,.$$

We have $dP^* = f_0(P)\Delta^{n-1}$, therefore,

$$\delta(P^*) = f_0(P)\frac{(s+t)^n - t^n}{s} - \frac{\partial}{\partial t}\sum_{k=0}^{n} f_k(P)s^{n-1-k}t^{k+1}.$$

The coefficient of $s^{n-1-k}t^k$ on the right hand side above is given by

$$\delta_k(P^*) = \binom{n}{k} f_0(P) - (k+1)f_k(P), \quad \text{for } 1 \leqslant k \leqslant n.$$

This is non-negative by Theorem 1.7.10 or by Exercise 1.4.15.

The following properties of $H(P)$ follow from the corresponding properties of $F(P)$ established above and the identity $H(P)(s,t) = F(P)(s-t,t)$.

THEOREM 1.7.12.

(a) *The ring homomorphism* $H\colon \mathfrak{P} \longrightarrow \mathbb{Z}[s,t]$ *satisfies* $H(P)\big|_{t=0} = s^n$. *The restriction of* H *to the subring* $\mathfrak{S}$ *of simple polytopes satisfies the equation*

$$H(dP) = \partial H(P)$$

where $\partial = \frac{\partial}{\partial s} + \frac{\partial}{\partial t}$.

(b) *The image of* H *is generated by* $H(\Delta^1) = s + t$ *and* $H(\Delta^2) = s^2 + st + t^2$.

(c) *If* $\widetilde{H}\colon \mathfrak{S} \longrightarrow \mathbb{Z}[s,t]$ *is a linear map satisfying* $\widetilde{H}(dP) = \partial\widetilde{H}(P)$ *and* $\widetilde{H}(P)\big|_{t=0} = s^n$ *for any simple* n*-polytope* P, *then* $\widetilde{H}(P) = H(P)$.

Ring of building sets. Let $\mathcal{B}_1$ and $\mathcal{B}_2$ be building sets on $[n_1]$ and $[n_2]$ respectively. A *map* $f\colon \mathcal{B}_1 \to \mathcal{B}_2$ *of building sets* is a map $f\colon [n_1] \to [n_2]$ satisfying $f^{-1}(S) \in \mathcal{B}_1$ for every $S \in \mathcal{B}_2$. Two building sets $\mathcal{B}_1$ and $\mathcal{B}_2$ are said to be *equivalent*, if there exist maps $f\colon \mathcal{B}_1 \to \mathcal{B}_2$ and $g\colon \mathcal{B}_2 \to \mathcal{B}_1$ of building sets such that the compositions $f \circ g$ and $g \circ f$ are the identity maps.

The *product* of $\mathcal{B}_1$ and $\mathcal{B}_2$ is the building set $\mathcal{B}_1 \cdot \mathcal{B}_2$ on $[n_1 + n_2]$ induced by appending the interval $[n_2]$ to the interval $[n_1]$.

EXAMPLE 1.7.13. The collection $\mathcal{S}$ of Example 1.5.3 and the complete collection $\mathcal{C}$ are both connected building sets on $[n+1]$. They are initial and terminal connected building sets respectively, since for every connected building set $\mathcal{B}$ on $[n+1]$ there are maps $\mathcal{C} \to \mathcal{B} \to \mathcal{S}$ of building sets induced by the identity map on $[n+1]$.

DEFINITION 1.7.14. Denote by $\mathfrak{B}^{2n}$ the free abelian group generated by the equivalence classes of building sets on $[k]$ for $k \leqslant n+1$. Since $\mathcal{B}_1 \cdot \mathcal{B}_2$ is equivalent to $\mathcal{B}_2 \cdot \mathcal{B}_1$, the product turns $\mathfrak{B} = \bigoplus_{n \geqslant 0} \mathfrak{B}^{2n}$ into a commutative associative ring.

Given a connected building set $\mathcal{B}$ on $[n+1]$, define

$$d\mathcal{B} = \sum_{S \in \mathcal{B} \setminus [n+1]} \mathcal{B}|_S \cdot \mathcal{B}/S \tag{1.33}$$

where the sum is taken in the ring $\mathfrak{B}$. Since $\mathfrak{B}$ is generated by connected building sets, we may extend d to a linear map $d\colon \mathfrak{B} \to \mathfrak{B}$ using the Leibniz formula

$$d(\mathcal{B}_1 \cdot \mathcal{B}_2) = (d\mathcal{B}_1) \cdot \mathcal{B}_2 + \mathcal{B}_1 \cdot (d\mathcal{B}_2). \tag{1.34}$$

We therefore get a derivation of $\mathfrak{B}$.

Proposition 1.7.15. *The map $\mathcal{B} \mapsto P_{\mathcal{B}}$ induces a differential ring homomorphism $\beta\colon \mathfrak{B} \to \mathfrak{P}$. Its image is a graded subring with unit in $\mathfrak{P}$ multiplicatively generated by nestohedra $P_{\mathcal{B}}$ corresponding to connected building sets $\mathcal{B}$.*

Proof. The fact that β is a ring homomorphism is obvious. It follows from Corollary 1.5.12 and (1.34) that β commutes with the differentials. Every building set consisting only of singletons is mapped to P^0 which represents a unit in $\mathfrak{P}$. Finally, it follows from Proposition 1.5.2 (a) that the image $\beta(\mathfrak{B})$ is generated by $P_{\mathcal{B}}$ with connected $\mathcal{B}$. □

Remark. The ring $\mathfrak{B}$ does not have unit, and the map β is not injective.

Given a subset $S \subset [n+1]$, denote by $\Gamma|_S$ the restriction of Γ to the vertex set S, and denote by Γ/S the graph with the vertex set $[n+1]\backslash S$ having an edge between two vertices i and j whenever they are path connected in $\Gamma_{S\cup\{i,j\}}$.

Proposition 1.7.16. *For a connected graph Γ on $n+1$ vertices, we have*

$$dP_\Gamma = \sum_{\substack{S\subsetneq[n+1] \\ \Gamma|_S \text{ is connected}}} P(\Gamma|_S) \times P(\Gamma/S).$$

Proof. Follows directly from (1.33). □

Example 1.7.17. We have the following formulae for the differentials of the four graph-associahedra defined in Section 1.5:

$$\begin{aligned}
d\,As^n &= \sum_{i+j=n-1} (i+2)As^i \times As^j;\\
d\,Pe^n &= \sum_{i+j=n-1} \binom{n+1}{i+1} Pe^i \times Pe^j;\\
d\,Cy^n &= (n+1) \sum_{i+j=n-1} As^i \times Cy^j;\\
d\,St^n &= n \cdot St^{n-1} + \sum_{i=0}^{n-1} \binom{n}{i} St^i \times Pe^{n-i-1}.
\end{aligned}$$

For example (see Figs. 1.3–1.7),

$$\begin{aligned}
d\,As^3 &= 2As^0 \times As^2 + 3As^1 \times As^1 + 4As^2 \times As^0\ ;\\
d\,Pe^3 &= 4Pe^0 \times Pe^2 + 6Pe^1 \times Pe^1 + 4Pe^2 \times Pe^0\ ;\\
d\,Cy^3 &= 4(As^0 \times Cy^2 + As^1 \times Cy^1 + As^2 \times Cy^0)\ ;\\
d\,St^3 &= 3St^2 + St^0 \times Pe^2 + 3St^1 \times Pe^1 + 3St^2 \times Pe^0\ .
\end{aligned}$$

Exercises.

1.7.18. Show that there are finitely many combinatorial polytopes with a given number of facets.

1.7.19. Show that $\delta = Fd - \frac{\partial}{\partial t}F$ is an F-derivation.

1.8. Families of polytopes and differential equations

In this section, using the language of generating series, we interpret the formulae for the differential of nestohedra and graph-associahedra as certain partial differential equations. These differential equations encode the combinatorial information of the face structure of nestohedra. This exposition builds upon the results of [**40**] and [**38**].

Denote by $\mathfrak{P}[q]$ the polynomial ring in an indeterminate q with coefficients in $\mathfrak{P}$.

PROPOSITION 1.8.1. *Let*

$$Q\colon \mathfrak{P} \to \mathfrak{P}[q], \quad P \mapsto Q(P;q)$$

be a linear map such that

$$Q(dP;q) = \frac{\partial}{\partial q} Q(P;q) \quad \text{and} \quad Q(P;0) = P$$

for any polytope P. Then

$$Q(P;q) = \sum_{k=0}^{n} (d^k P) \frac{q^k}{k!}.$$

PROOF. Use induction by dimension, as in the proof of Proposition 1.7.7. □

Now assume given a sequence $\mathcal{P} = \{P^n,\ n \geqslant 0\}$ of polytopes, one in each dimension. We define its *generating series* as the formal power series

$$\mathcal{P}(x) = \sum_{n \geqslant 0} \lambda_n P^n x^{n+n_0}$$

in $\mathfrak{P} \otimes \mathbb{Q}[[x]]$. The parameter n_0 and the coefficients λ_n will be chosen depending on a particular sequence $\mathcal{P}$. Using the transformation Q of the previous proposition we may define the following 2-parameter extension of the generating series:

$$\mathcal{P}(q,x) = \sum_{n \geqslant 0} \lambda_n Q(P^n;q) x^{n+n_0}. \tag{1.35}$$

We have $\mathcal{P}(0,x) = \mathcal{P}(x)$.

We consider the following generating series of the six sequences of nestohedra:

$$\begin{aligned}
&\Delta(x) = \sum_{n \geqslant 0} \Delta^n \frac{x^{n+1}}{(n+1)!} \ ; \qquad && I(x) = \sum_{n \geqslant 0} I^n \frac{x^n}{n!} \ ; \\
&As(x) = \sum_{n \geqslant 0} As^n x^{n+2} \ ; && Pe(x) = \sum_{n \geqslant 0} Pe^n \frac{x^{n+1}}{(n+1)!} \ ; \\
&Cy(x) = \sum_{n \geqslant 0} Cy^n \frac{x^{n+1}}{n+1} \ ; && St(x) = \sum_{n \geqslant 0} St^n \frac{x^n}{n!} \ .
\end{aligned} \tag{1.36}$$

LEMMA 1.8.2. *The differentials of the generating series above are given by*

$$\begin{aligned}
&d\Delta(x) = x\Delta(x) \ ; \qquad && dI(x) = 2xI(x) \ ; \\
&d\,As(x) = As(x)\frac{d}{dx}As(x) \ ; && d\,Pe(x) = Pe^2(x) \ ; \\
&d\,Cy(x) = As(x)\frac{d}{dx}Cy(x) \ ; && d\,St(x) = \big(x + Pe(x)\big)St(x) \ .
\end{aligned}$$

PROOF. This follows from the formulae of Examples 1.7.4 and 1.7.17. □

THEOREM 1.8.3. *The two-parameter extensions of the generating series* (1.36) *satisfy the following partial differential equations:*

$$\frac{\partial}{\partial q}\Delta(q,x) = x\Delta(q,x) \;; \qquad \frac{\partial}{\partial q} I(q,x) = 2xI(q,x) \;;$$

$$\frac{\partial}{\partial q} As(q,x) = As(q,x)\frac{\partial}{\partial x} As(q,x) \;; \qquad \frac{\partial}{\partial q} Pe(q,x) = Pe^2(q,x) \;;$$

$$\frac{\partial}{\partial q} Cy(q,x) = As(q,x)\frac{\partial}{\partial x} Cy(q,x) \;; \qquad \frac{\partial}{\partial q} St(q,x) = \big(x + Pe(q,x)\big)St(q,x) \;.$$

PROOF. A direct calculation using formulae (1.35) and (1.36). □

REMARK. The role of the parameters λ_n in (1.35) can be illustrated as follows. If we replace the first series $\Delta(x)$ of (1.36) by

$$\widehat{\Delta}(x) = \sum_{n\geqslant 0} \Delta^n \frac{x^{n+1}}{n+1},$$

then the first equations of Lemma 1.8.2 and Theorem 1.8.3 take the form

$$d\widehat{\Delta}(x) = x^2\frac{d}{dx}\widehat{\Delta}(x), \quad \frac{\partial}{\partial q}\widehat{\Delta}(q,x) = x^2\frac{d}{dx}\widehat{\Delta}(q,x).$$

Four of the equations of Theorem 1.8.3, namely those corresponding to the series Δ, I, Pe and St, are ordinary differential equations. Their solutions are completely determined by the initial data $\mathcal{P}(0,x) = \mathcal{P}(x)$ and are given by the explicit formulae

$$\Delta(q,x) = \Delta(x)e^{qx} \;; \qquad I(q,x) = I(x)e^{2qx} \;;$$

$$Pe(q,x) = \frac{Pe(x)}{1-qPe(x)} \;; \qquad St(q,x) = St(x)\frac{e^{qx}}{1-qPe(x)} \;.$$

The equation for $U = As(q,x)$ has the form $U_q = UU_x$. This classical quasilinear partial differential equation was considered by E. Hopf, and therefore became known as the *Hopf equation*.

THEOREM 1.8.4.

(a) *The series $As(q,x)$ is given by the solution of the functional equation (*equation on characteristics*)*

$$As(q,x) = As(x + qAs(q,x)), \tag{1.37}$$

where $As(x) = As(0,x)$.

(b) *The series $Cy(q,x)$ is given by the solution of*

$$Cy(q,x) = Cy(x + qAs(q,x)), \tag{1.38}$$

where $Cy(x) = Cy(0,x)$.

PROOF. Set $U = As(q,x)$ and $As_x = \frac{d}{dx}As(x)$. If U is a solution to (1.37), then we obtain by differentiating

$$U_q = (U + qU_q)As_x, \quad U_x = (1 + qU_x)As_x.$$

Therefore, $(1 - qAs_x)U_q = UAs_x$ and $(1 - qAs_x)U_x = As_x$, which implies that U satisfies the Hopf equation $U_q = UU_x$. Its solution with the initial condition $U(0, x) = As(x)$ is unique by the general theory of quasilinear equations (in our case the uniqueness can be also verified using standard arguments with power series).

Similarly, by differentiating (1.38) we obtain for $V = Cy(q, x)$:

$$V_q = (U + qU_q)Cy_x, \quad V_x = (1 + qU_x)Cy_x.$$

Using that $U_q = UU_x$ we rewrite the first equation above as $V_q = U(1 + qU_x)Cy_x$, which implies $V_q = UV_x$ as claimed. This is exactly the equation for $V = Cy(q, x)$ given by Theorem 1.8.3, and its solution with $V(0, x) = Cy(x)$ is unique. $\square$

We can also use Lemma 1.8.2 to calculate the face-polynomials $F(s, t)$ of graph-associahedra. Let $\mathcal{P}(x)$ be one of the generating series (1.36), and set

$$F_{\mathcal{P}} = F(\mathcal{P}(x)) = \sum_{n\geqslant 0} \lambda^n x^{n+n_0} \sum_{k=0}^{n} f_k(P^n)s^k t^{n-k}.$$

We refer to $F_{\mathcal{P}} = F_{\mathcal{P}}(s, t; x)$ as the *generating series of face-polynomials*; it is a series in x whose coefficients are polynomials in s and t.

THEOREM 1.8.5. *The generating series of face-polynomials corresponding to* (1.36) *satisfy the following differential equations, with the initial conditions given in the second column:*

$$\begin{aligned}
&\frac{\partial}{\partial t} F_{\Delta} = xF_{\Delta}, && F_{\Delta}(s, 0; x) = \frac{e^{sx} - 1}{s}\ ;\\
&\frac{\partial}{\partial t} F_{I} = 2xF_{I}, && F_{I}(s, 0; x) = e^{sx}\ ;\\
&\frac{\partial}{\partial t} F_{As} = F_{As}\frac{\partial}{\partial x} F_{As}, && F_{As}(s, 0; x) = \frac{x^2}{1 - sx}\ ;\\
&\frac{\partial}{\partial t} F_{Pe} = F_{Pe}^2, && F_{Pe}(s, 0; x) = \frac{e^{sx} - 1}{s}\ ;\\
&\frac{\partial}{\partial t} F_{Cy} = F_{As}\frac{\partial}{\partial x} F_{Cy}, && F_{Cy}(s, 0; x) = -\frac{\ln(1 - sx)}{s}\ ;\\
&\frac{\partial}{\partial t} F_{St} = \big(x + F_{Pe}\big)F_{St}, && F_{St}(s, 0; x) = e^{sx}\ .
\end{aligned}$$

PROOF. The differential equations are obtained by applying F to the equations of Lemma 1.8.2 and using the fact that $F(dP) = \frac{\partial}{\partial t}F(P)$. The initial conditions follow by substituting s^n for P^n in (1.36) and calculating the resulting series. $\square$

Again, the four equations for the series F_{Δ}, F_I, F_{Pe} and F_{St} are ordinary differential equations. Their solutions are completely determined by the initial data; the explicit formulae are left as exercises. The remaining two partial differential equations for the series F_{As} and F_{Cy} can be explicitly solved as follows.

THEOREM 1.8.6.

(a) *The series $U = F_{As}$ satisfies the quadratic equation*

$$t(s + t)U^2 + (2tx + sx - 1)U + x^2 = 0. \tag{1.39}$$

The initial condition $F_{As}(s, 0; x) = \frac{x^2}{1-sx}$ determines its solution uniquely.

(b) *The series* F_{Cy} *is given by*

$$F_{Cy} = -\frac{1}{s}\ln\bigl(1 - s(x + tF_{As}))\bigr).$$

PROOF. (a) By analogy with (1.37) we show that $U = F_{As}$ satisfies

$$U = \varphi(x + tU),$$

where $\varphi(x) = F_{As}(s, 0; x) = \frac{x^2}{1-sx}$. It is equivalent to (1.39).

(b) We have that $V = F_{Cy}$ is given by the solution to $V_t = UV_x$. By analogy with (1.38) we show that it is given by

$$V = \psi(x + tU),$$

where $U = F_{As}$ and $\psi(x) = F_{Cy}(s, 0; x) = -\frac{\ln(1-sx)}{s}$. □

As a corollary, we shall derive a formula for the number of k-faces in As^n, which equals the number of bracketing of a word of $n+2$ letters with $n-k$ pairs of brackets. These numbers were first calculated by Cayley in 1891:

THEOREM 1.8.7. *The number of* k*-dimensional faces in an* n*-dimensional associahedron is given by*

$$f_k(As^n) = \frac{1}{n+2}\binom{n}{k}\binom{2n-k+2}{n+1}.$$

PROOF. We use the fact that F_{As} satisfies the Hopf equation, whose solutions may be obtained using conservation laws. Let

$$U(t, x) = \sum_{k\geqslant 0} U_k(x)t^k$$

be the solution of the Cauchy problem for the Hopf equation:

$$U_t = UU_x, \qquad U(0, x) = \varphi(x). \tag{1.40}$$

This equation has the following conservation laws

$$\left(\frac{U^k}{k}\right)_t = \left(\frac{U^{k+1}}{k+1}\right)_x, \quad \text{for } k \geqslant 1.$$

Hence,

$$\frac{d^k}{dt^k}U = \frac{d^{k-1}}{dt^{k-1}}\left(\frac{U^2}{2}\right)_x = \frac{d^{k-2}}{dt^{k-2}}\left(\frac{U^3}{3}\right)_{xx} = \cdots = \frac{d^k}{dx^k}\left(\frac{U^{k+1}}{k+1}\right), \quad \text{for } k \geqslant 1.$$

Evaluating at $t = 0$ we obtain

$$\left.\frac{d^k}{dt^k}U\right|_{t=0} = k!\,U_k(x) = \frac{d^k}{dx^k}\left(\frac{U_0^{k+1}(x)}{k+1}\right).$$

Therefore,

$$U_k(x) = \frac{1}{(k+1)!}\,\frac{d^k}{dx^k}\,\varphi^{k+1}(x).$$

By Theorem 1.8.5 the function

$$U = F_{As}(s, t; x) = \sum_{n\geqslant 0}\sum_{k=0}^{n} f_{n-k}(As^n)s^{n-k}t^k x^{n+2} \tag{1.41}$$

satisfies the Hopf equation (1.40) with the initial function $\varphi(s;x) = \frac{x^2}{1-sx}$. We therefore calculate

$$\begin{aligned}
U_k(s;x) &= \frac{1}{(k+1)!}\frac{d^k}{dx^k}\left(\frac{x^{2(k+1)}}{(1-sx)^{k+1}}\right) \\
&= \frac{1}{(k+1)!}\frac{d^k}{dx^k}\left(x^{2(k+1)}\sum_{l\geqslant 0}\binom{l+k}{l}s^l x^l\right) \\
&= \frac{1}{(k+1)!}\sum_{l\geqslant 0}\binom{l+k}{k}\frac{(2k+l+2)!}{(k+l+2)!}s^l x^{k+l+2} \\
&= \sum_{n\geqslant k}\frac{1}{n+2}\binom{n}{k}\binom{n+k+2}{n+1}s^{n-k}x^{n+2}.
\end{aligned}$$

On the other hand, it follows from (1.41) that

$$U_k(s;x) = \sum_{n\geqslant k} f_{n-k}(As^n)s^{n-k}x^{n+2}.$$

Comparing the last two formulae we obtain

$$f_{n-k}(As^n) = \frac{1}{n+2}\binom{n}{k}\binom{n+k+2}{n+1},$$

which is equivalent to the required formula. □

Exercises.

1.8.8. By solving the first two differential equations of Theorem 1.8.5 show that the generating series for the F-polynomials of simplices and cubes are given by

$$\begin{aligned}
F_\Delta(s,t;x) &= \sum_{n\geqslant 0} F(\Delta^n)\frac{x^{n+1}}{(n+1)!} = e^{tx}\frac{e^{sx}-1}{s}, \\
F_I(s,t;x) &= \sum_{n\geqslant 0} F(I^n)\frac{x^n}{n!} = e^{tx}e^{(s+t)x}.
\end{aligned}$$

Compare this with the formulae for $F(\Delta^n)$ and $F(I^n)$.

1.8.9. Show by solving the corresponding differential equations from Theorem 1.8.5 that the generating series for the face-polynomials of permutahedra and stellahedra are given by

$$\begin{aligned}
F_{Pe}(s,t;x) &= \sum_{n\geqslant 0} F(Pe^n)\frac{x^{n+1}}{(n+1)!} = \frac{e^{sx}-1}{s-t(e^{sx}-1)}, \\
F_{St}(s,t;x) &= \sum_{n\geqslant 0} F(St^n)\frac{x^n}{n!} = \frac{se^{(s+t)x}}{s-t(e^{sx}-1)}.
\end{aligned}$$

Compute the face-polynomials $F(Pe^n)$ and $F(St^n)$ and the face numbers explicitly.

1.8.10. Define the generating series $H_{\mathcal{P}}(s,t;x)$ of H-polynomials by analogy with the generating series of face-polynomials. Deduce the following formulae for sequences of nestohedra:

$$H_\Delta(s,t;x) = \sum_{n\geqslant 0} H(\Delta^n)\frac{x^{n+1}}{(n+1)!} = \frac{e^{sx}-e^{tx}}{s-t},$$

$$H_I(s,t;x) = \sum_{n\geqslant 0} H(I^n)\frac{x^n}{n!} = e^{(s+t)x},$$

$$H_{Pe}(s,t;x) = \sum_{n\geqslant 0} H(Pe^n)\frac{x^{n+1}}{(n+1)!} = \frac{e^{sx}-e^{tx}}{se^{tx}-te^{sx}},$$

$$H_{St}(s,t;x) = \sum_{n\geqslant 0} H(St^n)\frac{x^n}{n!} = \frac{(s-t)e^{(s+t)x}}{se^{tx}-te^{sx}}.$$

1.8.11. The series $Y = H_{As}(s,t;x) = \sum_{n\geqslant 0} H(As^n)x^{n+2}$ satisfies the quadratic equation

$$Y = (x+sY)(x+tY).$$

The initial condition $H_{As}(s,0;x) = \frac{x^2}{1-sx}$ determines its solution uniquely.

1.8.12. The components of the h-vector of As^n are given by

$$h_k = \frac{1}{n+1}\binom{n+1}{k}\binom{n+1}{k+1}, \qquad 0 \leqslant k \leqslant n.$$

CHAPTER 2

Combinatorial structures

The face poset of a convex polytope is a classical example of a combinatorial structure underlying a decomposition of a geometric object. With the development of combinatorial topology several new combinatorial structures emerged, such as simplicial and cubical complexes, simplicial posets and other types of regular cellular (or CW) complexes. Many of these structures have eventually become objects of independent study in geometric combinatorics.

A simplicial complex is the abstract combinatorial structure behind a simplicial subdivision (or *triangulation*) of a topological space. Triangulations were first introduced by Poincaré and provide a rigorous and convenient tool for studying topological invariants of smooth manifolds by combinatorial methods. The notion of a *nerve* of a covering of a topological space, introduced by Alexandroff, provides another source of examples of simplicial complexes.

The study of triangulations stimulated the development of first combinatorial and then algebraic topology in the first half of the XXth century. With the appearance of cellular complexes algebraic tools gradually replaced the combinatorial ones in mainstream topology. Simplicial complexes still play a pivotal role in PL (piecewise linear) topology, however nowadays the main source of interest in them is in discrete and computational geometry. One reason for that is the emergence of computers, since simplicial complexes provide the most effective way to translate geometric and topological structures into machine language.

We therefore may distinguish two different views on the role of simplicial complexes and triangulations. In topology, simplicial complexes and their different derivatives such as singular chains and simplicial sets are used as technical tools to study the topology of the underlying space. Most combinatorial invariants of nerves or triangulations (such as the number of faces of a given dimension) do not have meaning in topology, as they do not reflect any topological feature of the underlying space. Topologists therefore tend not to distinguish between simplicial complexes that have the same underlying topology. For instance, refining a triangulation (such as passing to the barycentric subdivision) changes the combinatorics drastically, but does not affect the underlying topology. On the other hand, in combinatorial geometry the combinatorics of a simplicial complex is what really matters, while the underlying topology is often simple or irrelevant.

In toric topology the combinatorist's point of view on triangulations and similar decompositions is enriched by elaborate topological techniques. Combinatorial invariants of triangulations therefore can be analysed by topological methods, and at the same time combinatorial structures such as simplicial complexes or posets become a source of examples of topological spaces and manifolds with nice features and lots of symmetry, e.g. bearing a torus action. The combinatorial structures are the subject of this chapter; the associated topological objects will come later.

Here we assume only minimal knowledge of topology. The reader may also check Appendix B for the definition of simplicial homology groups, etc.

2.1. Polyhedral fans

Like convex polytopes, polyhedral fans encode both geometrical and combinatorial information. The geometry of fans is poorer than that of convex polytopes, but this geometry is still a part of the structure of a fan, which distinguishes fans from the purely combinatorial objects considered later in this chapter.

Although fans were considered in convex geometry independently, the main source of interest to them is in the theory of *toric varieties*, which are classified by rational fans. Toric varieties are the subject of Chapter 5, and here we describe the terminology and constructions related to fans.

Definition 2.1.1. A set of vectors $\boldsymbol{a}_1, \ldots, \boldsymbol{a}_k \in \mathbb{R}^n$ defines a *convex polyhedral cone*, or simply *cone*,

$$\sigma = \{\mu_1 \boldsymbol{a}_1 + \cdots + \mu_k \boldsymbol{a}_k \colon \mu_i \in \mathbb{R}_{\geqslant}\}.$$

Here $\boldsymbol{a}_1, \ldots \boldsymbol{a}_k$ are referred to as *generating vectors* (or *generators*) of σ. A cone is *rational* if its generators can be chosen from the integer lattice $\mathbb{Z}^n \subset \mathbb{R}^n$. If σ is a rational cone, then its generators $\boldsymbol{a}_1, \ldots \boldsymbol{a}_k$ are usually chosen to be *primitive*, i.e. each $\boldsymbol{a}_i$ is the smallest lattice vector in the ray defined by it.

A cone is *strongly convex* if it does not contain a line. A strongly convex cone is *simplicial* if it is generated by a part of basis of $\mathbb{R}^n$, and is *regular* if it is generated by a part of basis of $\mathbb{Z}^n$. (Regular cones play a special role in the theory of toric varieties, see Chapter 5.)

A cone is also an (unbounded) intersection of finitely many halfspaces in $\mathbb{R}^n$, so it is a convex polyhedron in the sense of Definition 1.1.2. We therefore may define face of a cone in the same way as we did for polytopes, as intersections of σ with supporting hyperplanes. We can only consider supporting hyperplanes containing $\mathbf{0}$; such a hyperplane is defined by a linear function $\boldsymbol{u}$ and will be denoted by $\boldsymbol{u}^\perp$. A *face* τ of a cone σ is therefore an intersection of σ with a supporting hyperplane $\boldsymbol{u}^\perp$, i.e. $\tau = \sigma \cap \boldsymbol{u}^\perp$. Every face of a cone is itself a cone. If $\sigma \neq \mathbb{R}^n$, then σ has the smallest face $\sigma \cap (-\sigma)$; it is a vertex $\mathbf{0}$ if and only if σ strongly convex. Any cone is generated by a set of nonzero vectors along its edges.

A *fan* is a finite collection $\Sigma = \{\sigma_1, \ldots, \sigma_s\}$ of strongly convex cones in some $\mathbb{R}^n$ such that every face of a cone in Σ belongs to Σ and the intersection of any two cones in Σ is a face of each. A fan Σ is *rational* (respectively, *simplicial*, *regular*) if every cone in Σ is rational (respectively, simplicial, regular). A fan $\Sigma = \{\sigma_1, \ldots, \sigma_s\}$ is called *complete* if $\sigma_1 \cup \cdots \cup \sigma_s = \mathbb{R}^n$.

Given a cone $\sigma \subset \mathbb{R}^n$, define its *dual* as

$$\check{\sigma} = \bigl\{\boldsymbol{x} \in \mathbb{R}^n \colon \langle \boldsymbol{u}, \boldsymbol{x} \rangle \geqslant 0 \text{ for all } \boldsymbol{u} \in \sigma\bigr\}. \tag{2.1}$$

(Note the difference with the definition of the polar set of a polyhedron, see (1.8).) Observe that if $\boldsymbol{u} \in \check{\sigma}$, then $\boldsymbol{u}^\perp$ is a supporting hyperplane of σ. It can be shown that $\check{\sigma}$ is indeed a cone, $(\check{\sigma})\check{} = \sigma$, and $\check{\sigma}$ is strongly convex if and only if $\dim \sigma = n$.

Cones in a fan can be separated by hyperplanes (or linear functions):

LEMMA 2.1.2 (Separation Lemma). *Let σ and σ' be two cones whose intersection τ is a face of each. Then there exists $\boldsymbol{u} \in \check{\sigma} \cap (-\sigma')\check{}$ such that*

$$\tau = \sigma \cap \boldsymbol{u}^{\perp} = \sigma' \cap \boldsymbol{u}^{\perp}.$$

In other words, a supporting hyperplane defining τ can be chosen so as to separate σ and σ'.

PROOF. We only sketch a proof; the details can be found, e.g. in [**96**, §1.2]. The fact that σ and σ' intersect in a face implies that the cone $\xi = \sigma - \sigma' = \sigma + (-\sigma')$ is not the whole space. Let $\boldsymbol{u}^{\perp}$ be a supporting hyperplane defining the smallest face of ξ, that is,

$$\xi \cap \boldsymbol{u}^{\perp} = \xi \cap (-\xi) = (\sigma - \sigma') \cap (\sigma' - \sigma).$$

We claim that this $\boldsymbol{u}$ has the required properties. Indeed, $\sigma \subset \xi$ implies $\boldsymbol{u} \in \check{\sigma}$, and $\tau \subset \xi \cap (-\xi)$ implies $\tau \subset \sigma \cap \boldsymbol{u}^{\perp}$. Conversely, if $\boldsymbol{x} \in \sigma \cap \boldsymbol{u}^{\perp}$, then $\boldsymbol{x}$ is in $\sigma' - \sigma$, so that $\boldsymbol{x} = \boldsymbol{y}' - \boldsymbol{y}$ for $\boldsymbol{y}' \in \sigma'$, $\boldsymbol{y} \in \sigma$. Then $\boldsymbol{x} + \boldsymbol{y} \in \sigma \cap \sigma' = \tau$, which implies that both $\boldsymbol{x}$ and $\boldsymbol{y}$ are in τ. Hence $\sigma \cap \boldsymbol{u}^{\perp} = \tau$, and the same argument for $-\boldsymbol{u}$ shows that $\boldsymbol{u} \in (-\sigma')\check{}$ and $\sigma' \cap \boldsymbol{u}^{\perp} = \tau$. $\square$

Miraculously, the convex-geometrical separation property above will translate into topological separation (Hausdorffness) of algebraic varieties and topological spaces constructed from fans in the latter chapters.

The next construction assigns a complete fan to every convex polytope.

CONSTRUCTION 2.1.3 (Normal fan). Let P be a polytope (1.1) with m facets $F_1, \ldots, F_m$ and normal vectors $\boldsymbol{a}_1, \ldots, \boldsymbol{a}_m$. Given a face $Q \subset P$ define the cone

$$\sigma_Q = \{\boldsymbol{u} \in \mathbb{R}^n \colon \langle \boldsymbol{u}, \boldsymbol{x}' \rangle \leqslant \langle \boldsymbol{u}, \boldsymbol{x} \rangle \text{ for all } \boldsymbol{x}' \in Q \text{ and } \boldsymbol{x} \in P\}. \tag{2.2}$$

The dual cone $\check{\sigma_Q}$ is generated by vectors $\boldsymbol{x} - \boldsymbol{x}'$ where $\boldsymbol{x} \in P$ and $\boldsymbol{x}' \in Q$. In other words, $\check{\sigma_Q}$ is the 'polyhedral angle' at the face Q consisting of all vectors pointing from points of Q to points of P.

We say that a vector $\boldsymbol{a}_i$ is *normal* to the face Q if $Q \subset F_i$. The cone σ_Q is generated by those $\boldsymbol{a}_i$ which are normal to Q (this is an exercise). Then

$$\Sigma_P = \{\sigma_Q \colon Q \text{ is a face of } P\}$$

is a complete fan Σ_P in $\mathbb{R}^n$ (this is another exercise), which is denoted by Σ_P and is referred to as the *normal fan* of the polytope P. If $\mathbf{0}$ is contained in the interior of P then Σ_P may be also described as the set of cones over the faces of the polar polytope P^* (yet another exercise).

It is clear from the above descriptions that the normal fan Σ_P is simplicial if and only if P is simple. In this case the definition of Σ_P may be simplified: the cones of Σ_P are generated by those sets $\{\boldsymbol{a}_{i_1}, \ldots, \boldsymbol{a}_{i_k}\}$ for which the intersection $F_{i_1} \cap \cdots \cap F_{i_k}$ is nonempty.

If all vertices of P are in the lattice $\mathbb{Z}^n$ then the normal fan Σ_P is rational, but the converse is not true. Polytopes whose normal fans are regular are called *Delzant* (this name comes from a symplectic geometry construction discussed in Section 5.5). Therefore, P is a Delzant polytope if and only if for every vertex $\boldsymbol{x} \in P$ the normal vectors to the facets meeting at $\boldsymbol{x}$ can be chosen to form a basis of $\mathbb{Z}^n$. In this definition one may replace 'normal vectors to the facets meeting at $\boldsymbol{x}$' by 'vectors along the edges meeting at $\boldsymbol{x}$'. A Delzant polytope is necessarily simple.

The normal fan Σ_P of a polytope P contains the information about the normals to the facets (the generators $\boldsymbol{a}_i$ of the edges of Σ_P) and the combinatorial structure of P (which sets of vectors $\boldsymbol{a}_i$ span a cone of Σ_P is determined by which facets intersect at a face), however the scalars b_i in (1.1) are lost. Not any fan can be obtained by 'forgetting the numbers b_i' from a presentation of a polytope by inequalities, i.e. not any fan is a normal fan of a polytope. This is fails even for regular fans, as is shown by the next example, which is taken from [**96**].

EXAMPLE 2.1.4. Consider the complete three-dimensional fan Σ with 7 edges generated by the vectors $\boldsymbol{a}_1 = \boldsymbol{e}_1$, $\boldsymbol{a}_2 = \boldsymbol{e}_2$, $\boldsymbol{a}_3 = \boldsymbol{e}_3$, $\boldsymbol{a}_4 = -\boldsymbol{e}_1 - \boldsymbol{e}_2 - \boldsymbol{e}_3$, $\boldsymbol{a}_5 = -\boldsymbol{e}_1 - \boldsymbol{e}_2$, $\boldsymbol{a}_6 = -\boldsymbol{e}_2 - \boldsymbol{e}_3$, $\boldsymbol{a}_7 = -\boldsymbol{e}_1 - \boldsymbol{e}_3$, and 10 three-dimensional cones with vertex $\mathbf{0}$ over the faces of the triangulated boundary of the tetrahedron shown in Fig. 2.1. It is easy to verify that Σ is regular.



FIGURE 2.1. Complete regular fan not coming from a polytope.

Assume that $\Sigma = \Sigma_P$ is the normal fan of a polytope P. Consider the function $\psi\colon \mathbb{R}^n \to \mathbb{R}$ given by

$$\psi(\boldsymbol{u}) = \min_{\boldsymbol{x}\in P}\langle \boldsymbol{u}, \boldsymbol{x}\rangle = \min_{\boldsymbol{v}\in V(P)}\langle \boldsymbol{u}, \boldsymbol{v}\rangle,$$

where $V(P)$ is the set of vertices of P. This function is continuous and its restriction to every 3-dimensional cone of Σ_P is linear. Indeed, 3-dimensional cones $\sigma_{\boldsymbol{v}}$ correspond to vertices $\boldsymbol{v} \in V(P)$, and we have $\psi(\boldsymbol{u}) = \langle \boldsymbol{u}, \boldsymbol{v}\rangle$ for $\boldsymbol{u} \in \sigma_{\boldsymbol{v}}$ by definition (2.2) of $\sigma_{\boldsymbol{v}}$.

Now consider the two 3-dimensional cones of Σ_P generated by the triples $\boldsymbol{a}_1, \boldsymbol{a}_3, \boldsymbol{a}_5$ and $\boldsymbol{a}_1, \boldsymbol{a}_5, \boldsymbol{a}_6$, and let $\boldsymbol{v}$ and $\boldsymbol{v}'$ be the corresponding vertices of P. Then $\psi(\boldsymbol{a}_1) = \langle \boldsymbol{a}_1, \boldsymbol{v}\rangle$, $\psi(\boldsymbol{a}_3) = \langle \boldsymbol{a}_3, \boldsymbol{v}\rangle$, $\psi(\boldsymbol{a}_5) = \langle \boldsymbol{a}_5, \boldsymbol{v}\rangle$, $\psi(\boldsymbol{a}_6) = \langle \boldsymbol{a}_6, \boldsymbol{v}'\rangle$, hence,

$$\psi(\boldsymbol{a}_1) + \psi(\boldsymbol{a}_5) - \psi(\boldsymbol{a}_3) = \langle \boldsymbol{a}_1 + \boldsymbol{a}_5 - \boldsymbol{a}_3, \boldsymbol{v}\rangle = \langle \boldsymbol{a}_6, \boldsymbol{v}\rangle > \langle \boldsymbol{a}_6, \boldsymbol{v}'\rangle = \psi(\boldsymbol{a}_6).$$

Therefore,

$$\psi(\boldsymbol{a}_1) + \psi(\boldsymbol{a}_5) > \psi(\boldsymbol{a}_3) + \psi(\boldsymbol{a}_6).$$

Similarly,

$$\psi(\boldsymbol{a}_2) + \psi(\boldsymbol{a}_6) > \psi(\boldsymbol{a}_1) + \psi(\boldsymbol{a}_7),$$

$$\psi(\boldsymbol{a}_3) + \psi(\boldsymbol{a}_7) > \psi(\boldsymbol{a}_2) + \psi(\boldsymbol{a}_5).$$

Adding the last three inequalities together we get a contradiction.

Exercises.

2.1.5. Let σ be a cone in $\mathbb{R}^n$. Show that $\sigma^\vee$ is also a cone, $(\sigma^\vee)^\vee = \sigma$, and $\sigma^\vee$ is strongly convex if and only if $\dim \sigma = n$.

2.1.6. The cone σ_Q given by (2.2) is generated by those vectors among $\boldsymbol{a}_1, \ldots, \boldsymbol{a}_m$ which are normal to Q.

2.1.7. The set $\{\sigma_Q \colon Q \text{ is a face of } P\}$ is a complete fan in $\mathbb{R}^n$.

2.1.8. If $\mathbf{0}$ is contained in the interior of P then Σ_P consists of cones over the faces of the polar polytope P^*.

2.1.9. Let P be a convex polytope (not necessarily simple). Does the collection of cones generated by the sets $\{\boldsymbol{a}_{i_1}, \ldots, \boldsymbol{a}_{i_k}\}$ of normal vectors for which $F_{i_1} \cap \cdots \cap F_{i_k} \neq \varnothing$ form a fan?

2.2. Simplicial complexes

A simplex is the convex hull of a set of affinely independent points in $\mathbb{R}^n$.

Definition 2.2.1. A *geometric simplicial complex* in $\mathbb{R}^n$ is a collection $\mathcal{P}$ of simplices of arbitrary dimension such that every face of a simplex in $\mathcal{P}$ belongs to $\mathcal{P}$ and the intersection of any two simplices in $\mathcal{P}$ is either empty or a face of each.

To make the exposition more streamlined and without creating much ambiguity we shall not distinguish between the collection $\mathcal{P}$ (which is an abstract set of simplices) and the union of its simplices (which is a subset in $\mathbb{R}^n$). In PL (piecewise linear) topology the latter union is usually referred to as 'the *polyhedron* of $\mathcal{P}$'. Although we have already reserved the term 'polyhedron' for a finite intersection of halfspaces (1.1), we shall also occasionally use it in the PL topological sense (when it creates no ambiguity).

A *face* of $\mathcal{P}$ is a face of any of its simplices. The dimension of $\mathcal{P}$ is the maximal dimension of its faces.

If we know the set of vertices of $\mathcal{P}$ in $\mathbb{R}^n$ then we may recover the whole $\mathcal{P}$ by specifying which subsets of vertices span simplices. This observation leads to the following definition.

Definition 2.2.2. An *abstract simplicial complex* on a set $\mathcal{V}$ is a collection $\mathcal{K}$ of subsets $I \subset \mathcal{V}$ such that if $I \in \mathcal{K}$ then any subset of I also belongs to $\mathcal{K}$. We always assume that the empty set $\varnothing$ belongs to $\mathcal{K}$. We refer to $I \in \mathcal{K}$ as an (abstract) *simplex* of $\mathcal{K}$.

One-element simplices are called *vertices* of $\mathcal{K}$. If $\mathcal{K}$ contains all one-element subsets of $\mathcal{V}$, then we say that $\mathcal{K}$ is a simplicial complex *on the vertex set* $\mathcal{V}$.

It is sometimes convenient to consider simplicial complexes $\mathcal{K}$ whose vertex sets are proper subsets of $\mathcal{V}$. In this case we refer to a one-element subset of $\mathcal{V}$ which is not a vertex of $\mathcal{K}$ as a *ghost vertex*.

The *dimension* of a simplex $I \in \mathcal{K}$ is $\dim I = |I| - 1$, where $|I|$ denotes the number of elements in I. A simplicial complex $\mathcal{K}$ is *pure* if all its maximal simplices have the same dimension. A subcollection $\mathcal{K}' \subset \mathcal{K}$ which is itself a simplicial complex is called a *subcomplex* of $\mathcal{K}$.

A geometric simplicial complex $\mathcal{P}$ is said to be a *geometric realisation* of an abstract simplicial complex $\mathcal{K}$ on a set $\mathcal{V}$ if there is a bijection between $\mathcal{V}$ and the vertex set of $\mathcal{P}$ which maps abstract simplices of $\mathcal{K}$ to vertex sets of faces of $\mathcal{P}$.

Both geometric and abstract simplicial complexes will be assumed to be *finite*, unless we explicitly specify otherwise. In most constructions we identify the set $\mathcal{V}$ with the index set $[m] = \{1, \ldots, m\}$ and consider abstract simplicial complexes on $[m]$. An identification of $\mathcal{V}$ with $[m]$ fixes an order of vertices, although this order will be irrelevant in most cases. We drop the parentheses in the notation of one-element subsets $\{i\} \subset [m]$, so that $i \in \mathcal{K}$ means that $\{i\}$ is a vertex of $\mathcal{K}$, and $i \notin \mathcal{K}$ means that $\{i\}$ is a ghost vertex.

We shall use the common notation Δ^{m-1} for any of the following three objects: an $(m-1)$-simplex (a convex polytope), the geometric simplicial complex consisting of all faces in an $(m-1)$-simplex, and the abstract simplicial complex consisting of all subsets of $[m]$.

Construction 2.2.3. Every abstract simplicial complex $\mathcal{K}$ on $[m]$ can be realised geometrically in $\mathbb{R}^m$ as follows. Let $\boldsymbol{e}_1, \ldots, \boldsymbol{e}_m$ be the standard basis in $\mathbb{R}^m$, and for each $I \subset [m]$ denote by Δ^I the convex hull of points $\boldsymbol{e}_i$ with $i \in I$. Then

$$\bigcup_{I \in \mathcal{K}} \Delta^I \subset \mathbb{R}^m$$

is a geometric realisation of $\mathcal{K}$.

The above construction is just a geometrical interpretation of the fact that any simplicial complex on $[m]$ is a subcomplex of the simplex Δ^{m-1}. Also, by a classical result [**185**], a d-dimensional abstract simplicial complex admits a geometric realisation in $\mathbb{R}^{2d+1}$.

Example 2.2.4. The boundary of a simplicial n-polytope is a simplicial complex of dimension $n-1$. For a simple polytope P, we shall denote by $\mathcal{K}_P$ the boundary complex ∂P^* of the dual polytope. It coincides with the nerve of the covering of ∂P by the facets. That is, the vertices of $\mathcal{K}_P$ are the facets of P, and a set of vertices spans a simplex whenever the intersection of the corresponding facets is nonempty. We refer to $\mathcal{K}_P$ as the *nerve complex* of P.

Definition 2.2.5. The *f-vector* of an $(n-1)$-dimensional simplicial complex $\mathcal{K}$ is $\boldsymbol{f}(\mathcal{K}) = (f_0, f_1, \ldots, f_{n-1})$, where f_i is the number of i-dimensional simplices in $\mathcal{K}$. We also set $f_{-1} = 1$; to justify this convention one can assign dimension -1 to the empty simplex. The *h-vector* $\boldsymbol{h}(\mathcal{K}) = (h_0, h_1, \ldots, h_n)$ is defined by the identity

$$h_0 s^n + h_1 s^{n-1} + \cdots + h_n = (s-1)^n + f_0 (s-1)^{n-1} + \cdots + f_{n-1}. \tag{2.3}$$

(Warning: this is not the identity obtained by substituting $t = 1$ in (1.10).) The *g-vector* $\boldsymbol{g}(\mathcal{K}) = (g_0, g_1, \ldots, g_{[n/2]})$ is defined by $g_0 = 1$ and $g_i = h_i - h_{i-1}$ for $i = 1, \ldots, [n/2]$.

Remark. If $\mathcal{K} = \mathcal{K}_P$ is the nerve complex of a simple polytope P, then $\boldsymbol{f}(\mathcal{K}) = \boldsymbol{f}(P^*)$ and $\boldsymbol{h}(\mathcal{K}) = \boldsymbol{h}(P)$. This our notational convention may look a bit artificial, but it seems to be the best possible way to treat the face vectors of both polytopes and simplicial complexes consistently.

Definition 2.2.6. Let $\mathcal{K}_1$, $\mathcal{K}_2$ be simplicial complexes on the sets $[m_1]$, $[m_2]$ respectively, and $\mathcal{P}_1$, $\mathcal{P}_2$ their geometric realisations. A map $\varphi\colon [m_1] \to [m_2]$ induces a *simplicial map* between $\mathcal{K}_1$ and $\mathcal{K}_2$ if $\varphi(I) \in \mathcal{K}_2$ for any $I \in \mathcal{K}_1$. A simplicial map φ is said to be *nondegenerate* if $|\varphi(I)| = |I|$ for any $I \in \mathcal{K}_1$. On the geometric level, a simplicial map extends linearly on the faces of $\mathcal{P}_1$ to a map $\mathcal{P}_1 \to \mathcal{P}_2$, which we continue to denote by φ. A *simplicial isomorphism* is a simplicial map for which there exists a simplicial inverse.

There is an obvious isomorphism between any two geometric realisations of an abstract simplicial complex $\mathcal{K}$. We therefore shall use the common notation $|\mathcal{K}|$ for any geometric realisation of $\mathcal{K}$. Whenever it is safe, we shall not distinguish between abstract simplicial complexes and their geometric realisations. For example, we shall say 'simplicial complex $\mathcal{K}$ is homeomorphic to X' instead of 'the geometric realisation of $\mathcal{K}$ is homeomorphic to X'.

We shall refer to a simplicial complex homeomorphic to a topological space X as a *triangulation of* X, or a *simplicial subdivision of* X.

A *subdivision* of a geometric simplicial complex $\mathcal{P}$ is a geometric simplicial complex $\mathcal{P}'$ such that each simplex of $\mathcal{P}'$ is contained in a simplex of $\mathcal{P}$ and each simplex of $\mathcal{P}$ is a union of finitely many simplices of $\mathcal{P}'$. A *PL map* $\varphi\colon \mathcal{P}_1 \to \mathcal{P}_2$ is a simplicial map from a subdivision of $\mathcal{P}_1$ to a subdivision of $\mathcal{P}_2$. A *PL homeomorphism* is a PL map for which there exists a PL inverse. Two PL homeomorphic simplicial complexes are also referred to as *combinatorially equivalent*. In other words, two simplicial complexes are PL homeomorphic if there is a simplicial complex isomorphic to a subdivision of each of them.

Remark. For a topological approach to PL maps (where a PL map is defined between spaces rather than their triangulations) we refer to standard sources on PL topology, such as [**125**] and [**197**].

Example 2.2.7.

1. If P is a simple n-polytope then the nerve complex $\mathcal{K}_P$ (see Example 2.2.4) is a triangulation of an $(n-1)$-dimensional sphere S^{n-1}.

2. Let Σ be a simplicial fan in $\mathbb{R}^n$ with m edges generated by vectors $\boldsymbol{a}_1, \ldots, \boldsymbol{a}_m$. Its *underlying simplicial complex* is defined by

$$\mathcal{K}_\Sigma = \bigl\{\{i_1, \ldots, i_k\} \subset [m]\colon \boldsymbol{a}_{i_1}, \ldots, \boldsymbol{a}_{i_k} \text{ span a cone of } \Sigma\bigr\}.$$

Informally, $\mathcal{K}_\Sigma$ may be viewed as the intersection of Σ with a unit sphere. The fan Σ is complete if and only if $\mathcal{K}_\Sigma$ is a triangulation of S^{n-1}. If Σ is a normal fan of a simple n-polytope P, then $\mathcal{K}_\Sigma = \mathcal{K}_P$.

Construction 2.2.8 (join). Let $\mathcal{K}_1$ and $\mathcal{K}_2$ be simplicial complexes on sets $\mathcal{V}_1$ and $\mathcal{V}_2$ respectively. The *join* of $\mathcal{K}_1$ and $\mathcal{K}_2$ is the simplicial complex

$$\mathcal{K}_1 * \mathcal{K}_2 = \bigl\{I \subset \mathcal{V}_1 \sqcup \mathcal{V}_2\colon I = I_1 \cup I_2,\ I_1 \in \mathcal{K}_1, I_2 \in \mathcal{K}_2\bigr\}$$

on the set $\mathcal{V}_1 \sqcup \mathcal{V}_2$. The join operation is associative by inspection.

Example 2.2.9.

1. If $\mathcal{K}_1 = \Delta^{m_1-1}$, $\mathcal{K}_2 = \Delta^{m_2-1}$, then $\mathcal{K}_1 * \mathcal{K}_2 = \Delta^{m_1+m_2-1}$.

2. The simplicial complex $\Delta^0 * \mathcal{K}$ (the join of $\mathcal{K}$ and a point) is called the *cone* over $\mathcal{K}$ and denoted $\operatorname{cone}\mathcal{K}$.

3. Let S^0 be a pair of disjoint points (a 0-sphere). Then $S^0 * \mathcal{K}$ is called the *suspension* of $\mathcal{K}$ and denoted $\Sigma\mathcal{K}$. Geometric realisations of $\operatorname{cone}\mathcal{K}$ and $\Sigma\mathcal{K}$ are the topological cone and suspension over $|\mathcal{K}|$ respectively.

4. Let P_1 and P_2 be simple polytopes. Then

$$\mathcal{K}_{P_1 \times P_2} = \mathcal{K}_{P_1} * \mathcal{K}_{P_2}.$$

(see Construction 1.1.12).

Construction 2.2.10. The fact that the product of two simplices is not a simplex makes triangulations of products of spaces more subtle. There is the following canonical way to triangulate the product of two simplicial complexes whose vertices are ordered. Let $\mathcal{K}_1$ and $\mathcal{K}_2$ be simplicial complexes on $[m_1]$ and $[m_2]$ respectively (this is one of the few constructions where the order of vertices is important: here it is an additional structure). We construct a simplicial complex $\mathcal{K}_1 \tilde{\times} \mathcal{K}_2$ on $[m_1] \times [m_2]$ as follows. By definition, simplices of $\mathcal{K}_1 \tilde{\times} \mathcal{K}_2$ are those subsets in products $I_1 \times I_2$ of $I_1 \in \mathcal{K}_1$ and $I_2 \in \mathcal{K}_2$ which establish non-decreasing relations between I_1 and I_2. More formally,

$$\begin{aligned}\mathcal{K}_1 \tilde{\times} \mathcal{K}_2 = \bigl\{ & I \subset I_1 \times I_2 \colon I_1 \in \mathcal{K}_1,\ I_2 \in \mathcal{K}_2, \\ & \text{and } i \leqslant i' \text{ implies } j \leqslant j' \text{ for any two pairs } (i,j),(i',j') \in I \bigr\}.\end{aligned}$$

We leave it as an exercise to check that $|\mathcal{K}_1 \tilde{\times} \mathcal{K}_2|$ defines a triangulation of $|\mathcal{K}_1| \times |\mathcal{K}_2|$. Note that $\mathcal{K}_1 \tilde{\times} \mathcal{K}_2 \neq \mathcal{K}_2 \tilde{\times} \mathcal{K}_1$ in general. Note also that if $\mathcal{K}_1 = \mathcal{K}_2 = \mathcal{K}$, then the diagonal is naturally a subcomplex in the triangulation of $|\mathcal{K}| \times |\mathcal{K}|$.

Construction 2.2.11 (connected sum of simplicial complexes). Let $\mathcal{K}_1$ and $\mathcal{K}_2$ be two pure d-dimensional simplicial complexes on sets $\mathcal{V}_1$ and $\mathcal{V}_2$ respectively, where $|\mathcal{V}_1| = m_1$, $|\mathcal{V}_2| = m_2$. Choose two maximal simplices $I_1 \in \mathcal{K}_1$, $I_2 \in \mathcal{K}_2$ and fix an identification of I_1 with I_2. Let $\mathcal{V}_1 \cup_I \mathcal{V}_2$ be the union of $\mathcal{V}_1$ and $\mathcal{V}_2$ in which I_1 is identified with I_2, and denote by I the subset created by the identification. We have $|\mathcal{V}_1 \cup_I \mathcal{V}_2| = m_1 + m_2 - d - 1$. Both $\mathcal{K}_1$ and $\mathcal{K}_2$ now can be viewed as simplicial complexes on the set $\mathcal{V}_1 \cup_I \mathcal{V}_2$. We define the *connected sum* of $\mathcal{K}_1$ and $\mathcal{K}_2$ at I_1 and I_2 as the simplicial complex

$$\mathcal{K}_1 \#_{I_1,I_2} \mathcal{K}_2 = (\mathcal{K}_1 \cup \mathcal{K}_2) \setminus \{I\}$$

on the set $\mathcal{V}_1 \cup_I \mathcal{V}_2$. When the choices are clear, or their effect on the result irrelevant, we use the abbreviation $K_1 \# K_2$. Geometrically, the connected sum of $|\mathcal{K}_1|$ and $|\mathcal{K}_2|$ is produced by attaching $|\mathcal{K}_1|$ to $|\mathcal{K}_2|$ along I_1 and I_2 and then removing the face I obtained by the identification.

Example 2.2.12. Connected sum of simple polytopes defined in Construction 1.1.14 is dual to the operation described above. Namely, if P and Q are two simple n-polytopes, then

$$\mathcal{K}_P \# \mathcal{K}_Q = \mathcal{K}_{P \# Q}.$$

Definition 2.2.13. Let $\mathcal{K}$ be a simplicial complex on a set $\mathcal{V}$. The *link* and the *star* of a simplex $I \in \mathcal{K}$ are the subcomplexes

$$\begin{aligned}\operatorname{lk}_{\mathcal{K}} I &= \bigl\{ J \in \mathcal{K} \colon I \cup J \in \mathcal{K},\ I \cap J = \varnothing \bigr\};\\ \operatorname{st}_{\mathcal{K}} I &= \bigl\{ J \in \mathcal{K} \colon I \cup J \in \mathcal{K} \bigr\}.\end{aligned}$$

We also define the subcomplex

$$\partial \operatorname{st}_{\mathcal{K}} I = \bigl\{ J \in \mathcal{K} \colon I \cup J \in \mathcal{K},\ I \not\subset J \bigr\}.$$

Then we have a sequence of inclusions

$$\operatorname{lk}_{\mathcal{K}} I \subset \partial \operatorname{st}_{\mathcal{K}} I \subset \operatorname{st}_{\mathcal{K}} I.$$

For any vertex $v \in \mathcal{K}$, the subcomplex $\operatorname{st}_{\mathcal{K}} v$ is the cone over $\operatorname{lk}_{\mathcal{K}} v = \partial \operatorname{st}_{\mathcal{K}} v$. Also, $|\operatorname{st}_{\mathcal{K}} v|$ is the union of all faces of $|\mathcal{K}|$ that contain v. We omit the subscript $\mathcal{K}$ in the notation of link and star whenever the ambient simplicial complex is clear.

The links of simplices determine the topological structure of the space $|\mathcal{K}|$ near any of its points. In particular, the following proposition describes the 'local cohomology' of $|\mathcal{K}|$.

PROPOSITION 2.2.14. *Let x be an interior point of a simplex $I \in \mathcal{K}$. Then*

$$H^i\bigl(|\mathcal{K}|, |\mathcal{K}| \setminus x\bigr) \cong \widetilde{H}^{i-|I|}(\operatorname{lk} I),$$

where $H^i(X, A)$ denotes the ith relative singular cohomology group of a pair $A \subset X$, and $\widetilde{H}^i(\mathcal{K})$ denotes the ith reduced simplicial cohomology group of $\mathcal{K}$.

PROOF. We have

$$\begin{aligned} H^i\bigl(|\mathcal{K}|, |\mathcal{K}| \setminus x\bigr) &\cong H^i\bigl(\operatorname{st} I, (\operatorname{st} I) \setminus x\bigr) \cong H^i\bigl(\operatorname{st} I, (\partial I) * (\operatorname{lk} I)\bigr) \cong \\ &\cong \widetilde{H}^{i-1}\bigl((\partial I) * (\operatorname{lk} I)\bigr) \cong \widetilde{H}^{i-|I|}(\operatorname{lk} I). \end{aligned}$$

Here the first isomorphism follows from the excision property, the second uses the fact that $(\partial I) * (\operatorname{lk} I)$ is a deformation retract of $(\operatorname{st} I) \setminus x$, the third follows from the homology sequence of pair, and the fourth is by the suspension isomorphism. □

Given a subcomplex $\mathcal{L} \subset \mathcal{K}$, define the *closed combinatorial neighbourhood* of $\mathcal{L}$ in $\mathcal{K}$ by

$$U_{\mathcal{K}}(\mathcal{L}) = \bigcup_{I \in \mathcal{L}} \operatorname{st}_{\mathcal{K}} I.$$

That is, $U_{\mathcal{K}}(\mathcal{L})$ consists of all simplices of $\mathcal{K}$, together with all their faces, having some simplex of $\mathcal{L}$ as a face. Define also the *open combinatorial neighbourhood* $\overset{\circ}{U}_{\mathcal{K}}(\mathcal{L})$ of $|\mathcal{L}|$ in $|\mathcal{K}|$ as the union of relative interiors of simplices of $|\mathcal{K}|$ having some simplex of $|\mathcal{L}|$ as a face.

DEFINITION 2.2.15. Given a subset $I \subset \mathcal{V}$, define the corresponding *full subcomplex* of $\mathcal{K}$ (or the *restriction* of $\mathcal{K}$ to I) as

$$\mathcal{K}_I = \bigl\{J \in \mathcal{K} \colon J \subset I\bigr\}. \tag{2.4}$$

Set $\operatorname{core} \mathcal{V} = \{v \in \mathcal{V} \colon \operatorname{st} v \neq \mathcal{K}\}$. The *core* of $\mathcal{K}$ is the subcomplex $\operatorname{core} \mathcal{K} = \mathcal{K}_{\operatorname{core} \mathcal{V}}$. Then we may write $\mathcal{K} = \operatorname{core}(\mathcal{K}) * \Delta^{s-1}$, where Δ^{s-1} is the simplex on the set $\mathcal{V} \setminus \operatorname{core} \mathcal{V}$.

EXAMPLE 2.2.16.

1. $\operatorname{lk}_{\mathcal{K}} \varnothing = \mathcal{K}$.
2. Let $K = \partial\Delta^3$ be the boundary of the tetrahedron on four vertices $1, 2, 3, 4$, and $I = \{1, 2\}$. Then $\operatorname{lk} I$ consists of two disjoint points 3 and 4.
3. Let $\mathcal{K}$ be the cone over $\mathcal{L}$ with vertex v. Then $\operatorname{lk} v = \mathcal{L}$, $\operatorname{st} v = \mathcal{K}$, and $\operatorname{core} \mathcal{K} = \operatorname{core} \mathcal{L}$.

Exercises.

2.2.17. Assume that $\mathcal{K}_1$ is realised geometrically in $\mathbb{R}^{n_1}$ and $\mathcal{K}_2$ in $\mathbb{R}^{n_2}$. Construct a realisation of the join $\mathcal{K}_1 * \mathcal{K}_2$ in $\mathbb{R}^{n_1+n_2+1}$.

2.2.18. Show that $|\mathcal{K}_1 \mathbin{\tilde{\times}} \mathcal{K}_2|$ is a triangulation of $|\mathcal{K}_1| \times |\mathcal{K}_2|$.

2.2.19. Let $\mathcal{K}$ be a pure simplicial complex. Then $\operatorname{lk} I$ is pure of dimension $\dim \mathcal{K} - |I|$ for any $I \in \mathcal{K}$.

2.3. Barycentric subdivision and flag complexes

Definition 2.3.1. The *barycentric subdivision* of an abstract simplicial complex $\mathcal{K}$ is the simplicial complex $\mathcal{K}'$ defined as follows. The vertex set of $\mathcal{K}'$ is the set $\{I \in \mathcal{K},\ I \neq \varnothing\}$ of nonempty simplices of $\mathcal{K}$. Simplices of $\mathcal{K}'$ are chains of embedded simplices of $\mathcal{K}$. That is, $\{I_1, \ldots, I_r\} \in \mathcal{K}'$ if and only if $I_1 \subset I_2 \subset \cdots \subset I_r$ in $\mathcal{K}$ (after possible reordering) and $I_1 \neq \varnothing$.

The *barycentre* of a geometric simplex in $\mathbb{R}^d$ with vertices $\boldsymbol{v}_1, \ldots, \boldsymbol{v}_{d+1}$ is the point $\frac{1}{d+1}(\boldsymbol{v}_1 + \cdots + \boldsymbol{v}_{d+1})$. A geometric realisation of $\mathcal{K}'$ may be obtained by mapping every vertex of $\mathcal{K}'$ to the barycentre of the corresponding simplex of $|\mathcal{K}|$; simplices of $|\mathcal{K}'|$ are therefore spanned by the sets of barycentres of chains of embedded simplices of $|\mathcal{K}|$.

Example 2.3.2. For any $(n-1)$-dimensional simplicial complex $\mathcal{K}$ on $[m]$, there is a nondegenerate simplicial map $\mathcal{K}' \to \Delta^{n-1}$ defined on the vertices by $I \mapsto |I|$ for $I \in \mathcal{K}$, where $|I|$ denotes the cardinality of I. Here I is viewed as a vertex of $\mathcal{K}'$ and $|I|$ as a vertex of Δ^{n-1}.

Example 2.3.3. Let $\mathcal{K}$ be a simplicial complex on a set $\mathcal{V}$, and assume we are given a choice function $c\colon \mathcal{K} \to \mathcal{V}$ assigning to each simplex $I \in \mathcal{K}$ one of its vertices. For instance, if $\mathcal{V} = [m]$ we can define $c(I)$ as the minimal element of I. For every such f there is a canonical simplicial map $\varphi_c\colon \mathcal{K}' \to \mathcal{K}$ constructed as follows. We define φ_c on the vertices of $\mathcal{K}'$ by $\varphi_c(I) = c(I)$ and then extend it to simplices of $\mathcal{K}'$ by the formula

$$\varphi_c(I_1 \subset I_2 \subset \cdots \subset I_r) = \{c(I_1), c(I_2), \ldots, c(I_r)\}.$$

The right hand side is a subset of I_r, and therefore it is a simplex of $\mathcal{K}$.

We shall need explicit formulae for the transformation of the f- and h-vectors under the barycentric subdivision. Introduce the matrix

$$B = (b_{ij}), \quad 0 \leqslant i, j \leqslant n-1; \quad b_{ij} = \sum_{k=0}^{i} (-1)^k \binom{i+1}{k} (i-k+1)^{j+1}.$$

It can be shown that $b_{ij} = 0$ for $i > j$ and $b_{ii} = (i+1)!$, so that B is a nonsingular upper triangular matrix.

Lemma 2.3.4. *Let $\mathcal{K}'$ be the barycentric subdivision of an $(n-1)$-dimensional simplicial complex $\mathcal{K}$. Then the f-vectors of $\mathcal{K}$ and $\mathcal{K}'$ are related by the identity*

$$f_i(\mathcal{K}') = \sum_{j=i}^{n-1} b_{ij} f_j(\mathcal{K}), \quad 0 \leqslant i \leqslant n-1,$$

that is, $\boldsymbol{f}^t(\mathcal{K}') = B\boldsymbol{f}^t(\mathcal{K})$ where $\boldsymbol{f}^t(\mathcal{K})$ is the column vector with entries $f_i(\mathcal{K})$.

Proof. Consider the barycentric subdivision of a j-simplex Δ^j, and let b'_{ij} be the number of its i-simplices which lie inside Δ^j, i.e. not contained in $\partial\Delta^j$. Then we have $f_i(\mathcal{K}') = \sum_{j=i}^{n-1} b'_{ij} f_j(\mathcal{K})$. It remains to show that $b_{ij} = b'_{ij}$. Indeed, it is easy to see that the numbers b'_{ij} satisfy the following recurrence relation:

$$b'_{ij} = (j+1)b'_{i-1,j-1} + \tbinom{j+1}{2} b'_{i-1,j-2} + \cdots + \tbinom{j+1}{j-i+1} b'_{i-1,i-1}.$$

It follows by induction that b'_{ij} is given by the same formula as b_{ij}. □

Now introduce the matrix

$$D=(d_{pq}),\quad 0\leqslant p,q\leqslant n;\quad d_{pq}=\sum_{k=0}^{p}(-1)^k\binom{n+1}{k}(p-k)^q(p-k+1)^{n-q},$$

where we set $0^0=1$.

Lemma 2.3.5. *The h-vectors of $\mathcal{K}$ and $\mathcal{K}'$ are related by the identity:*

$$h_p(\mathcal{K}')=\sum_{q=0}^{n}d_{pq}h_q(\mathcal{K}),\quad 0\leqslant p\leqslant n,$$

that is, $\boldsymbol{h}^t(\mathcal{K}')=D\boldsymbol{h}^t(\mathcal{K})$. Moreover, the matrix D is nonsingular.

Proof. The formula for $h_p(\mathcal{K}')$ is established by a routine check using Lemma 2.3.4, relations (2.3) and identities for the binomial coefficients. If we add the component $f_{-1}=1$ to the f-vector and change the matrix B appropriately, then we obtain $D=C^{-1}BC$, where C is the transition matrix from the h-vector to the f-vector (its explicit form can be obtained easily from relations (2.3)). This implies the nonsingularity of D. □

Definition 2.3.6. Let $\mathcal{P}$ be a poset (partially ordered set) with strict order relation $<$. Its *order complex* $\operatorname{ord}(\mathcal{P})$ is the collection of all totally ordered chains $x_1<x_2<\cdots<x_k$ (or *flags*), $x_i\in\mathcal{P}$. Clearly, $\operatorname{ord}(\mathcal{P})$ is a simplicial complex.

The following proposition is clear from the definition.

Proposition 2.3.7. *Let $\mathcal{K}$ be a simplicial complex, viewed as the poset of its simplices with respect to inclusion. Then $\operatorname{ord}(\mathcal{K}\setminus\varnothing)$ is the barycentric subdivision $\mathcal{K}'$. The order complex of $\mathcal{K}$ (with the empty simplex included) is $\operatorname{cone}(\mathcal{K}')$.*

This observation may be used to define the barycentric subdivision of other combinatorial objects. For example, let Q be a convex polytope, and $\mathcal{Q}$ the poset of its proper faces. Then $\operatorname{ord}(\mathcal{Q})$ is a simplicial complex; moreover, it is the boundary complex of a simplicial polytope Q' (an exercise), called the *barycentric subdivision* of Q. The vertices of Q' correspond to the barycentres of proper faces of Q.

Proposition 2.3.8. *Let P be a simple polytope and let $\mathcal{K}=\mathcal{K}_P$ be its nerve complex. Given a facet $F\subset P$, let v be the corresponding vertex of $\mathcal{K}$. Then $\operatorname{st}_{\mathcal{K}'}v$ is a triangulation of F.*

Proof. We identify ∂P with $\mathcal{K}'$ by mapping the barycentre of each proper face of P to the corresponding vertex of $\mathcal{K}'$. Under this identification, F is mapped to the union of simplices of $\mathcal{K}'$ corresponding to chains $G_1\subset\cdots\subset G_k=F$ of faces of P ending at F. This union is exactly the star of v in $\mathcal{K}'$. □

Definition 2.3.9. A simplicial complex $\mathcal{K}$ is called a *flag complex* if any set of vertices of $\mathcal{K}$ which are pairwise connected by edges spans a simplex.

A flag complex is therefore completely determined by its 1-skeleton, which is a simple graph. Given such a graph Γ, we may reconstruct the flag complex $\mathcal{K}_\Gamma$, whose simplices are the vertex sets of complete subgraphs (or *cliques*) of Γ.

Flag complexes may be characterised in terms of their missing faces. A *missing face* of $\mathcal{K}$ is a subset $I\subset[m]$ such that $I\notin\mathcal{K}$, but every proper subset of I is a simplex of $\mathcal{K}$. Then $\mathcal{K}$ is flag if and only if each of its missing faces has two vertices.

Therefore, P is a flag simple n-polytope if and only if any set of its facets whose pairwise intersections are nonempty intersects at a face.

Example 2.3.10.

1. The order complex of a poset is a flag complex.

2. A simple polytope P is a flag polytope (see Definition 1.6.1) if and only if its nerve complex $\mathcal{K}_P$ is a flag complex.

3. The boundary of a 5-gon is a flag complex, but it is not the order complex of a poset.

4. The boundary of a d-simplex is not flag for $d > 1$.

5. The join $\mathcal{K}_1 * \mathcal{K}_2$ of two flag complexes (see Construction 2.2.8) is flag (an exercise). Therefore, the product $P \times Q$ of two flag simple polytopes is flag.

6. The connected sum $\mathcal{K}_1 \# \mathcal{K}_2$ of two d-dimensional complexes (see Construction 2.2.11) is not flag if $d > 1$.

Since the h-vector of a sphere triangulation is symmetric, the *γ-vector* $\gamma(\mathcal{K}) = (\gamma_0, \gamma_1, \ldots, \gamma_{[n/2]})$ of an $(n-1)$-dimensional sphere triangulation $\mathcal{K}$ can be defined by the equation

$$\sum_{i=0}^{n} h_i s^i t^{n-i} = \sum_{i=0}^{[n/2]} \gamma_i (s+t)^{n-2i} (st)^i. \tag{2.5}$$

Conjecture 2.3.11 (Gal [**97**]). *Let $\mathcal{K}$ be a flag triangulation of an $(n-1)$-sphere. Then $\gamma_i(\mathcal{K}) \geqslant 0$ for $i = 0, \ldots, \left[\frac{n}{2}\right]$.*

Substituting $n = 2q$, $s = 1$ and $t = -1$ into (2.5) we obtain

$$\sum_{i=0}^{2q} (-1)^i h_i = (-1)^q \gamma_q\,.$$

The top inequality $\gamma_q \geqslant 0$ from the Gal conjecture therefore implies the following:

Conjecture 2.3.12 (Charney–Davis [**58**]). *The inequality*

$$(-1)^q \sum_{i=0}^{2q} (-1)^i h_i(\mathcal{K}) \geqslant 0$$

holds for any flag triangulation $\mathcal{K}$ of a $(2q-1)$-sphere.

The Charney–Davis conjecture is a discrete analogue of the well-known *Hopf conjecture* of differential geometry, which states that the Euler characteristic χ of a closed nonpositively curved Riemannian manifold M^{2q} satisfies the inequality $(-1)^q \chi \geqslant 0$. Due to a theorem of Gromov [**105**], a piecewise Euclidean cubical complex satisfies the discrete analogue of the nonpositive curvature condition, the so-called *CAT(0) inequality*, if and only if the links of all vertices are flag complexes. As it was observed in [**58**], the Hopf conjecture for piecewise Euclidean cubical manifolds translates into Conjecture 2.3.12.

The Hopf and Charney–Davis conjectures are valid for $q = 1, 2$. In [**97**] the Gal conjecture was verified for $n \leqslant 5$.

Exercises.

2.3.13. Show that the matrix B of Lemma 2.3.4 satisfies $b_{ij} = 0$ for $i > j$ and $b_{ii} = (i+1)!$.

2.3.14. Prove the formula for $h_p(\mathcal{K}')$ of Lemma 2.3.5.

2.3.15. Show that the order complex of the poset of proper faces of a polytope is the boundary complex of a simplicial polytope. (Hint: use stellar subdivisions, see Section 2.7 and the exercises there.)

2.3.16. The join of two flag complexes is flag.

2.3.17. Links of simplices in a flag complex are flag.

2.4. Alexander duality

For any simplicial complex, a dual complex may be defined on the same set. This duality has many important combinatorial and topological consequences.

Definition 2.4.1 (dual complex). Let $\mathcal{K}$ be a simplicial complex on $[m]$ and $\mathcal{K} \ne \Delta^{m-1}$. Define

$$\widehat{\mathcal{K}} = \bigl\{ I \subset [m]\colon \quad [m] \setminus I \notin \mathcal{K} \bigr\}.$$

Then $\widehat{\mathcal{K}}$ is also a simplicial complex on $[m]$, which we refer to as the *dual* of $\mathcal{K}$. Obviously, the dual of $\widehat{\mathcal{K}}$ is $\mathcal{K}$.

Construction 2.4.2. The barycentric subdivisions of both $\mathcal{K}$ and $\widehat{\mathcal{K}}$ can be realised as subcomplexes in the barycentric subdivision of the boundary of the standard simplex on the set $[m]$ in the following way.

A face of $(\partial\Delta^{m-1})'$ corresponds to a chain $I_1 \subset \cdots \subset I_r$ of included subsets in $[m]$ with $I_1 \ne \varnothing$ and $I_r \ne [m]$. We denote this face by $\Delta_{I_1 \subset \cdots \subset I_r}$. (For example, $\Delta_{\{i\}}$ is the ith vertex of Δ^{m-1} regarded as a vertex of $(\partial\Delta^{m-1})'$.) Then

$$G(\mathcal{K}) = \bigcup_{I_1 \subset \cdots \subset I_r,\ I_r \in \mathcal{K}} \Delta_{I_1 \subset \cdots \subset I_r} \tag{2.6}$$

is a geometric realisation of $\mathcal{K}'$. Denote $\widehat{i} = [m] \setminus \{i\}$ and, more generally, $\widehat{I} = [m] \setminus I$ for any subset $I \subset [m]$. Define the following subcomplex in $(\partial\Delta^{m-1})'$:

$$D(\mathcal{K}) = \bigcup_{I_1 \subset \cdots \subset I_r,\ \widehat{I}_r \notin \mathcal{K}} \Delta_{\widehat{I}_r \subset \cdots \subset \widehat{I}_1}.$$

Proposition 2.4.3. *For any simplicial complex $\mathcal{K} \ne \Delta^{m-1}$ on the set $[m]$, $D(\mathcal{K})$ is a geometric realisation of the barycentric subdivision of the dual complex:*

$$D(\mathcal{K}) = |\widehat{\mathcal{K}}'|. \tag{2.7}$$

Moreover, the open combinatorial neighbourhood of complex (2.6) *realising $\mathcal{K}'$ in $(\partial\Delta^{m-1})'$ coincides with the complement of the complex $D(\mathcal{K})$ realising $\widehat{\mathcal{K}}'$:*

$$\overset{\circ}{U}_{(\partial\Delta^{m-1})'}\bigl(|\mathcal{K}'|\bigr) = \bigl(\partial\Delta^{m-1}\bigr)' \setminus |\widehat{\mathcal{K}}'|.$$

In particular, $|\mathcal{K}'|$ is a deformation retract of the complement to $|\widehat{\mathcal{K}}'|$ in $(\partial\Delta^{m-1})'$.

Proof. We map a vertex $\{i\}$ of $\widehat{\mathcal{K}}$ to the vertex $\widehat{i} = \Delta_{\widehat{i}}$ of $(\partial\Delta^{m-1})'$, and map the barycentre of a face $I \in \widehat{\mathcal{K}}$ to the vertex $\Delta_{\widehat{I}}$. Then the whole complex $\widehat{\mathcal{K}}'$ is mapped to the subcomplex

$$\bigcup_{I_1 \subset \cdots \subset I_r,\ I_r \in \widehat{\mathcal{K}}} \Delta_{\widehat{I}_r \subset \cdots \subset \widehat{I}_1},$$

which is the same as $D(\mathcal{K})$. The second statement is left as an exercise. □

Example 2.4.4. Let $\mathcal{K}$ be the boundary of a 4-gon with vertices $1, 2, 3, 4$ (see Fig. 2.2). Then $\widehat{\mathcal{K}}$ consists of two disjoint segments. The picture shows both $\mathcal{K}'$ and $\widehat{\mathcal{K}}'$ as subcomplexes in $(\partial\Delta^3)'$.



Figure 2.2. Dual complex and Alexander duality.

Theorem 2.4.5 (Combinatorial Alexander duality). *For any simplicial complex $\mathcal{K} \neq \Delta^{m-1}$ on the set $[m]$ there is an isomorphism*

$$\widetilde{H}^j(\mathcal{K}) \cong \widetilde{H}_{m-3-j}(\widehat{\mathcal{K}}), \quad \text{for } -1 \leqslant j \leqslant m-2,$$

where $\widetilde{H}_k(\cdot)$ and $\widetilde{H}^k(\cdot)$ denote the kth reduced simplicial homology and cohomology group (with integer coefficients) respectively. Here we assume that $\widetilde{H}_{-1}(\varnothing) = \widetilde{H}^{-1}(\varnothing) = \mathbb{Z}$.

Proof. Since $(\partial\Delta^{m-1})'$ is homeomorphic to S^{m-2}, the Alexander duality theorem [**115**, Theorem 3.44] and Proposition 2.4.3 imply that

$$\begin{aligned}\widetilde{H}^j(\mathcal{K}) = \widetilde{H}^j\big(\overset{\circ}{U}_{(\partial\Delta^{m-1})'}(|\mathcal{K}'|)\big) = \widetilde{H}^j\big((\partial\Delta^{m-1})' \setminus |\widehat{\mathcal{K}}'|\big)\\ = \widetilde{H}^j\big(S^{m-2} \setminus |\widehat{\mathcal{K}}|\big) \cong \widetilde{H}_{m-3-j}(\widehat{\mathcal{K}}). \quad \square\end{aligned}$$

A more direct topological proof is outlined in Exercise 2.4.10. Theorem 2.4.5 can be also proved in a purely combinatorial way, see [**25**]. There is also a proof within 'combinatorial commutative algebra' (which is the subject of Chapter 3), see Exercise 3.2.15 or [**160**, Theorem 5.6].

The duality between $\mathcal{K}$ and $\widehat{\mathcal{K}}$ extends to a duality between full subcomplexes of $\mathcal{K}$ and links of simplices in $\widehat{\mathcal{K}}$:

Corollary 2.4.6. *Let $\mathcal{K} \neq \Delta^{m-1}$ be a simplicial complex on $[m]$ and $I \notin \mathcal{K}$, that is, $\widehat{I} \in \widehat{\mathcal{K}}$. Then there is an isomorphism*

$$\widetilde{H}^j(\mathcal{K}_I) \cong \widetilde{H}_{|I|-3-j}\bigl(\mathrm{lk}_{\widehat{\mathcal{K}}}\,\widehat{I}\bigr), \quad \text{for } -1 \leqslant j \leqslant |I|-2.$$

Proof. We apply Theorem 2.4.5 to the complex $\mathcal{K}_I$, viewed as a simplicial complex on the set I of $|I|$ elements. It follows from the definition that the dual complex is $\mathrm{lk}_{\widehat{\mathcal{K}}}\,\widehat{I}$, which also can be viewed as a simplicial complex on the set I. □

Example 2.4.7. Let $\mathcal{K}$ be the boundary of a pentagon. Then $\widehat{\mathcal{K}}$ is a Möbius band triangulated as shown on Fig. 2.3. Note that this $\widehat{\mathcal{K}}$ can be realised as a subcomplex in $\partial\Delta^4$, and therefore it can be realised in $\mathbb{R}^3$ as a subcomplex in the Schlegel diagram of Δ^4, see Construction 2.5.2.



Figure 2.3. The boundary of a pentagon and its dual complex.

Adding a ghost vertex to $\mathcal{K}$ results in suspending $\widehat{\mathcal{K}}$, up to homotopy. The precise statement is as follows (the proof is clear and is omitted).

Proposition 2.4.8. *Let $\mathcal{K}$ be a simplicial complex on $[m]$ and $\mathcal{K}^\circ$ be the complex on $[m+1]$ obtained by adding one ghost vertex $\circ = m+1$ to $\mathcal{K}$. Then the maximal simplices of $\widehat{\mathcal{K}^\circ}$ are $[m]$ and $I \cup \circ$ for $I \in \widehat{\mathcal{K}}$, that is,*

$$\widehat{\mathcal{K}^\circ} = \Delta^{m-1} \cup_{\widehat{\mathcal{K}}} \operatorname{cone}\widehat{\mathcal{K}}.$$

In particular, $\widehat{\mathcal{K}^\circ}$ is homotopy equivalent to the suspension $\Sigma\widehat{\mathcal{K}}$.

Exercises.

2.4.9. Show that

$$\overset{\circ}{U}_{(\partial\Delta^{m-1})'}\bigl(|\mathcal{K}'|\bigr) = \bigl(\partial\Delta^{m-1}\bigr)' \setminus |\widehat{\mathcal{K}}'|.$$

2.4.10. Complete the details in the following direct proof of combinatorial Alexander duality (Theorem 2.4.5).

There is a simplicial map

$$\varphi\colon (\partial\Delta^{m-1})' \to \mathcal{K}' * \widehat{\mathcal{K}}'$$

which is constructed as follows. We identify $|\mathcal{K}'|$ with $G(\mathcal{K})$ and $|\widehat{\mathcal{K}}'|$ with $D(\mathcal{K})$, see (2.6) and (2.7). The vertex sets of these two subcomplexes split the vertex set of $(\partial\Delta^{m-1})'$ into two nonintersecting subsets. Therefore, φ is uniquely determined on the vertices. Check that φ is indeed a simplicial map. It induces a map of simplicial cochains

$$\varphi^*\colon C^j(\mathcal{K}')\otimes C^{m-3-j}(\widehat{\mathcal{K}}')\to C^{m-2}\bigl((\partial\Delta^{m-1})'\bigr),$$

or, equivalently,

$$C^j(\mathcal{K}')\to C_{m-3-j}(\widehat{\mathcal{K}}')\otimes C^{m-2}\bigl((\partial\Delta^{m-1})'\bigr).$$

By evaluating on the fundamental cycle of $(\partial\Delta^{m-1})'$ in C_{m-2} and passing to simplicial (co)homology, we obtain the required isomorphism

$$H^j(\mathcal{K})\xrightarrow{\cong}H_{m-3-j}(\widehat{\mathcal{K}}).$$

2.5. Classes of triangulated spheres

Boundary complexes of simplicial polytopes form an important albeit restricted class of triangulated spheres. In this section we review several other classes of triangulated spheres and related complexes, and discuss their role in topology, geometry and combinatorics.

Definition 2.5.1. A *triangulated sphere* (also known as a *sphere triangulation* or *simplicial sphere*) of dimension d is a simplicial complex $\mathcal{K}$ homeomorphic to a d-sphere S^d. A *PL sphere* is a triangulated sphere $\mathcal{K}$ which is PL homeomorphic to the boundary of a simplex (equivalently, there exists a subdivision of $\mathcal{K}$ isomorphic to a subdivision of the boundary of a simplex).

Remark. A PL sphere is not the same as a 'PL manifold homeomorphic to a sphere', but rather a 'PL manifold which is PL homeomorphic to the *standard* sphere', where the standard PL structure on a sphere is defined by triangulating it as the boundary of a simplex. Nevertheless, the two notions coincide in dimensions other than 4, see the discussion in the next section.

In small dimensions, triangulated spheres can be effectively visualised using *Schlegel diagrams* and their generalisations, which we describe below.

Definition 2.5.2. By analogy with the notion of a geometric simplicial complex, we define a *polyhedral complex* as a collection $\mathcal{C}$ of convex polytopes in a space $\mathbb{R}^n$ such that every face of a polytope in $\mathcal{C}$ belongs to $\mathcal{C}$ and the intersection of any two polytopes in $\mathcal{C}$ is either empty or a face of each. Two polyhedral complexes $\mathcal{C}_1$ and $\mathcal{C}_2$ are said to be *combinatorially equivalent* if there is a one-to-one correspondence between their polytopes respecting the inclusion of faces.

The boundary ∂P of a convex polytope P is a polyhedral complex. Another important polyhedral complex associated with P can be constructed as follows. Choose a facet F of P and a point $p\notin P$ 'close enough' to F so that any segment connecting p to a point in $P\setminus F$ intersects the relative interior of F (see Fig. 2.4 for the case $P=I^3$). Now project the complex ∂P onto F from the point p. The projection images of faces of P different from F form a polyhedral complex $\mathcal{C}$ subdividing F. We refer to $\mathcal{C}$, and also to any polyhedral complex combinatorially equivalent to it, as a *Schlegel diagram* of the polytope P.

An *n-diagram* is a polyhedral complex $\mathcal{C}$ consisting of n-polytopes and their faces and satisfying the following conditions:

(a) the union of all polytopes in $\mathcal{C}$ is an n-polytope Q;



Figure 2.4. Schlegel diagram of I^3.

(b) every nonempty intersection of a polytope from $\mathcal{C}$ with the boundary of Q belongs to $\mathcal{C}$.

We refer to Q as the *base* of the n-diagram $\mathcal{C}$. An n-diagram is *simplicial* if it consists of simplices and its base is a simplex.

By definition, a Schlegel diagram of an n-polytope is an $(n-1)$-diagram.

Proposition 2.5.3. *Let $\mathcal{C}$ be a simplicial $(n-1)$-diagram with base Q. Then $\mathcal{C}\cup_{\partial Q} Q$ is an $(n-1)$-dimensional PL sphere.*

Proof. We need to construct a simplicial complex which is a subdivision of both $\mathcal{C}\cup_{\partial Q} Q$ and $\partial\Delta^n$. This can be done as follows. Replace one of the facets of $\partial\Delta^n$ by the diagram $\mathcal{C}$. The resulting simplicial complex $\mathcal{K}$ is a subdivision of $\partial\Delta^n$. On the other hand, $\mathcal{K}$ is isomorphic to the subdivision of $\mathcal{C}\cup_{\partial Q} Q$ obtained by replacing Q by a Schlegel diagram of Δ^n. □

Corollary 2.5.4. *The boundary of a simplicial n-polytope is an $(n-1)$-dimensional PL sphere.*

In dimensions $n \leqslant 3$ every $(n-1)$-diagram is a Schlegel diagram of an n-polytope. Indeed, for $n \leqslant 2$ this is obvious, and for $n = 3$ this is one of equivalent formulations of the well-known *Steinitz theorem* (see [**228**, Theorem 5.8]). The first example of a 3-diagram which is not a Schlegel diagram of a 4-polytope was found by Grünbaum ([**107**, §11.5], see also [**108**]) as a correction of Brückner's result of 1909 on the classification of simplicial 4-polytopes with 8 vertices. Another example was found by Barnette [**14**]. We describe Barnette's example below. For the original example of Grünbaum, see Exercise 2.5.15.

Construction 2.5.5 (Barnette's 3-diagram). Here a certain simplicial 3-diagram $\mathcal{C}$ will be constructed. Consider the octahedron Q obtained by twisting the top face (abc) of a triangular prism (Fig. 2.5 (a)) slightly so that the vertices a, b, c, d, e and f are in general position. Assume that the edges (bd), (ce) and



(a) (b)

FIGURE 2.5. Barnette's 3-diagram.

(af) lie inside Q. The tetrahedra $(abde)$, $(bcef)$ and $(acdf)$ will be included in the complex $\mathcal{C}$. Each of these tetrahedra has two faces which lie inside Q. These 6 triangles together with (abc) and (def) form a triangulated 2-sphere inside Q, which we denote by $\mathcal{S}$. Now place a point p inside $\mathcal{S}$ so that the line segments from p to the vertices of $\mathcal{S}$ lie inside $\mathcal{S}$. We add to $\mathcal{C}$ the eight tetrahedra obtained by taking cones with vertex p over the faces of $\mathcal{S}$ (namely, the tetrahedra $(pabc)$, $(pdef)$, $(pabd)$, $(pbed)$, $(pbce)$, $(pcef)$, $(pacf)$ and $(padf)$). Note that $\mathcal{S} = \operatorname{lk} p$. Applying a projective transformation if necessary, we may assume that there is a point p' outside the octahedron Q with the property that the segments joining p' with the vertices of Q lie outside Q (see Fig. 2.5 (b)). Finally, we add to $\mathcal{C}$ the tetrahedra obtained by taking cones with vertex p' over the faces of Q other than the face (def) (there are 7 such tetrahedra: $(p'abc)$, $(p'abe)$, $(p'ade)$, $(p'acd)$, $(p'cdf)$, $(p'bcf)$ and $(p'bef)$). The union of all tetrahedra in $\mathcal{C}$ is the tetrahedron $(p'def)$; hence, $\mathcal{C}$ is a simplicial 3-diagram. It has 8 vertices and 18 tetrahedra.

PROPOSITION 2.5.6. *The 3-diagram $\mathcal{C}$ from the previous construction is not a Schlegel diagram.*

PROOF. Suppose there is a polytope P whose Schlegel diagram is $\mathcal{C}$. Since P is simplicial, we may assume that its vertices are in general position. We label the vertices of P with the same letters as the corresponding vertices in $\mathcal{C}$. Consider the complex $\mathcal{S} = \operatorname{lk} p$. Let P' be the convex hull of all vertices of P other than p. Then P' is still a simplicial polytope, and $\mathcal{S}$ is a subcomplex in $\partial P'$. In the complex $\partial P'$ the sphere $\mathcal{S}$ is filled with tetrahedra whose vertices belong to $\mathcal{S}$. Then at least one edge of one of these tetrahedra lies inside $\mathcal{S}$. However, any two vertices of $\mathcal{S}$ which are not joined by an edge on $\mathcal{S}$ are joined by an edge of $\mathcal{C}$ lying outside $\mathcal{S}$. Since the polytope P' cannot contain a double edge we have reached a contradiction. ☐

Now we can introduce two more classes of triangulated spheres.

Definition 2.5.7. A *polytopal sphere* is a triangulated sphere isomorphic to the boundary complex of a simplicial polytope.

A *star-shaped sphere* is a triangulated sphere isomorphic to the underlying complex of a complete simplicial fan. Equivalently, a triangulated sphere $\mathcal{K}$ of dimension $n-1$ is star-shaped if there is a geometric realisation of $\mathcal{K}$ in $\mathbb{R}^n$ and a point $p \in \mathbb{R}^n$ with the property that each ray emanating from p meets $|\mathcal{K}|$ in exactly one point. The set of points such $p \in \mathbb{R}^n$ is called the *kernel* of $|\mathcal{K}|$.

Example 2.5.8. The triangulated 3-sphere coming from Barnette's 3-diagram is known as the *Barnette sphere*. It is star-shaped. Indeed, in the construction of Barnette's 3-diagram we have the octahedron $Q \subset \mathbb{R}^3$ and a vertex p' outside Q such that $\operatorname{lk} p' = \partial Q$. If we choose the vertex p' in $\mathbb{R}^4 \setminus \mathbb{R}^3$, then the Barnette sphere can be realised in $\mathbb{R}^4$ as the boundary complex of the pyramid with vertex p' and base Q (subdivided as described in Construction 2.5.5). This realisation is obviously star-shaped.

We therefore have the following hierarchy of triangulations:

$$\begin{array}{ccccccc} \text{polytopal} & & \text{star-shaped} & & PL & & \text{triangulated} \\ \text{spheres} & \subset & \text{spheres} & \subset & \text{spheres} & \subset & \text{spheres} \end{array} \tag{2.8}$$

Here the first inclusion follows from the construction of the normal fan (Construction 2.1.3), and the second is left as an exercise.

In dimension 2 any triangulated sphere is polytopal (this is another corollary of the Steinitz theorem). Also, by the result of Mani [**147**], any triangulated d-dimensional sphere with up to $d+4$ vertices is polytopal. However, in general all inclusions above are strict.

The first inclusion in (2.8) is strict already in dimension 3, as is seen from Example 2.5.8. There are 39 combinatorially different triangulations of a 3-sphere with 8 vertices, among which exactly two are nonpolytopal (namely, the Barnette and Brückner spheres); this classification was completed by Barnette [**17**].

The second inclusion in (2.8) is also strict in dimension 3. The first example of a non-star-shaped sphere triangulation was found by Ewald and Schulz [**87**]. We sketch this example below, following [**86**, Theorem 5.5].

Example 2.5.9 (Non-star-shaped sphere triangulation). We use the fact, observed by Barnette, that not every tetrahedron in Barnette's 3-diagram can be chosen as the base of a 3-diagram of the Barnette sphere (see Exercise 2.5.16). For instance, the tetrahedron $(abcd)$ cannot be chosen as the base.

Now let $\mathcal{K}$ be a connected sum of two copies of the Barnette sphere along the tetrahedra $(abcd)$ (the identification of vertices in the two tetrahedra is irrelevant). Assume that $\mathcal{K}$ has a star-shaped realisation in $\mathbb{R}^4$. The hyperplane H through the points a, b, c, d splits $|\mathcal{K}|$ into two parts $|\mathcal{K}_1|$ and $|\mathcal{K}_2|$. Since the kernel of $|\mathcal{K}|$ is an open set, it contains a point p not lying in H. Then by projecting either $|\mathcal{K}_1|$ or $|\mathcal{K}_2|$ onto H from p, we obtain a 3-diagram of the Barnette sphere with base $(abcd)$. This is a contradiction.

The fact that $\mathcal{K}$ is a PL sphere is an exercise.

The third inclusion in (2.8) is the subtlest one. It is known that in dimension 3 any triangulated sphere is PL. In dimension 4 the corresponding question is open, but starting from dimension 5 there exist *non-PL* sphere triangulations. See the discussion in the next section and Example 2.6.3.

Many important open problems of combinatorial geometry arise from analysing the relationships between different classes of sphere triangulations. We end this section by discussing some of these problems.

In connection with the condition of realisability of a triangulated $(n-1)$-sphere in $\mathbb{R}^n$ in the definition of a star-shaped sphere, we note that the existence of such a realisation is open in general:

PROBLEM 2.5.10. Does every PL $(n-1)$-sphere admit a geometric realisation in an n-dimensional space?

The g-theorem (Theorem 1.4.14) gives a complete characterisation of integral vectors arising as the f-vectors of polytopal spheres. It is therefore natural to ask whether the g-theorem extends to all sphere triangulations. This question was posed by McMullen [**155**] as an extension of his conjecture for simplicial polytopes. Since 1980, when McMullen's conjecture for simplicial polytopes was proved by Billera, Lee, and Stanley, its generalisation to spheres has been regarded as the main open problem in the theory of f-vectors:

PROBLEM 2.5.11 (g-conjecture for triangulated spheres). Does Theorem 1.4.14 hold for triangulated spheres?

The g-conjecture is open even for star-shaped spheres. Note that only the necessity of the conditions in the g-theorem (that is, the fact that every g-vector is an M-vector) has to be verified for triangulated spheres. If correct, the g-conjecture would imply a complete characterisation of f-vectors of triangulated spheres.

The Dehn–Sommerville relations (condition (a) in Theorem 1.4.14) hold for arbitrary sphere triangulations (see Corollary 3.4.7 below). The f-vectors of triangulated spheres also satisfy the UBT and LBT inequalities given in Theorems 1.4.4 and 1.4.9 respectively. The proof of the Lower Bound Theorem for simplicial polytopes given by Barnette in [**16**] extends to all triangulated spheres (see also [**130**]). In particular, this implies the second GLBC inequality $h_1 \leqslant h_2$, see (1.19). The Upper Bound Theorem for triangulated spheres was proved by Stanley [**203**] (we shall give his argument in Section 3.3). This implies that the g-conjecture is true for triangulated spheres of dimension $\leqslant 4$. The third GLBC inequality $h_2 \leqslant h_3$ (for spheres of dimension $\geqslant 5$) is open.

Many attempts to prove the g-conjecture were made after 1980. Though unsuccessful, these attempts resulted in some very interesting reformulations of the g-conjecture. The results of Pachner [**175**] reduce the g-conjecture (for PL spheres) to some properties of *bistellar moves*; see the discussion after Theorem 2.7.3 below.

The lack of progress in proving the g-conjecture motivated Björner and Lutz to launch a computer-aided search for counterexamples [**24**]. Though their bistellar flip algorithm and BISTELLAR software produced many remarkable results on triangulations of manifolds, no counterexamples to the g-conjecture were found. More information on the g-conjecture and related questions may be found in [**208**] and [**228**, Lecture 8].

We may extend the hierarchy (2.8) by considering polyhedral complexes homeomorphic to spheres (the so-called *polyhedral spheres*) instead of triangulated spheres. There are obvious analogues of polytopal and PL spheres in this generality. However, unlike the case of triangulations, the two definitions of a star-shaped sphere (namely, the one using fans and the one using the kernel points) no longer produce the same classes of objects, see [**86**, § III.5] for the corresponding examples. One of the most notorious and long standing problems is to find a proper higher dimensional analogue to the Steinitz theorem. This theorem characterises graphs of 3-dimensional polytopes, and one of its equivalent formulations is that every polyhedral 2-sphere is polytopal. In higher dimensions, identification of the class of polytopal spheres inside all polyhedral spheres is known as the *Steinitz problem*:

PROBLEM 2.5.12 (Steinitz problem). Find necessary and sufficient conditions for a polyhedral decomposition of a sphere to be combinatorially equivalent to the boundary complex of a convex polytope.

This is far from being solved even in the case of triangulated spheres. For more information on the relationships between different classes of polyhedral spheres and complexes see the above cited book of Ewald [**86**] and the survey article by Klee and Kleinschmidt [**136**].

Exercises.

2.5.13. Show that $\mathcal{K}$ is the underlying complex of a complete simplicial fan if and only if there is a geometric realisation of $\mathcal{K}$ in $\mathbb{R}^n$ and a point $p \in \mathbb{R}^n$ with the property that each ray emanating from p meets $|\mathcal{K}|$ in exactly one point.

2.5.14. Prove that every star-shaped sphere is a PL sphere.

2.5.15 (Brückner sphere). The *Brückner sphere* is obtained by replacing two tetrahedra $(pabc)$ and $(p'abc)$ in the Barnette sphere by three tetrahedra $(pp'ab)$, $(pp'ac)$ and $(pp'bc)$ (this is an example of a bistellar 1-move considered in Section 2.7). Show that the Brückner sphere is star-shaped but not polytopal. Note that the 1-skeleton of the Brückner sphere is a complete graph (that is, the Brückner sphere is a *neighbourly* triangulation, see Definition 1.1.16).

2.5.16. Show that the tetrahedron $(abcd)$ in Barnette's 3-diagram (Construction 2.5.5) cannot be chosen as the base of a 3-diagram of the Barnette sphere. Which tetrahedra can be chosen as the base?

2.5.17. The connected sum of two PL spheres is a PL sphere.

2.6. Triangulated manifolds

Piecewise linear topology experienced an intensive development during the second half of the 20th century, thanks to the efforts of many topologists. Surgery theory for simply connected manifolds of dimension $\geqslant 5$ originated from the early work of Milnor, Kervaire, Browder, Novikov and Wall, culminated in the proof of the topological invariance of rational Pontrjagin classes given by Novikov in 1965, and was further developed in the work of Lashof, Rothenberg, Sullivan, Kirby, Siebenmann, and others. It led to a better understanding of the place of PL manifolds between the topological and smooth categories. Without attempting to overview the current state of the subject, which is generally beyond the scope of this book, we include several important results on the triangulation of topological manifolds, with a particular emphasis on various nonexamples. We also provide references for further reading.

All manifolds here are compact, connected and closed, unless otherwise stated.

Definition 2.6.1. A *triangulated manifold* (or *simplicial manifold*) is a simplicial complex $\mathcal{K}$ whose geometric realisation $|\mathcal{K}|$ is a topological manifold.

A *PL manifold* (or *combinatorial manifold*) is a simplicial complex $\mathcal{K}$ of dimension d such that $\operatorname{lk} I$ is a PL sphere of dimension $d-|I|$ for every nonempty simplex $I \in \mathcal{K}$.

Every PL manifold $|\mathcal{K}|$ of dimension d is a triangulated manifold: it has an atlas whose change of coordinates functions are piecewise linear. Indeed, for each vertex $v \in |\mathcal{K}|$ the $(d-1)$-dimensional PL sphere $\operatorname{lk} v$ bounds an open neighbourhood U_v which is homeomorphic to an open d-ball. Since any point of $|\mathcal{K}|$ is contained in U_v for some v, this defines an atlas for $|\mathcal{K}|$.

Remark. The term 'PL manifold' is often used for a manifold with a PL atlas, while its particular triangulation with the property above is referred to as a combinatorial manifold. We shall not distinguish between these two notions.

Does every triangulation of a topological manifold yield a simplicial complex which is a PL manifold? The answer is 'no', and the question itself ascends to a famous conjecture of the dawn of topology, known as the *Hauptvermutung*, which is German for 'main conjecture'. Below we briefly review the current status of this conjecture, referring to the survey article [**194**] by Ranicki for a much more detailed historical account and more references.

In the early days of topology all of the known topological invariants were defined in combinatorial terms, and it was very important to find out whether the topology of a triangulated space fully determines the combinatorial equivalence class of the triangulation (in the sense of Definition 2.2.6). In the modern terminology, the Hauptvermutung states that any two homeomorphic simplicial complexes are combinatorially equivalent (PL homeomorphic). This is valid in dimensions $\leqslant 3$; the result is due to Rado (1926) for 2-manifolds, Papakyriakopoulos (1943) for 2-complexes, Moise (1953) for 3-manifolds, and E. Brown (1963) for 3-complexes [**33**]; see [**164**] for a detailed exposition. The first examples of complexes disproving the Hauptvermutung in dimensions $\geqslant 6$ were found by Milnor in the early 1960s. However, the *manifold Hauptvermutung*, namely the question of whether two homeomorphic triangulated manifolds are combinatorially equivalent, remained open until the end of the 1960s. The first counterexamples were found by Siebenmann in 1969, and relied heavily on the topological surgery theory. The 'double suspension theorem', which we state as Theorem 2.6.2 below, appeared around 1975 and provided much more explicit counterexamples to the manifold Hauptvermutung.

A *d-dimensional homology sphere* (or simply *homology d-sphere*) is a topological d-manifold whose integral homology groups are isomorphic to those of a d-sphere S^d.

Theorem 2.6.2 (Edwards, Cannon). *The double suspension of any homology d-sphere is homeomorphic to S^{d+2}.*

This theorem was proved for most double suspensions and all triple suspensions by Edwards [**81**]; the general case was done by Cannon [**55**]. One of its most important consequences is the existence of non-PL triangulations of 5-spheres, which also disproves the manifold Hauptvermutung in dimensions $\geqslant 5$.

Example 2.6.3 (non-PL triangulated 5-sphere). Let M be a triangulated homology 3-sphere which is not homeomorphic to S^3. An example of such M is provided by the *Poincaré sphere*. It is the homogeneous space $SO(3)/A_5$, where the alternating group A_5 is represented in $\mathbb{R}^3$ as the group of self-transformations of a dodecahedron. A particular symmetric triangulation of the Poincaré sphere is given in [**24**]. By Theorem 2.6.2, the double suspension $\Sigma^2 M$ is homeomorphic to S^5 (and, more generally, $\Sigma^k M$ is homeomorphic to S^{k+3} for $k \geqslant 2$). However, $\Sigma^2 M$ cannot be a PL sphere, since M appears as the link of a 1-simplex in $\Sigma^2 M$.

Also, according to a result of Björner and Lutz [**24**], for any $d \geqslant 5$ there is a non-PL triangulation of S^d with $d+13$ vertices.

Theorem 2.6.2 led to progress in the following 'manifold recognition problem': given a simplicial complex, how one can decide whether its geometric realisation is a topological manifold? In higher dimensions there is the following result, which can be viewed as a generalisation of the double suspension theorem.

Theorem 2.6.4 (Edwards [**82**]). *For $d \geqslant 5$ the realisation of a simplicial complex $\mathcal{K}$ is a topological manifold of dimension d if and only if $\operatorname{lk} I$ has the homology of a $(d-|I|)$-sphere for each nonempty simplex $I \in \mathcal{K}$, and $\operatorname{lk} v$ is simply connected for each vertex v of $\mathcal{K}$.*

Remark. From the algorithmic point of view, the homology of links is easily computable, but their simply connectedness seems to be undecidable. There is a related result of Novikov [**225**, Appendix] that a triangulated 5-sphere cannot be algorithmically recognised. On the other hand, the algorithmic recognition problem for a triangulated 3-sphere has a positive solution, with the first algorithm provided by Rubinstein [**198**]. See detailed exposition in Matveev's book [**152**], which also contains a proof that all 3-dimensional *Haken manifolds* can be recognised and fully classified algorithmically.

With the discovery of exotic smooth structures on 7-spheres by Milnor and the disproval of the Hauptvermutung it had become important to understand better the relationship between PL and smooth structures on topological manifolds. Since a PL structure implies the existence of a particular sort of triangulation, the related question of whether a topological manifold admits *any* triangulation (not necessarily PL) had also become important.

Triangulations of 2-manifolds have been known from the early days of topology. A proof that any 3-manifold can be triangulated was obtained independently by Moise and Bing in the end of 1950s (the proof can be found in [**164**]). Since the link of a vertex in a triangulated 3-sphere is a 2-sphere, and a 2-sphere is always PL, all topological 3-manifolds are PL.

A smooth manifold of any dimension has a PL triangulation by a theorem of Whitney (a proof can be found in [**165**]). Moreover, in dimensions $\leqslant 3$ every topological manifold has a unique smooth structure, see [**164**] for a proof. All these considerations show that in dimensions up to 3 the categories of topological, PL and smooth manifolds are equivalent.

The situation in dimension 4 is quite different. There exist topological 4-manifolds that do not admit a PL triangulation. An example is provided by Freedman's fake $\mathbb{C}P^2$ [**94**, §8.3, §10.1], a topological manifold which is homeomorphic, but not diffeomorphic to the complex projective plane $\mathbb{C}P^2$. This example also shows that the Hauptvermutung is false in dimension 4. Even worse, some topological 4-manifolds do not admit any triangulation; an example is the topological 4-manifold with the intersection form E_8, see [**3**].

In dimension 4 the categories of PL and smooth manifolds agree, that is, there is exactly one smooth structure on every PL manifold. However, the classification of PL (or equivalently, smooth) structures is wide open even for the simplest topological manifolds. The most notable problem here is the following.

PROBLEM 2.6.5. Is a PL (or smooth) structure on a 4-sphere unique?

In dimensions $\geqslant 5$ the PL structure on a topological sphere is unique (that is, a PL manifold which is homeomorphic to a sphere is a PL sphere), but the triangulation problem is open:

PROBLEM 2.6.6 (Triangulation Conjecture). Is it true that any topological manifold of dimension $\geqslant 5$ can be triangulated?

For the discussion of the current status of the classification of PL structures on topological manifolds we refer to Ranicki's survey [**194**], the original essay [**134**] by Kirby and Siebenmann, and a more recent survey by Rudyak [**199**].

2.7. Stellar subdivisions and bistellar moves

By a theorem of Alexander, a common subdivision of two PL homeomorphic PL manifolds can be obtained by iterating operations from a very simple and explicit list, known as *stellar subdivisions*. An even more concrete iterative description of PL homeomorphisms was obtained by Pachner [**174**], who introduced the notion of *bistellar moves* (in other terminology, *bistellar flips* or *bistellar operations*), generalising the 2- and 3-dimensional *flips* from low-dimensional topology. These operations allow us to decompose a PL homeomorphism into a sequence of simple 'moves' and thus provide a very convenient way to compute and handle topological invariants of PL manifolds. Starting from a given PL triangulation, bistellar operations may be used to construct new triangulations with some good properties, such as ones that are symmetric or have a small number of vertices. On the other hand, bistellar moves can be used to produce some nasty triangulations if we start from a non-PL triangulation. Both approaches were used in the work of Björner–Lutz [**24**] and Lutz [**143**] to find many interesting triangulations of low-dimensional manifolds. In our exposition of bistellar moves we follow the terminology of [**143**].

Bistellar moves also provide a combinatorial interpretation for algebraic *flop* operations for projective *toric varieties* and for certain surgery operations on moment-angle complexes and torus manifolds. Finally, bistellar moves may be used to define a metric on the space of PL triangulations of a given PL manifold, see [**167**].

DEFINITION 2.7.1. Let $I \in \mathcal{K}$ be a nonempty simplex of a simplicial complex $\mathcal{K}$. The *stellar subdivision* of $\mathcal{K}$ at I is obtained by replacing the star of I by the cone over its boundary:

$$\operatorname{ss}_I \mathcal{K} = (\mathcal{K} \setminus \operatorname{st}_{\mathcal{K}} I) \cup (\operatorname{cone} \partial \operatorname{st}_{\mathcal{K}} I).$$

If $\dim I = 0$ then $\operatorname{ss}_I \mathcal{K} = \mathcal{K}$. Otherwise the complex $\operatorname{ss}_I \mathcal{K}$ acquires an additional vertex (the vertex of the cone) whose link is $\partial \operatorname{st}_{\mathcal{K}} I$. Two possible stellar subdivisions of a 2-dimensional complex are shown in Fig. 2.6.

Now let $\mathcal{K}$ be a triangulated manifold of dimension d. Assume that $I \in \mathcal{K}$ is a $(d-j)$-face such that the simplicial complex $\operatorname{lk}_{\mathcal{K}} I$ is the boundary of a j-simplex J which is not a face of $\mathcal{K}$. Then the operation bm_I on $\mathcal{K}$ defined by

$$\operatorname{bm}_I \mathcal{K} = \bigl(\mathcal{K} \setminus (I * \partial J)\bigr) \cup (\partial I * J)$$



FIGURE 2.6. Stellar subdivisions at a 2-simplex and at an edge.

is called a *bistellar j-move*. Since $I * \partial J = \mathrm{st}_{\mathcal{K}}\, I$ and $\partial I * J = \mathrm{st}_{\widetilde{\mathcal{K}}}\, J$, where $\widetilde{\mathcal{K}} = \mathrm{bm}_I\,\mathcal{K}$, the bistellar j-move is the composition of a stellar subdivision at I and the inverse stellar subdivision at J, which explains the term. In particular, the stellar subdivision $\mathrm{ss}_I\,\mathcal{K}$ is a common subdivision of $\mathcal{K}$ and $\widetilde{\mathcal{K}}$, so that $\mathcal{K}$ and $\widetilde{\mathcal{K}}$ are combinatorially equivalent. Note that a 0-move is the stellar subdivision at a maximal simplex (we assume that the boundary of a 0-simplex is empty).

Bistellar j-moves with $i \geqslant \left[\frac{d}{2}\right]$ are also called *reverse $(d-j)$-moves*. A 0-move adds a new vertex to the triangulation, a d-move (reverse 0-move) deletes a vertex, and all other bistellar moves do not change the number of vertices. The bistellar moves in dimension 2 and 3 are shown in Figures 2.7 and 2.8. The bistellar 1-move in dimension 3 replaces two tetrahedra with a common face by 3 tetrahedra with a common edge.

Two simplicial complexes are said to be *bistellarly equivalent* if one can be transformed to another by a finite sequence of bistellar moves.

We have seen that bistellar equivalence implies PL homeomorphism. The following result shows that for PL manifolds the converse is also true.

THEOREM 2.7.2 (Pachner [**174**, Theorem 1], [**175**, (5.5)]). *Two PL manifolds are bistellarly equivalent if and only if they are PL homeomorphic.*

The behaviour of the face numbers of a triangulation under bistellar moves is easily controlled. It can be most effectively described in terms of the g-vector, $g_i(\mathcal{K}) = h_i(\mathcal{K}) - h_{i-1}(\mathcal{K}),\ \ 0 < i \leqslant \left[\frac{d}{2}\right]$:

THEOREM 2.7.3 (Pachner [**174**]). *If a triangulated d-manifold $\widetilde{\mathcal{K}}$ is obtained from $\mathcal{K}$ by a bistellar k-move, $0 \leqslant k \leqslant \left[\frac{d-1}{2}\right]$, then*

$$
\begin{aligned}
g_{k+1}(\widetilde{\mathcal{K}}) &= g_{k+1}(\mathcal{K}) + 1;\\
g_i(\widetilde{\mathcal{K}}) &= g_i(\mathcal{K}) \quad \textit{for all } i \neq k+1.
\end{aligned}
$$

Furthermore, if d is even and $\widetilde{\mathcal{K}}$ is obtained from $\mathcal{K}$ by a bistellar $\left[\frac{d}{2}\right]$-move, then

$$
g_i(\widetilde{\mathcal{K}}) = g_i(\mathcal{K}) \quad \textit{for all } i.
$$



FIGURE 2.7. Bistellar moves for $q = 2$.



FIGURE 2.8. Bistellar moves for $q = 3$.

This theorem allows us to interpret the inequalities from the g-conjecture for PL spheres (see Theorem 1.4.14) in terms of the numbers of bistellar k-moves needed to transform a given PL sphere to the boundary of a simplex. For instance, the inequality $h_1 \leqslant h_2, \;\; n \geqslant 4$, is equivalent to the statement that the number of 1-moves in the sequence of bistellar moves taking a given $(n-1)$-dimensional

PL sphere to the boundary of an n-simplex is less than or equal to the number of reverse 1-moves. (Note that the g-vector of $\partial\Delta^n$ is $(1,0,\ldots,0)$.)

Remark. There is also a generalisation of Theorem 2.7.2 to PL manifolds with boundary, see [**175**, (6.3)].

In the case of polytopal sphere triangulations a stellar subdivision is related to another familiar operation:

Proposition 2.7.4. *Assume given a simple polytope P and a proper face $G \subset P$. Let P^* be the dual simplicial polytope, $\mathcal{K}_P = \partial P^*$ its nerve complex, and $J \subset P^*$ the face dual to G. Then the stellar subdivision $\mathrm{ss}_J\,\mathcal{K}_P$ is the nerve complex of the polytope $\widetilde{P}$ obtained by the face truncation at G.*

Proof. This follows directly by comparing the face poset of $\widetilde{P}$, described in Construction 1.1.13, with that of $\mathrm{ss}_J\,\mathcal{K}_P$. □

Exercises.

2.7.5. The barycentric subdivision of $\mathcal{K}$ can be obtained as a sequence of stellar subdivisions at all faces $I \in \mathcal{K}$, starting from the maximal ones.

2.7.6. Deduce formulae for the transformation of the f-, h- and g-vector of $\mathcal{K}$ under a stellar subdivision. Deduce similar formulae for a bistellar move (the case of the g-vector is Theorem 2.7.3).

2.8. Simplicial posets and simplicial cell complexes

Simplicial posets describe the combinatorial structures underlying 'generalised simplicial complexes' whose faces are still simplices, but two faces are allowed to intersect in any subcomplex of their boundary, rather than just in a single face. These are also known as 'ideal triangulations' in low-dimensional topology, or as 'simplicial cell complexes'.

Definition 2.8.1. A poset (partially ordered set) $\mathcal{S}$ with order relation $\leqslant$ is called *simplicial* if it has an initial element $\hat{0}$ and for each $\sigma \in \mathcal{S}$ the lower segment

$$[\hat{0},\sigma] = \{\tau \in \mathcal{S}\colon \hat{0} \leqslant \tau \leqslant \sigma\}$$

is the face poset of a simplex. (The latter poset is also known as a *Boolean lattice*, and simplicial posets are sometimes called *Boolean posets*.) We assume all our posets to be finite. The *rank function* $|\cdot|$ on $\mathcal{S}$ is defined by setting $|\sigma| = k$ if $[\hat{0},\sigma]$ is the face poset of a $(k-1)$-dimensional simplex. The rank of $\mathcal{S}$ is the maximum of ranks of its elements, and the *dimension* of $\mathcal{S}$ is its rank minus one. A *vertex* of $\mathcal{S}$ is an element of rank one. We assume that $\mathcal{S}$ has m vertices, denote the vertex set by $V(\mathcal{S})$, and usually identify it with $[m] = \{1,\ldots,m\}$. Similarly, we denote by $V(\sigma)$ the vertex set of σ, that is the set of i with $i \leqslant \sigma$.

The face poset of a simplicial complex is a simplicial poset, but there are many simplicial posets that do not arise in this way (see Example 2.8.2 below). We identify a simplicial complex with its face poset, thereby regarding simplicial complexes as particular cases of simplicial posets.

To each $\sigma \in \mathcal{S}$ we assign a geometric simplex Δ^σ whose face poset is $[\hat{0},\sigma]$, and glue these geometric simplices together according to the order relation in $\mathcal{S}$. As a result we get a regular cell complex in which the closure of each cell is identified with a simplex preserving the face structure, and all attaching and characteristic maps are inclusions (see [**115**, Appendix] for the terminology of cell complexes). We call it the *simplicial cell complex* associated with $\mathcal{S}$ and denote its underlying space by $|\mathcal{S}|$.

In the case when $\mathcal{S}$ is (the face poset of) a simplicial complex $\mathcal{K}$ the space $|\mathcal{S}|$ is the geometric realisation $|\mathcal{K}|$.

REMARK. Using a more formal categorical language, we consider the *face category* $\mathrm{CAT}(\mathcal{S})$ whose objects are elements $\sigma \in \mathcal{S}$ and there is a morphism from σ to τ whenever $\sigma \leqslant \tau$. Define a diagram (covariant functor) $\Delta^{\mathcal{S}}$ from $\mathrm{CAT}(\mathcal{S})$ to topological spaces by sending $\sigma \in \mathcal{S}$ to the geometric simplex Δ^σ and sending every morphism $\sigma \leqslant \tau$ to the inclusion $\Delta^\sigma \hookrightarrow \Delta^\tau$. Then we may write

$$|\mathcal{S}| = \operatorname{colim} \Delta^{\mathcal{S}},$$

where the colimit (or direct limit) is taken in the category of topological spaces. This is the first example of colimit construction over the face category $\mathrm{CAT}(\mathcal{S})$. Many other examples of this sort will appear later.

In most circumstances we shall not distinguish between simplicial posets and simplicial cell complexes. We shall also sometimes refer to elements $\sigma \in \mathcal{S}$ as *simplices* or *faces* of $\mathcal{S}$.

EXAMPLE 2.8.2. Consider the simplicial cell complex obtained by attaching two d-dimensional simplices along their boundaries. Its corresponding simplicial poset is not the face poset of a simplicial complex if $d > 0$.

Three cellular subdivisions of a circle are shown on Fig. 2.9. The first is not a simplicial cell complex. The second is a simplicial cell complex, but not a simplicial complex (it corresponds to $d = 1$ in the previous paragraph). The third one is a simplicial complex.



FIGURE 2.9. Cellular subdivisions of a circle.

CONSTRUCTION 2.8.3 (folding a simplicial poset onto a simplicial complex). For every simplicial poset $\mathcal{S}$ there is the associated simplicial complex $\mathcal{K}_{\mathcal{S}}$ on the same vertex set $V(\mathcal{S})$, whose simplices are sets $V(\sigma)$, $\sigma \in \mathcal{S}$. There is a *folding map* of simplicial posets

$$\mathcal{S} \longrightarrow \mathcal{K}_{\mathcal{S}}, \quad \sigma \mapsto V(\sigma). \tag{2.9}$$

It is identical on the vertices, and every simplex in $\mathcal{K}_{\mathcal{S}}$ gets covered by a finite number of simplices of $\mathcal{S}$.

For any two elements $\sigma, \tau \in \mathcal{S}$, denote by $\sigma \vee \tau$ the set of their least common upper bounds (*joins*), and denote by $\sigma \wedge \tau$ the set of their greatest common lower bounds (*meets*). Since $\mathcal{S}$ is a simplicial poset, $\sigma \wedge \tau$ consists of a single element whenever $\sigma \vee \tau$ is nonempty. It is easy to observe that $\mathcal{S}$ is a simplicial complex if and only if for any $\sigma, \tau \in \mathcal{S}$ the set $\sigma \vee \tau$ is either empty or consists of a single element. In this case $\mathcal{S}$ coincides with $\mathcal{K}_{\mathcal{S}}$.

Applying barycentric subdivision to every simplex $\sigma \in \mathcal{S}$ we obtain a new simplicial cell complex $\mathcal{S}'$, called the *barycentric subdivision* of $\mathcal{S}$. From Proposition 2.3.7 it is clear that $\mathcal{S}'$ can be identified with the (geometric realisation of the) order complex $\operatorname{ord}(\mathcal{S} \setminus \hat{0})$. We therefore obtain the following.

Proposition 2.8.4. *The barycentric subdivision $\mathcal{S}'$ of a simplicial cell complex is a simplicial complex.*

Exercises.

2.8.5. Show that the following conditions are equivalent for a simplicial poset $\mathcal{S}$:

(a) $\mathcal{S}$ is (the face poset of) a simplicial complex;
(b) for any $\sigma, \tau \in \mathcal{S}$ the set $\sigma \wedge \tau$ consists of a single element;
(c) for any $\sigma, \tau \in \mathcal{S}$ the set $\sigma \vee \tau$ is either empty or consists of a single element.

2.9. Cubical complexes

At some stage of the development of combinatorial topology, cubical complexes were considered as an alternative to triangulations, a new way to study topological invariants combinatorially. Their nice feature is that the product of cubes is again a cube, which makes subdivisions of products easier and leads to a more straightforward definition of the multiplication in cohomology. Later it turned out, however, that the cubical (co)homomology itself is not particularly advantageous in comparison with the simplicial one. Currently combinatorial geometry is the main field of applications of cubical complexes; moreover, subdivisions into cubes sometimes are very helpful in various geometrical and topological considerations. In this section we collect the necessary definitions and notation, and then proceed to describe some important cubical decompositions of simple polytopes and simplical complexes.

Definitions and examples. As in the case of simplicial complexes, a cubical complex can be defined either abstractly (as a poset) or geometrically (as a cell complex).

Definition 2.9.1. An *abstract cubical complex* is a finite poset $(\mathcal{C}, \subset)$ containing an initial element $\varnothing$ and satisfying the following two conditions:

(a) for every element $G \in \mathcal{C}$ the segment $[\varnothing, G]$ is isomorphic to the face poset of a cube;
(b) for every two elements $G_1, G_2 \in \mathcal{C}$ there is a unique meet (greatest lower bound).

Elements $G \in \mathcal{C}$ are *faces* of the cubical complex. If $[\varnothing, G]$ is the face poset of the k-cube $\mathbb{I}^k$, then the face G is of *k-dimensional.* The dimension of $\mathcal{C}$ is the maximal dimension of its faces. The meet of any two faces G_1, G_2 is also called their *intersection* and denoted $G_1 \cap G_2$.

A d-dimensional *topological cube* is a d-ball with a face structure defined by a homeomorphism with the standard cube $\mathbb{I}^d$. A *face* of a topological d-cube is thus the homeomorphic image of a face of $\mathbb{I}^d$.

Definition 2.9.2. A *topological cubical complex* is a set $\mathcal{U}$ of topological cubes of arbitrary dimensions which are all embedded in the same space $\mathbb{R}^n$ and satisfy the following conditions:

(a) every face of a cube in $\mathcal{U}$ belongs to $\mathcal{U}$;
(b) The intersection of any two cubes in $\mathcal{U}$ is a face of each.

Every abstract cubical complex $\mathcal{C}$ has a *geometric realisation*, a topological cubical complex $\mathcal{U}$ whose faces form a poset isomorphic to $\mathcal{C}$. Such $\mathcal{U}$ can be constructed by taking a disjoint union of topological cubes corresponding to all segments $[\varnothing, G] \subset \mathcal{C}$ and identifying faces according to the poset relation.

From now on we shall not distinguish between abstract cubical complexes and their geometric realisations.

By analogy with simplicial complexes, define the *f-vector* of a cubical complex $\mathcal{C}$ by $\boldsymbol{f}(\mathcal{C}) = (f_0, f_1, \ldots)$ where f_i is the number of i-dimensional faces. There are also notions of h- and g-vectors, and cubical analogues of the UBC, LBC and g-conjecture. See **[2]**, **[10]** and **[209]** for more details and references.

The difference between Definition 2.9.2 of a geometric cubical complex and Definition 2.2.1 of a geometric simplicial complex is that we realise abstract cubes by topological complexes rather than polytopes. This difference is substantial: if we replace topological cubes by combinatorial ones (i.e. by convex polytopes combinatorially equivalent to a cube) in Definition 2.9.2, then we obtain the definition of a *polyhedral cubical complex*. Although this notion is also important in combinatorial geometry, not every abstract cubical complex can be realised by a polyhedral complex, as shown by the next example.

Example 2.9.3. Consider the decomposition of a Möbius strip into 3 squares shown in Fig. 2.10.



Figure 2.10. Cubical complex which does not admit a polyhedral realisation.

Proposition 2.9.4. *The topological cubical complex shown in Fig. 2.10 does not admit a polyhedral cubical realisation.*

Proof. Assume that such a realisation exists. Then since $ABED$ is a convex 4-gon, the points A and D are in the same halfplane with respect to the line BE, and therefore A and D are in the same halfspace defined by the plane BCE. Similarly, since $ABCF$ is a convex 4-gon, the points A and F are in the same halfspace with respect to BCE. Hence, D and F are also in the same halfspace with respect to BCE. On the other hand, since $CDEF$ is a convex 4-gon, the points D and F must be in different subspaces with respect to BCE. A contradiction. $\square$

The example above shows that, unlike the case of simplicial complexes, the theory of abstract cubical complexes cannot be described by using convex-geometric considerations only. Another simple manifestation of this is the fact that not every cubical complex may be realised as a subcomplex in a standard cube, in contrast to simplicial complexes which are always embeddable in a standard simplex. The boundary of a triangle is the simplest example of a cubical complex not embeddable in a cube. It is also not embeddable into the standard cubical lattice in $\mathbb{R}^n$ (for any n). On the other hand, every cubical complex admits a cubical subdivision which is embeddable in a standard cube, as shown in the next subsection. Without subdivision the question of embeddability in a standard cube or cubical lattice is nontrivial. The importance of studying cubical maps (in particular, cubical embeddings) of 2-dimensional cubical complexes to the cubical lattice in $\mathbb{R}^3$ was pointed out by S. Novikov in connection with the 3-*dimensional Ising model*. A significant advance on this problem has been achieved in **[77]**, where necessary and sufficient conditions were obtained for a cubical complex to admit a cubical map to a standard lattice.

Cubical subdivisions of simple polytopes and simplicial complexes. The particular constructions of cubical complexes given here will be important in the definition of moment-angle complexes. Neither of these constructions is particularly new, but they are probably not well recorded in the literature (see however the references at the end of the section).

Any face of $\mathbb{I}^m$ has the form

$$C_{J\subset I}=\{(y_1,\dots,y_m)\in\mathbb{I}^m\colon y_j=0\text{ for }j\in J,\ y_j=1\text{ for }j\notin I\}$$

where $J\subset I$ is a pair of embedded (possibly empty) subsets of $[m]$. We also set

$$C_I=C_{\varnothing\subset I}=\{(y_1,\dots,y_m)\in\mathbb{I}^m\colon y_j=1\text{ for }j\notin I\}\tag{2.10}$$

to simplify the notation.

Construction 2.9.5 (canonical triangulation of $\mathbb{I}^m$). Denote by $\Delta=\Delta^{m-1}$ the simplex on $[m]$. We assign to a subset $I=\{i_1,\dots,i_k\}\subset[m]$ the vertex $v_I=C_{I\subset I}$ of $\mathbb{I}^m$. That is, $v_I=(\varepsilon_1,\dots,\varepsilon_m)$ where $\varepsilon_i=0$ if $i\in I$ and $\varepsilon_i=1$ otherwise. Regarding each I as a vertex of the barycentric subdivision of Δ, we can extend the correspondence $I\mapsto v_I$ to a piecewise linear embedding $i_c\colon\Delta'\to\mathbb{I}^m$. Under this embedding the vertices of Δ are mapped to the vertices of $\mathbb{I}^m$ with exactly one zero coordinate, and the barycentre of Δ is mapped to $(0,\dots,0)\in\mathbb{I}^m$ (see Fig. 2.11). The image $i_c(\Delta')$ is the union of m facets of $\mathbb{I}^m$ meeting at the vertex $(0,\dots,0)$. For each pair $I\subset J$, all simplices of Δ' of the form $I=I_1\subset I_2\subset\cdots\subset I_k=J$ are mapped to the same face $C_{I\subset J}$ of $\mathbb{I}^m$. The map $i_c\colon\Delta'\to\mathbb{I}^m$ extends to $\operatorname{cone}(\Delta')$ by mapping the cone vertex to $(1,\dots,1)\in\mathbb{I}^m$. The image of the resulting map $\operatorname{cone}(i_c)$ is the whole cube $\mathbb{I}^m$. Thus, $\operatorname{cone}(i_c)\colon\operatorname{cone}(\Delta')\to\mathbb{I}^m$ is a PL homeomorphism which is linear on simplices of $\operatorname{cone}(\Delta')$. This defines a canonical triangulation of $\mathbb{I}^m$, the 'triangulation along the main diagonal'.

Subdivisions appeared in the construction above can be summarised as follows:

Proposition 2.9.6. *The PL map* $\operatorname{cone}(i_c)\colon\operatorname{cone}(\Delta')\to\mathbb{I}^m$ *gives rise to*

(a) *cubical subdivision of Δ^{m-1} isomorphic to 'half of the boundary of $\mathbb{I}^m$' (the union of facets of $\mathbb{I}^m$ containing the zero vertex);*

(b) *cubical subdivision of* $\operatorname{cone}\Delta^{m-1}$ *(which is Δ^m) isomorphic to $\mathbb{I}^m$;*



FIGURE 2.11. Taking cone over the barycentric subdivision of simplex defines a triangulation of the cube.

(c) *simplicial subdivision of* $\mathbb{I}^m$ *isomorphic to* $\mathrm{cone}((\Delta^{m-1})')$.

CONSTRUCTION 2.9.7 (cubical subdivision of a simple polytope). Let P be a simple n-polytope with m facets $F_1, \ldots, F_m$. We shall embed P piecewise linearly into the standard cube $\mathbb{I}^m$, thereby inducing a cubical subdivision $\mathcal{C}(P)$ of P by the preimages of faces of $\mathbb{I}^m$.

Denote by $\mathcal{S}$ the set of barycentres of faces of P, including the vertices and the barycentre of the whole polytope. This will be the vertex set of $\mathcal{C}(P)$. Every $(n-k)$-face G of P is an intersection of k facets: $G = F_{i_1} \cap \cdots \cap F_{i_k}$. We map the barycentre of G to the vertex $(\varepsilon_1, \ldots, \varepsilon_m) \in \mathbb{I}^m$, where $\varepsilon_i = 0$ if $i \in \{i_1, \ldots, i_k\}$ and $\varepsilon_i = 1$ otherwise. The resulting map $\mathcal{S} \to \mathbb{I}^m$ can be extended linearly on the simplices of the barycentric subdivision of P to an embedding $c_P\colon P \to \mathbb{I}^m$. The case $n = 2$, $m = 3$ is shown in Fig. 2.12.



FIGURE 2.12. Embedding $c_P\colon P \to \mathbb{I}^m$ for $n = 2$, $m = 3$.

The image $c_P(P) \subset \mathbb{I}^m$ is the union of all faces $C_{J\subset I}$ such that $\bigcap_{i\in I} F_i \neq \varnothing$. For such $C_{J\subset I}$, the preimage $c_P^{-1}(C_{J\subset I})$ is declared to be a face of the cubical complex $\mathcal{C}(P)$. The vertex set of $c_P^{-1}(C_{J\subset I})$ is the subset of $\mathcal{S}$ consisting of barycentres of all faces between the faces G and H of P, where $G = \bigcap_{j\in J} F_j$ and $H = \bigcap_{i\in I} F_i$. Therefore, the faces of $\mathcal{C}(P)$ correspond to pairs of embedded faces $G \supset H$, and we denote them by $C_{G\supset H}$. In particular, maximal (n-dimensional) faces of $\mathcal{C}(P)$ correspond to pairs $G = P$, $H = v$, where v is a vertex of P. For these maximal faces we use the abbreviated notation $C_v = C_{P\supset v}$.

For every vertex $v = F_{i_1} \cap \dots \cap F_{i_n} \in P$ with $I_v = \{i_1, \dots, i_n\}$ we have

$$c_P(C_v) = C_{I_v} = \bigl\{(y_1, \dots, y_m) \in \mathbb{I}^m \colon y_j = 1 \text{ for } j \notin I_v\bigr\}. \tag{2.11}$$

We therefore obtain:

PROPOSITION 2.9.8. *A simple polytope P with m facets admits a cubical decomposition whose maximal faces C_v correspond to the vertices $v \in P$. The resulting cubical complex $\mathcal{C}(P)$ embeds canonically into $\mathbb{I}^m$, as described by* (2.11).

LEMMA 2.9.9. *The number of k-faces of the cubical complex $\mathcal{C}(P)$ is given by*

$$f_k\bigl(\mathcal{C}(P)\bigr) = \sum_{i=0}^{n-k} \binom{n-i}{k} f_i(P), \quad \textit{for } 0 \leqslant k \leqslant n.$$

PROOF. The formula follows from the fact that the k-faces of $\mathcal{C}(P)$ are in one-to-one correspondence with the pairs $G^{i+k} \supset H^i$ of faces of P. □

CONSTRUCTION 2.9.10 (cubical subdivision of a simplicial complex). Let $\mathcal{K}$ be a simplicial complex on $[m]$. Then $\mathcal{K}$ is naturally a subcomplex of Δ^{m-1} and the barycentric subdivision $\mathcal{K}'$ is a subcomplex of $(\Delta^{m-1})'$. Restricting the PL map from Construction 2.9.5 to $\mathcal{K}'$, we obtain the embedding $i_c|_{\mathcal{K}'}\colon |\mathcal{K}'| \to \mathbb{I}^m$. Its image is a cubical subcomplex in $\mathbb{I}^m$, which we denote $\operatorname{cub}(\mathcal{K})$. Then $\operatorname{cub}(\mathcal{K})$ is the union of faces $C_{I\subset J}\subset\mathbb{I}^m$ over all pairs $I\subset J$ of nonempty simplices of $\mathcal{K}$:

$$\operatorname{cub}(\mathcal{K})=\bigcup_{\varnothing\ne I\subset J\in\mathcal{K}}C_{I\subset J}\subset\mathbb{I}^m. \tag{2.12}$$

Construction 2.9.11. Since $\operatorname{cone}(\mathcal{K}')$ is a subcomplex of $\operatorname{cone}((\Delta^{m-1})')$, Construction 2.9.5 also provides a PL embedding

$$\operatorname{cone}(i_c)|_{\operatorname{cone}(\mathcal{K}')}\colon|\operatorname{cone}(\mathcal{K}')|\to\mathbb{I}^m.$$

The image of this embedding is an n-dimensional cubical subcomplex of $\mathbb{I}^m$, which we denote $\operatorname{cc}(\mathcal{K})$. It is easy to see that

$$\operatorname{cc}(\mathcal{K})=\bigcup_{I\subset J\in\mathcal{K}}C_{I\subset J}=\bigcup_{J\in\mathcal{K}}C_J. \tag{2.13}$$

Remark. If $i\in[m]$ is not a vertex of $\mathcal{K}$ (a ghost vertex), then $\operatorname{cc}(\mathcal{K})$ is contained in the facet $\{y_i=1\}$ of $\mathbb{I}^m$.

The following statement summarises the results of two previous constructions.

Proposition 2.9.12. *For any simplicial complex $\mathcal{K}$ on the set $[m]$, there is a PL embedding of $|\mathcal{K}|$ into $\mathbb{I}^m$ linear on the simplices of $\mathcal{K}'$. The image of this embedding is the cubical subcomplex* (2.12)*. Moreover, there is a PL embedding of $|\operatorname{cone}\mathcal{K}|$ into $\mathbb{I}^m$ linear on the simplices of* $\operatorname{cone}(\mathcal{K}')$*, whose image is the cubical subcomplex* (2.13)*.*

A cubical complex $\mathcal{C}'$ is called a *cubical subdivision* of a cubical complex $\mathcal{C}$ if each face of $\mathcal{C}'$ is contained in a face of $\mathcal{C}$, and each face of $\mathcal{C}$ is a union of finitely many faces of $\mathcal{C}'$.

Proposition 2.9.13. *For every cubical complex $\mathcal{C}$ with q vertices, there exists a cubical subdivision that is embeddable as a subcomplex in $\mathbb{I}^q$.*

Proof. We first construct a simplicial complex $\mathcal{K}_\mathcal{C}$ which subdivides the cubical complex $\mathcal{C}$ and has the same vertices. This can be done by induction on the skeleta of $\mathcal{C}$, by extending the triangulation from the k-dimensional skeleton to the interiors of $(k+1)$-dimensional faces using a generic convex function $f\colon\mathbb{R}^k\to\mathbb{R}$, see [**228**, §5.1] (note that the 1-skeleton of $\mathcal{C}$ is already a simplicial complex). Then applying Construction 2.9.10 to $\mathcal{K}_\mathcal{C}$ we get cubical complex $\operatorname{cub}(\mathcal{K}_\mathcal{C})$ that subdivides $\mathcal{K}_\mathcal{C}$ and therefore $\mathcal{C}$. It is embeddable into $\mathbb{I}^q$ by Proposition 2.9.12. □

Example 2.9.14. Cubical complexes $\operatorname{cub}(\mathcal{K})$ and $\operatorname{cc}(\mathcal{K})$ when $\mathcal{K}$ is a disjoint union of 3 vertices or the boundary of a triangle are shown in Figs. 2.13–2.14.

Remark. Let P be a simple n-polytope, and let $\mathcal{K}_P$ be its nerve complex. Then $\operatorname{cc}(\mathcal{K}_P)=c_P(P)$, i.e. $\operatorname{cc}(\mathcal{K}_P)$ coincides with the cubical complex $\mathcal{C}(P)$ from Construction 2.9.7.

Different versions of Construction 2.9.11 can be found in [**10**, p. 299]. A similar construction was also considered in [**70**, p. 434]. Finally, a version of cubical subcomplex $\operatorname{cub}(\mathcal{K})\subset I^m$ appeared in [**77**] in connection with the problems described in the end of the previous subsection.



(a) $\mathcal{K} = 3$ points (b) $\mathcal{K} = \partial\Delta^2$

FIGURE 2.13. Cubical complex cub($\mathcal{K}$).



(a) $\mathcal{K} = 3$ points (b) $\mathcal{K} = \partial\Delta^2$

FIGURE 2.14. Cubical complex cc($\mathcal{K}$).

Exercises.

2.9.15. Show that the triangulation of $\mathbb{I}^m$ from Construction 2.9.5 coincides with the triangulation of the product of m one-dimensional simplices from Construction 2.2.10.

CHAPTER 3

Combinatorial algebra of face rings

With the appearance of the face rings in the beginning of the 1970s in the work of Reisner and Stanley many combinatorial problems were translated into the language of commutative algebra, which paved the way for their solution using the extensive machinery of algebraic and homological methods. Algebraic tools used for attacking combinatorial problems included regular sequences, Cohen–Macaulay and Gorenstein rings, Tor-algebras, local cohomology, etc. A whole new thriving field appeared on the borders of combinatorics and algebra, which has since become known as *combinatorial commutative algebra*. The basic reference here is Stanley's monograph [**208**].

In this chapter we collect the wealth of algebraic notions and constructions related to face rings. Our choice of material and notation was guided by the topological applications in the later chapters of the book. (This explains the unusual for algebraists even grading in the polynomial rings and their homogeneous quotients, and also the nonpositive homological grading in free resolutions and Tor.) In the first sections we review standard results and constructions of combinatorial commutative algebra, including the Tor-algebras and algebraic Betti numbers of face rings, Cohen–Macaulay and Gorenstein complexes. The later sections contain some more recent developments, including the face rings of simplicial posets, different characterisations of Cohen–Macaulay and Gorenstein simplicial posets in terms of their face rings and h-vectors, and generalisations of the Dehn–Sommerville relations. Although all these algebraic and combinatorial results have a strong topological flavour and were indeed originally motivated by topological constructions, we tried to keep this chapter mostly algebraic and do not require much topological knowledge from the reader here.

The preliminary algebraic material of a more general sort, not directly or exclusively related to the face rings (such as resolutions and the functor Tor, and Cohen–Macaulay rings) is collected in Appendices at the end of the book.

Alongside the above mentioned monograph by Stanley [**208**], an extensive survey of Cohen–Macaulay rings by Bruns and Herzog [**35**] and a more recent monograph [**160**] by Miller and Sturmfels may be recommended for a deeper study of algebraic methods in combinatorics.

We use the common notation $\mathbf{k}$ for the ground ring, which is always assumed to be the ring $\mathbb{Z}$ of integers or a field. The former is preferable for topological applications, but the latter is more common in the algebraic literature. We shall often refer to $\mathbf{k}$-modules as '$\mathbf{k}$-vector spaces'; in the case $\mathbf{k} = \mathbb{Z}$ the latter means 'abelian groups'.

We assume graded commutativity instead of commutativity; algebras commutative in the standard sense will be those whose nontrivial graded components appear only in even degrees. In particular, the polynomial algebra $\mathbf{k}[v_1,\dots,v_m]$, which we often abbreviate to $\mathbf{k}[m]$, has $\deg v_i = 2$. The exterior algebra $\Lambda[u_1,\dots,u_m]$ has $\deg u_i = 1$. Given a subset $I = \{i_1,\dots,i_k\} \subset [m]$ we denote by v_I the square-free monomial $v_{i_1}\cdots v_{i_k}$ in $\mathbf{k}[m]$. We also denote by u_I the exterior monomial $u_{i_1}\cdots u_{i_k}$ where $i_1 < \dots < i_k$.

3.1. Face rings of simplicial complexes

Definition 3.1.1. The *face ring* (or the *Stanley–Reisner ring*) of a simplicial complex $\mathcal{K}$ on the set $[m]$ is the quotient graded ring

$$\mathbf{k}[\mathcal{K}] = \mathbf{k}[v_1,\dots,v_m]/\mathcal{I}_{\mathcal{K}},$$

where $\mathcal{I}_{\mathcal{K}} = (v_I\colon I \notin \mathcal{K})$ is the ideal generated by those monomials v_I for which I is not a simplex of $\mathcal{K}$. The ideal $\mathcal{I}_{\mathcal{K}}$ is known as the *Stanley–Reisner ideal* of $\mathcal{K}$.

Example 3.1.2.

1. Let $\mathcal{K}$ be the 2-dimensional simplicial complex shown in Fig. 3.1. Then

$$\mathcal{I}_{\mathcal{K}} = (v_1v_5, v_3v_4, v_1v_2v_3, v_2v_4v_5).$$



Figure 3.1.

2. The face ring $\mathbf{k}[\mathcal{K}]$ is a *quadratic algebra* (that is, the ideal $\mathcal{I}_{\mathcal{K}}$ is generated by quadratic monomials) if and only if $\mathcal{K}$ is a flag complex (an exercise).

3. Let $\mathcal{K}_1 * \mathcal{K}_2$ be the join of $\mathcal{K}_1$ and $\mathcal{K}_2$ (see Construction 2.2.8). Then

$$\mathbf{k}[\mathcal{K}_1 * \mathcal{K}_2] = \mathbf{k}[\mathcal{K}_1] \otimes \mathbf{k}[\mathcal{K}_2].$$

Here and below $\otimes$ denotes the tensor product over $\mathbf{k}$.

We note that $\mathcal{I}_{\mathcal{K}}$ is a *monomial ideal*, and it has a basis consisting of square-free monomials v_I corresponding to the missing faces of $\mathcal{K}$.

Proposition 3.1.3. *Every square-free monomial ideal $\mathcal{I}$ in the polynomial ring is the Stanley–Reisner ideal of a simplicial complex $\mathcal{K}$.*

Proof. We set

$$\mathcal{K} = \{I \subset [m]\colon v_I \notin \mathcal{I}\}.$$

Then $\mathcal{K}$ is a simplicial complex and $\mathcal{I} = \mathcal{I}_{\mathcal{K}}$. □

Let P be a simple n-polytope and let $\mathcal{K}_P$ be its nerve complex (see Example 2.2.4). We define the *face ring* $\mathbf{k}[P]$ as the face ring of $\mathcal{K}_P$. Explicitly,

$$\mathbf{k}[P]=\mathbf{k}[v_1,\ldots,v_m]/\mathcal{I}_P,$$

where $\mathcal{I}_P$ is the ideal generated by those square-free monomials $v_{i_1}v_{i_2}\cdots v_{i_s}$ whose corresponding facets intersect trivially, $F_{i_1}\cap\cdots\cap F_{i_s}=\varnothing$.

Example 3.1.4.

1. Let P be an n-simplex (viewed as a simple polytope). Then

$$\mathbf{k}[P]=\mathbf{k}[v_1,\ldots,v_{n+1}]/(v_1v_2\cdots v_{n+1}).$$

2. Let P be a 3-cube I^3. Then

$$\mathbf{k}[P]=\mathbf{k}[v_1,v_2,\ldots,v_6]/(v_1v_4,v_2v_5,v_3v_6).$$

3. Let P be an m-gon, $m\geqslant 4$. Then

$$\mathcal{I}_P=(v_iv_j\colon i-j\neq 0,\pm1 \mod m).$$

4. Given two simple polytopes P_1 and P_2, we have

$$\mathbf{k}[P_1\times P_2]=\mathbf{k}[P_1]\otimes\mathbf{k}[P_2].$$

Proposition 3.1.5. *Let $\varphi\colon\mathcal{K}\to\mathcal{L}$ be a simplicial map between simplicial complexes $\mathcal{K}$ and $\mathcal{L}$ on the vertex sets $[m]$ and $[l]$ respectively. Define the map $\varphi^*\colon\mathbf{k}[w_1,\ldots,w_l]\to\mathbf{k}[v_1,\ldots,v_m]$ by*

$$\varphi^*(w_j)=\sum_{i\in\varphi^{-1}(j)}v_i.$$

Then φ^* descends to a homomorphism $\mathbf{k}[\mathcal{L}]\to\mathbf{k}[\mathcal{K}]$, which we continue to denote φ^*.

Proof. We only need to check that $\varphi^*(\mathcal{I}_\mathcal{L})\subset\mathcal{I}_\mathcal{K}$. Suppose $J=\{j_1,\ldots,j_s\}\subset[l]$ is not a simplex of $\mathcal{L}$. We have

$$\varphi^*(w_{j_1}\ldots w_{j_s})=\sum_{i_1\in\varphi^{-1}(j_1),\ldots,i_s\in\varphi^{-1}(j_s)}v_{i_1}\ldots v_{i_s}.$$

We claim that the right hand side above belongs to $\mathcal{I}_\mathcal{K}$, i.e. for any monomial $v_{i_1}\ldots v_{i_s}$ in the right hand side the set $I=\{i_1,\ldots,i_s\}$ is not a simplex of $\mathcal{K}$. Indeed, otherwise we would have $\varphi(I)=J\in\mathcal{L}$ by the definition of a simplicial map, which contradicts the assumption. $\square$

Example 3.1.6. The face ring of the barycentric subdivision $\mathcal{K}'$ of $\mathcal{K}$ is

$$\mathbf{k}[\mathcal{K}']=\mathbf{k}[b_I\colon I\in\mathcal{K}\setminus\varnothing]/\mathcal{I}_{\mathcal{K}'},$$

where b_I is the polynomial generator of degree 2 corresponding to a nonempty simplex $I\in\mathcal{K}$, and $\mathcal{I}_{\mathcal{K}'}$ is generated by quadratic monomials b_Ib_J for which $I\not\subset J$ and $J\not\subset I$. The simplicial map $\nabla\colon\mathcal{K}'\to\mathcal{K}$ from Example 2.3.3 induces a map ∇^* of the face ring, given on the generators $v_j\in\mathbf{k}[\mathcal{K}]$ by

$$\nabla^*(v_j)=\sum_{I\in\mathcal{K}\colon\min I=j}b_I.$$

Example 3.1.7. The nondegenerate map $\mathcal{K}' \to \Delta^{n-1}$ from Example 2.3.2 induces the following map of the corresponding face rings:

$$\begin{aligned} \mathbf{k}[v_1, \ldots, v_n] &\longrightarrow \mathbf{k}[\mathcal{K}'] \\ v_i &\longmapsto \sum_{|I|=i} b_I. \end{aligned}$$

This defines a canonical $\mathbf{k}[v_1, \ldots, v_n]$-module structure on $\mathbf{k}[\mathcal{K}']$.

An important tool arising from the functoriality of the face ring is the restriction homomorphism. For any simplex $I \in \mathcal{K}$, the corresponding full subcomplex $\mathcal{K}_I$ is $\Delta^{|I|-1}$ and $\mathbf{k}[\mathcal{K}_I]$ is the polynomial ring $\mathbf{k}[v_i \colon i \in I]$ on $|I|$ generators. The inclusion $\mathcal{K}_I \subset \mathcal{K}$ induces the *restriction homomorphism*

$$s_I \colon \mathbf{k}[\mathcal{K}] \to \mathbf{k}[v_i \colon i \in I],$$

which maps v_i to zero whenever $i \notin I$.

The following simple proposition will be used in several algebraic and topological arguments of the later chapters.

Proposition 3.1.8. *The direct sum*

$$s = \bigoplus_{I \in \mathcal{K}} s_I \colon \mathbf{k}[\mathcal{K}] \longrightarrow \bigoplus_{I \in \mathcal{K}} \mathbf{k}[v_i \colon i \in I]$$

of all restriction maps is a monomorphism.

Proof. Consider the composite map

$$\mathbf{k}[v_1, \ldots, v_m] \xrightarrow{\ p\ } \mathbf{k}[\mathcal{K}] \xrightarrow{\ s\ } \textstyle\bigoplus_{I \in \mathcal{K}} \mathbf{k}[v_i \colon i \in I]$$

where p is the quotient projection. Suppose $s\,p(Q) = 0$ where $Q = Q(v_1, \ldots, v_m)$ is a polynomial. Then for any monomial $v_{i_1}^{\alpha_1} \cdots v_{i_k}^{\alpha_k}$ which enters Q with a nonzero coefficient we have $I = \{i_1, \ldots, i_k\} \notin \mathcal{K}$ (as otherwise the Ith component of the image under $s\,p$ is nonzero). Hence $p(Q) = 0$ and s is injective. □

Proposition 3.1.9. *The face ring $\mathbf{k}[\mathcal{K}]$ has the $\mathbf{k}$-vector space basis consisting of monomials $v_{j_1}^{\alpha_1} \cdots v_{j_k}^{\alpha_k}$ where $\alpha_i > 0$ and $\{j_1, \ldots, j_k\} \in \mathcal{K}$.*

Proof. Indeed, the polynomial algebra $\mathbf{k}[m]$ has the $\mathbf{k}$-vector space basis consisting of all monomials $v_{j_1}^{\alpha_1} \cdots v_{j_k}^{\alpha_k}$, and such a monomial maps to zero under the projection $\mathbf{k}[m] \to \mathbf{k}[\mathcal{K}]$ precisely when $\{j_1, \ldots, j_k\} \notin \mathcal{K}$. □

Recall that the Poincaré series of a nonnegatively graded $\mathbf{k}$-vector space $V = \bigoplus_{i=0}^{\infty} V^i$ is given by $F(V; \lambda) = \sum_{i=0}^{\infty} (\dim_{\mathbf{k}} V^i) \lambda^i$. Since $\mathbf{k}[\mathcal{K}]$ is graded by even integers, its Poincaré series is even.

Theorem 3.1.10 (Stanley). *Let $\mathcal{K}$ be an $(n-1)$-dimensional simplicial complex with f-vector $(f_0, \ldots, f_{n-1})$ and h-vector $(h_0, \ldots, h_n)$. Then the Poincaré series of the face ring $\mathbf{k}[\mathcal{K}]$ is*

$$F\bigl(\mathbf{k}[\mathcal{K}]; \lambda\bigr) = \sum_{k=0}^{n} f_{k-1} \left(\frac{\lambda^2}{1-\lambda^2} \right)^k = \frac{h_0 + h_1 \lambda^2 + \cdots + h_n \lambda^{2n}}{(1-\lambda^2)^n}.$$

Proof. By Proposition 3.1.9, a $(k-1)$-dimensional simplex $\{i_1, \ldots, i_k\} \in \mathcal{K}$ contributes a summand $\frac{\lambda^{2k}}{(1-\lambda^2)^k}$ to the Poincaré series of $\mathcal{K}$ (this summand is just the Poincaré series of the subspace generated by monomials $v_{i_1}^{\alpha_1} \cdots v_{i_k}^{\alpha_k}$ with positive exponents α_i). This proves the first identity, and the second follows from (2.3). □

EXAMPLE 3.1.11.

1. Let $\mathcal{K} = \Delta^{n-1}$. Then $f_i = \binom{n}{i+1}$ for $-1 \leqslant i \leqslant n-1$, $h_0 = 1$ and $h_i = 0$ for $i > 0$. Since every subset of $[n]$ is a simplex of Δ^{n-1}, we have $\mathbf{k}[\Delta^{n-1}] = \mathbf{k}[v_1, \ldots, v_n]$ and $F(\mathbf{k}[\Delta^{n-1}]; \lambda) = (1-\lambda^2)^{-n}$.

2. Let $\mathcal{K} = \partial\Delta^n$ be the boundary of an n-simplex. Then $h_i = 1$ for $0 \leqslant i \leqslant n$, and $\mathbf{k}[\partial\Delta^n] = \mathbf{k}[v_1, \ldots, v_{n+1}]/(v_1 v_2 \cdots v_{n+1})$. By Theorem 3.1.10,

$$F\bigl(\mathbf{k}[\partial\Delta^n]; \lambda\bigr) = \frac{1 + \lambda^2 + \cdots + \lambda^{2n}}{(1-\lambda^2)^n}.$$

The affine algebraic variety corresponding to the commutative finitely generated $\mathbf{k}$-algebra $\mathbf{k}[\mathcal{K}] = \mathbf{k}[m]/\mathcal{I}_{\mathcal{K}}$ (i.e. the set of common zeros of elements of $\mathcal{I}_{\mathcal{K}}$, viewed as algebraic functions on $\mathbf{k}^m$) can be easily identified as follows.

PROPOSITION 3.1.12. *The affine variety corresponding to* $\mathbf{k}[\mathcal{K}]$ *is given by*

$$X(\mathcal{K}) = \bigcup_{I \in \mathcal{K}} S_I,$$

where $S_I = \mathbf{k}\langle \boldsymbol{e}_i \colon i \in I\rangle$ *is the coordinate subspace in* $\mathbf{k}^m$ *spanned by the set of standard basis vectors corresponding to* I.

PROOF. The statement obviously holds in the case $\mathcal{K} = \Delta^{m-1}$. So we assume $\mathcal{K} \neq \Delta^{m-1}$. We shall use the following notation from Section 2.4: $\widehat{I} = [m] \setminus I$, the complement of $I \subset [m]$, and $\widehat{\mathcal{K}} = \{\widehat{I} \in [m] \colon I \notin \mathcal{K}\}$, the dual complex of $\mathcal{K}$. Given a point $\boldsymbol{z} = (z_1, \ldots, z_m) \in \mathbf{k}^m$, we denote by

$$\omega(\boldsymbol{z}) = \{i \colon z_i = 0\} \subset [m],$$

the set of zero coordinates of $\boldsymbol{z}$.

By the definition of the algebraic variety $X(\mathcal{K})$ corresponding to $\mathbf{k}[\mathcal{K}]$,

$$\begin{aligned} X(\mathcal{K}) = \bigcap_{J \notin \mathcal{K}} \bigcup_{j \in J} \{\boldsymbol{z} \colon z_j = 0\} &= \bigcap_{J \notin \mathcal{K}} \bigl\{\boldsymbol{z} \colon \omega(\boldsymbol{z}) \cap J \neq \varnothing\bigr\} \\ &= \bigcap_{\widehat{J} \in \widehat{\mathcal{K}}} \bigl\{\boldsymbol{z} \colon \omega(\boldsymbol{z}) \not\subset \widehat{J}\bigr\} = \bigl\{\boldsymbol{z} \colon \omega(\boldsymbol{z}) \notin \widehat{\mathcal{K}}\bigr\}. \end{aligned}$$

On the other hand,

$$\begin{aligned} \bigcup_{I \in \mathcal{K}} S_I = \bigcup_{I \in \mathcal{K}} \bigcap_{j \in \widehat{I}} \{\boldsymbol{z} \colon z_j = 0\} &= \bigcup_{I \in \mathcal{K}} \bigl\{\boldsymbol{z} \colon \widehat{I} \subset \omega(\boldsymbol{z})\bigr\} \\ &= \bigcup_{\widehat{I} \notin \widehat{\mathcal{K}}} \bigl\{\boldsymbol{z} \colon \omega(\boldsymbol{z}) \supset \widehat{I}\bigr\} = \bigl\{\boldsymbol{z} \colon \omega(\boldsymbol{z}) \notin \widehat{\mathcal{K}}\bigr\}. \end{aligned}$$

The required identity follows by comparing the two formulae above. □

REMARK. The variety $X(\mathcal{K})$ is an example of an *arrangement of coordinate subspaces*, which will be studied further in Section 4.6.

We finish this section with a result showing that the face ring determines its underlying simplicial complex:

THEOREM 3.1.13 (Bruns–Gubeladze [**34**]). *Let* $\mathbf{k}$ *be a field, and* $\mathcal{K}_1$ *and* $\mathcal{K}_2$ *be two simplicial complexes on the vertex sets* $[m_1]$ *and* $[m_2]$ *respectively. Suppose* $\mathbf{k}[\mathcal{K}_1]$ *and* $\mathbf{k}[\mathcal{K}_2]$ *are isomorphic as* $\mathbf{k}$*-algebras. Then there exists a bijective map* $[m_1] \to [m_2]$ *which induces an isomorphism between* $\mathcal{K}_1$ *and* $\mathcal{K}_2$.

Proof. Let $f\colon \mathbf{k}[\mathcal{K}_1] \to \mathbf{k}[\mathcal{K}_2]$ be an isomorphism of $\mathbf{k}$-algebras. An easy argument shows that we can assume that f is a graded isomorphism (an exercise, or see [**34**, p. 316]).

Since f is graded, by restriction to the linear components we observe that $m_1 = m_2$ and that f is induced by a linear isomorphism $F\colon \mathbf{k}[m_1] \to \mathbf{k}[m_2]$. This is described by the commutative diagram

$$\begin{tikzcd}
\mathbf{k}[v_1,\dots,v_{m_1}] \arrow[r, "F"] \arrow[d] & \mathbf{k}[v_1,\dots,v_{m_2}] \arrow[d] \\
\mathbf{k}[\mathcal{K}_1] \arrow[r, "f"] & \mathbf{k}[\mathcal{K}_2]
\end{tikzcd}$$

By passing to the associated affine varieties, we observe that the isomorphism $f^*\colon X(\mathcal{K}_2) \to X(\mathcal{K}_1)$ is the restriction of the $\mathbf{k}$-linear isomorphism $F^*\colon \mathbf{k}^{m_2} \to \mathbf{k}^{m_1}$. This is described by the commutative diagram

$$\begin{tikzcd}
\mathbf{k}^{m_2} \arrow[r, "F^*"] & \mathbf{k}^{m_1} \\
X(\mathcal{K}_2) \arrow[r, "f^*"] \arrow[u] & X(\mathcal{K}_1) \arrow[u]
\end{tikzcd}$$

The isomorphism f^* establishes a bijective correspondence

$$\varPhi\colon \{\text{maximal faces of } \mathcal{K}_2\} \to \{\text{maximal faces of } \mathcal{K}_1\}$$

which is defined by the formula $f^*(S_I) = S_{\varPhi(I)}$, where I is a maximal face of $\mathcal{K}_2$. It is also clear that $|\varPhi(I)| = |I| = \dim S_I$.

We denote by $\mathcal{P}_1$ the intersection poset of the subspaces S_I, $I \in \mathcal{K}_1$, with respect to inclusion (i.e. the elements of $\mathcal{P}_1$ are nonempty intersections $S_{I_1}\cap\dots\cap S_{I_k}$ with $I_j \in \mathcal{K}_1$). The poset $\mathcal{P}_1$ can be also viewed as the intersection poset of the maximal faces of $\mathcal{K}_1$. We define the poset $\mathcal{P}_2$ corresponding to $\mathcal{K}_2$ similarly. The correspondence $\varPhi$ obviously extends to an isomorphism of posets $\varPhi\colon \mathcal{P}_2 \to \mathcal{P}_1$, which preserves the dimension of spaces (or the number of elements in the intersections of maximal faces).

Now introduce the following equivalence relation on the vertex sets $[m_1]$ and $[m_2]$: for $i_1, i_2 \in [m_1]$ (or $j_1, j_2 \in [m_2]$) we put $i_1 \sim i_2$ if and only if the two sets of maximal faces $\mathcal{K}_1$ containing i_1 and i_2 respectively coincide (and similarly for j_1 and j_2). The equivalence classes in $[m_1]$ are the minimal (with respect to inclusion) nonempty intersections of maximal faces of $\mathcal{K}_1$, and similarly for $[m_2]$. Since $\varPhi$ is an isomorphism of posets, the two systems of equivalence classes are in natural bijective correspondence, and the corresponding equivalence classes have the same numbers of elements. This gives rise to the bijective map $\varphi\colon [m_2] \to [m_1]$ which satisfies the condition that $i \in I$ if and only if $\varphi(i) \in \varPhi(I)$, where $i \in [m_2]$ and $I \in \mathcal{K}_2$ is a maximal face. Since any face of a simplicial complex is contained in a maximal face, we obtain that $\psi = \varphi^{-1}\colon [m_1] \to [m_2]$ is the required map. $\square$

Exercises.

3.1.14. Show that the Stanley–Reisner ideal $\mathcal{I}_{\mathcal{K}}$ is generated by quadratic monomials if and only if $\mathcal{K}$ is a flag complex.

3.1.15 (see [**183**, (4.7)]). Let $\mathrm{CAT}(\mathcal{K})$ be the face category of $\mathcal{K}$ (objects are simplices, morphisms are inclusions), $\mathrm{CAT}^{op}(\mathcal{K})$ the opposite category (in which the morphisms are reverted), and CGA the category of commutative graded algebras. Consider the diagram

$$\begin{aligned}\mathbf{k}[\cdot]_{\mathcal{K}}\colon \mathrm{CAT}^{op}(\mathcal{K}) &\longrightarrow \mathrm{CGA},\\ I &\longmapsto \mathbf{k}[v_i\colon i\in I]\end{aligned}$$

whose value on a morphism $I \subset J$ is the surjection $\mathbf{k}[v_j\colon j\in J]\to\mathbf{k}[v_i\colon i\in I]$ sending each v_j with $j\notin I$ to zero. Show that

$$\mathbf{k}[\mathcal{K}] = \lim \mathbf{k}[\cdot]_{\mathcal{K}}$$

where the limit is taken in the category CGA. (See Appendix C for the details on categorical constructions.)

3.1.16. If $\mathbf{k}[\mathcal{K}_1]$ and $\mathbf{k}[\mathcal{K}_2]$ are isomorphic as $\mathbf{k}$-algebras, then there is also a graded isomorphism $\mathbf{k}[\mathcal{K}_1]\to\mathbf{k}[\mathcal{K}_2]$. (Hint: Show first that $\mathbf{k}[\mathcal{K}_1]$ and $\mathbf{k}[\mathcal{K}_2]$ are isomorphic as augmented $\mathbf{k}$-algebras, and then pass to the associated graded algebras with respect to the augmentation ideals.)

3.2. Tor-algebras and Betti numbers

The algebraic Betti numbers of the face ring $\mathbf{k}[\mathcal{K}]$ are the dimensions of the Tor-groups of $\mathbf{k}[\mathcal{K}]$ viewed as a module over the polynomial ring. These basic homological invariants of a simplicial complex $\mathcal{K}$ appear to be of great importance both for combinatorial commutative algebra and toric topology.

The face ring $\mathbf{k}[\mathcal{K}]$ acquires a canonical $\mathbf{k}[m]$-module structure via the quotient projection $\mathbf{k}[m]\to\mathbf{k}[\mathcal{K}]$. We therefore may consider the corresponding Tor-modules (see Appendix, Section A.2):

$$\mathrm{Tor}_{\mathbf{k}[v_1,\dots,v_m]}\bigl(\mathbf{k}[\mathcal{K}],\mathbf{k}\bigr) = \bigoplus_{i,j\geqslant 0}\mathrm{Tor}^{-i,2j}_{\mathbf{k}[v_1,\dots,v_m]}\bigl(\mathbf{k}[\mathcal{K}],\mathbf{k}\bigr).$$

From Lemma A.2.10 we obtain that $\mathrm{Tor}_{\mathbf{k}[m]}(\mathbf{k}[\mathcal{K}],\mathbf{k})$ is a bigraded algebra in a natural way, and there is the following isomorphism of bigraded algebras:

$$\mathrm{Tor}_{\mathbf{k}[v_1,\dots,v_m]}\bigl(\mathbf{k}[\mathcal{K}],\mathbf{k}\bigr)\cong H\bigl[\Lambda[u_1,\dots,u_m]\otimes\mathbf{k}[\mathcal{K}],d\bigr],$$

where the bigrading and differential on the right hand side are given by

$$\begin{aligned}\operatorname{bideg}u_i=(-1,2),&\quad \operatorname{bideg}v_i=(0,2),\\ du_i=v_i,&\quad dv_i=0.\end{aligned}\tag{3.1}$$

Definition 3.2.1. We refer to $\mathrm{Tor}_{\mathbf{k}[v_1,\dots,v_m]}(\mathbf{k}[\mathcal{K}],\mathbf{k})$ as the Tor-*algebra* of a simplicial complex $\mathcal{K}$.

The *bigraded Betti numbers* of $\mathbf{k}[\mathcal{K}]$ are defined by

$$\beta^{-i,2j}\bigl(\mathbf{k}[\mathcal{K}]\bigr)=\dim_{\mathbf{k}}\mathrm{Tor}^{-i,2j}_{\mathbf{k}[v_1,\dots,v_m]}\bigl(\mathbf{k}[\mathcal{K}],\mathbf{k}\bigr),\qquad\text{for } i,j\geqslant 0.\tag{3.2}$$

We also set

$$\beta^{-i}\bigl(\mathbf{k}[\mathcal{K}]\bigr)=\dim_{\mathbf{k}}\mathrm{Tor}^{-i}_{\mathbf{k}[v_1,\dots,v_m]}\bigl(\mathbf{k}[\mathcal{K}],\mathbf{k}\bigr)=\sum_j\beta^{-i,2j}\bigl(\mathbf{k}[\mathcal{K}]\bigr).$$

The Tor-algebra has the following functorial property:

PROPOSITION 3.2.2. *A simplicial map* $\varphi\colon \mathcal{K} \to \mathcal{L}$ *between simplicial complexes on the sets* $[m]$ *and* $[l]$ *respectively induces a homomorphism*

$$\varphi^*_{\mathrm{Tor}}\colon\ \mathrm{Tor}_{\mathbf{k}[w_1,\ldots,w_l]}\big(\mathbf{k}[\mathcal{L}],\mathbf{k}\big) \to \mathrm{Tor}_{\mathbf{k}[v_1,\ldots,v_m]}\big(\mathbf{k}[\mathcal{K}],\mathbf{k}\big)$$

of the corresponding Tor-*algebras.*

PROOF. This follows from Proposition 3.1.5 and Theorem A.2.5 (b). □

Consider the minimal resolution $[R_{\min}, d]$ of the $\mathbf{k}[m]$-module $\mathbf{k}[\mathcal{K}]$ (see Construction A.2.2). Then $R^0_{\min} \cong 1 \cdot \mathbf{k}[m]$ is a free module with one generator of degree 0. The basis of $R^{-1}_{\min}$ is a minimal generator set for $\mathcal{I}_{\mathcal{K}}$, and these minimal generators correspond to the missing faces of $\mathcal{K}$. Given a missing face $\{i_1,\ldots,i_k\} \subset [m]$, denote by $r_{i_1,\ldots,i_k}$ the corresponding generator of $R^{-1}_{\min}$. Then the map $d\colon R^{-1}_{\min} \to R^0_{\min}$ takes $r_{i_1,\ldots,i_k}$ to $v_{i_1}\ldots v_{i_k}$. By Proposition A.2.6, $\beta^{-1,2j}(\mathbf{k}[\mathcal{K}])$ is equal to the number of missing faces with j elements.

EXAMPLE 3.2.3. Let $\mathcal{K} =$ 

, the boundary of a 4-gon. Then

$$\mathbf{k}[\mathcal{K}] \cong \mathbf{k}[v_1,\ldots,v_4]/(v_1v_3, v_2v_4).$$

Let us construct a minimal resolution of $\mathbf{k}[\mathcal{K}]$. The module $R^0_{\min}$ has one generator 1 (of degree 0). The module $R^{-1}_{\min}$ has two generators r_{13} and r_{24} of degree 4, and the differential $d\colon R^{-1}_{\min} \to R^0_{\min}$ takes r_{13} to v_1v_3 and r_{24} to v_2v_4. The kernel $R^{-1}_{\min} \to R^0_{\min}$ is generated by one element $v_2v_4r_{13} - v_1v_3r_{24}$. Hence, $R^{-2}_{\min}$ has one generator of degree 8, which we denote by a, and the map $d\colon R^{-2}_{\min} \to R^{-1}_{\min}$ is injective and takes a to $v_2v_4r_{13} - v_1v_3r_{24}$. Thus, the minimal resolution is

$$0 \longrightarrow R^{-2}_{\min} \longrightarrow R^{-1}_{\min} \longrightarrow R^0_{\min} \longrightarrow M \longrightarrow 0,$$

where $\operatorname{rank} R^0_{\min} = \beta^{0,0}(\mathbf{k}[\mathcal{K}]) = 1$, $\operatorname{rank} R^{-1}_{\min} = \beta^{-1,4}(\mathbf{k}[\mathcal{K}]) = 2$, $\operatorname{rank} R^{-2}_{\min} = \beta^{-2,8}(\mathbf{k}[\mathcal{K}]) = 1$.

The following fundamental result of Hochster reduces the calculation of the Betti numbers $\beta^{-i,2j}(\mathbf{k}[\mathcal{K}])$ to the calculation of reduced simplicial cohomology of full subcomplexes in $\mathcal{K}$.

THEOREM 3.2.4 (Hochster [**124**]). *We have*

$$\mathrm{Tor}^{-i,2j}_{\mathbf{k}[v_1,\ldots,v_m]}\big(\mathbf{k}[\mathcal{K}],\mathbf{k}\big) = \bigoplus_{J\subset[m]\colon\,|J|=j} \widetilde{H}^{j-i-1}(\mathcal{K}_J;\mathbf{k}),$$

where $\mathcal{K}_J$ *is the full subcomplex of* $\mathcal{K}$ *obtained by restricting to* $J \subset [m]$. *We assume* $\widetilde{H}^{-1}(\mathcal{K}_\varnothing;\mathbf{k}) = \mathbf{k}$ *above.*

We shall give a proof of Hochster's formula following [**179**]. The idea is to first reduce the Koszul algebra $[\Lambda[u_1,\ldots,u_m] \otimes \mathbf{k}[\mathcal{K}], d]$ to a certain finite dimensional quotient $R^*(\mathcal{K})$, without changing the cohomology, and then identify $R^*(\mathcal{K})$ with the sum of simplicial cochain complexes of all full subcomplexes in $\mathcal{K}$. The algebra $R^*(\mathcal{K})$ will be also used in the cohomological calculations for moment-angle complexes in Chapter 4.

We use simplified notation u_Jv_I for a monomial $u_J \otimes v_I$ in the Koszul algebra $\Lambda[u_1,\ldots,u_m] \otimes \mathbf{k}[\mathcal{K}]$.

CONSTRUCTION 3.2.5. We introduce the quotient algebra

$$R^*(\mathcal{K}) = \Lambda[u_1, \dots, u_m] \otimes \mathbf{k}[\mathcal{K}]/(v_i^2 = u_i v_i = 0,\ 1 \leqslant i \leqslant m).$$

Since the ideal generated by v_i^2 and $u_i v_i$ is homogeneous and invariant with respect to the differential (since $d(u_i v_i) = 0$ and $d(v_i^2) = 0$), we obtain that $R^*(\mathcal{K})$ has differential and bigrading (3.1). We also have the quotient projection

$$\varrho\colon \Lambda[u_1, \dots, u_m] \otimes \mathbf{k}[\mathcal{K}] \to R^*(\mathcal{K}).$$

By definition, the algebra $R^*(\mathcal{K})$ has a $\mathbf{k}$-vector space basis consisting of monomials $u_J v_I$ where $J \subset [m]$, $I \in \mathcal{K}$ and $J \cap I = \varnothing$. Therefore,

$$\dim_{\mathbf{k}} R^{-p,2q} = f_{q-p-1}\tbinom{m-q+p}{p}, \tag{3.3}$$

where $(f_0, f_1, \dots, f_{n-1})$ is the f-vector of $\mathcal{K}$ and $f_{-1} = 1$. We have a $\mathbf{k}$-linear map

$$\iota\colon R^*(\mathcal{K}) \to \Lambda[u_1, \dots, u_m] \otimes \mathbf{k}[\mathcal{K}]$$

sending each $u_J v_I$ identically. The map ι commutes with the differentials, and therefore defines a homomorphism of bigraded differential $\mathbf{k}$-vector spaces satisfying the relation $\varrho \cdot \iota = \mathrm{id}$. Note that ι is not a map of algebras.

LEMMA 3.2.6. *The projection homomorphism* $\varrho\colon \Lambda[u_1, \dots, u_m] \otimes \mathbf{k}[\mathcal{K}] \to R^*(\mathcal{K})$ *induces an isomorphism in cohomology.*

PROOF. The argument is similar to that used for the Koszul resolution (see Construction A.2.4). We shall construct a cochain homotopy between the maps id and $\iota \cdot \varrho$ from $\Lambda[u_1, \dots, u_m] \otimes \mathbf{k}[\mathcal{K}]$ to itself, that is, a map s satisfying the identity

$$ds + sd = \mathrm{id} - \iota \cdot \varrho. \tag{3.4}$$

We first consider the case $\mathcal{K} = \Delta^{m-1}$. Then $\Lambda[u_1, \dots, u_m] \otimes \mathbf{k}[\Delta^{m-1}]$ is the Koszul resolution (A.5), which will be denoted by

$$E = E_m = \Lambda[u_1, \dots, u_m] \otimes \mathbf{k}[v_1, \dots, v_m], \tag{3.5}$$

and the algebra $R^*(\Delta^{m-1})$ is isomorphic to

$$\bigl(\Lambda[u] \otimes \mathbf{k}[v]/(v^2, uv)\bigr)^{\otimes m}. \tag{3.6}$$

For $m = 1$, we define the map $s_1\colon E_1^{0,*} = \mathbf{k}[v] \to E_1^{-1,*}$ by the formula

$$s_1(a_0 + a_1 v + \dots + a_j v^j) = u(a_2 v + a_3 v^2 + \dots + a_j v^{j-1}).$$

We need to check identity (3.4) for $x = a_0 + a_1 v + \dots + a_j v^j \in E_1^{0,*}$ and for $ux \in E_1^{-1,*}$, as each element of E_1 is the sum of elements of these two types. In the first case we have $ds_1 x = x - a_0 - a_1 v = x - \iota\varrho x$, and $s_1 dx = 0$. In the second case, i.e. for $ux \in E_1^{-1,*}$, we have $ds_1(ux) = 0$, and $s_1 d(ux) = ux - a_0 u = ux - \iota\varrho(ux)$. In both cases (3.4) holds.

Now we may assume by induction that a cochain homotopy $s_m\colon E_m \to E_m$ has been already constructed for $m = k - 1$. Since $E_k = E_{k-1} \otimes E_1$, $\varrho_k = \varrho_{k-1} \otimes \varrho_1$ and $\iota_k = \iota_{k-1} \otimes \iota_1$, a direct calculation shows that the map

$$s_k = s_{k-1} \otimes \mathrm{id} + \iota_{k-1}\varrho_{k-1} \otimes s_1 \tag{3.7}$$

is a cochain homotopy between id and $\iota_k\varrho_k$, which finishes the proof for $\mathcal{K} = \Delta^{m-1}$.

In the case of arbitrary $\mathcal{K}$ the algebras $\Lambda[u_1, \dots, u_m] \otimes \mathbf{k}[\mathcal{K}]$ and $R^*(\mathcal{K})$ are obtained by factorising (3.5) and (3.6) respectively by the ideal $\mathcal{I}_{\mathcal{K}}$ in $\Lambda[u_1, \dots, u_m] \otimes \mathbf{k}[m]$. Observe that $\mathcal{I}_{\mathcal{K}}$ is generated by v_I with $I \notin \mathcal{K}$ as an ideal, and it has a $\mathbf{k}$-vector space basis of monomials $u_J v_{i_1}^{\alpha_1}\cdots v_{i_k}^{\alpha_k}$ with $I=\{i_1,\ldots,i_k\}\notin\mathcal{K}$ and $\alpha_i>0$. We need to check that

$$d(\mathcal{I}_\mathcal{K})\subset\mathcal{I}_\mathcal{K},\quad \iota\varrho(\mathcal{I}_\mathcal{K})\subset\mathcal{I}_\mathcal{K},\quad s(\mathcal{I}_\mathcal{K})\subset\mathcal{I}_\mathcal{K}.$$

The first inclusion is obvious: since d is a derivation, we only need to check that $dv_I\in\mathcal{I}_\mathcal{K}$ for $I\notin\mathcal{K}$, but $dv_I=0$. The second inclusion is also clear, since

$$\iota\varrho(u_J v_{i_1}^{\alpha_1}\cdots v_{i_k}^{\alpha_k})=\begin{cases}u_J v_{i_1}\cdots v_{i_k}, & \text{if } \alpha_i=1 \text{ and } J\cap\{i_1,\ldots,i_k\}=\varnothing;\\ 0, & \text{otherwise.}\end{cases}$$

It remains to check the third inclusion. By expanding the inductive formula (3.7) we obtain

$$s_m=s_1\otimes\mathrm{id}\otimes\cdots\otimes\mathrm{id}+\iota_1\varrho_1\otimes s_1\otimes\mathrm{id}\otimes\cdots\otimes\mathrm{id}+\iota_1\varrho_1\otimes\cdots\otimes\iota_1\varrho_1\otimes s_1.$$

It follows that

$$s_m(u_J v_{i_1}^{\alpha_1}\cdots v_{i_k}^{\alpha_k})=\sum_{p\colon\,\alpha_p>1}\pm u_J u_{i_p} v_{i_1}^{\alpha_1}\cdots v_{i_p}^{\alpha_p-1}\cdots v_{i_k}^{\alpha_k}.$$

Therefore, $s(\mathcal{I}_\mathcal{K})\subset\mathcal{I}_\mathcal{K}$, and identity (3.4) holds in $\Lambda[u_1,\ldots,u_m]\otimes\mathbf{k}[\mathcal{K}]$. □

As an immediate consequence of Lemma 3.2.6 we obtain

COROLLARY 3.2.7. *We have that* $\beta^{-i,2j}\big(\mathbf{k}[\mathcal{K}]\big)=0$ *if* $i>m$ *or* $j>m$.

PROOF. Indeed, $R^{-i,2j}(\mathcal{K})=0$ if either i or j is greater than m. □

Now, in order to prove Theorem 3.2.4, we need to show that the cohomology of $R^*(\mathcal{K})$ is isomorphic to the direct sum of the reduced cohomology of the full subcomplexes on the right hand side of Hochster's formula. We shall see that this is true even without passing to cohomology, i.e. $R^*(\mathcal{K})$ is isomorphic to $\bigoplus_{I\subset m}C^*(\mathcal{K}_I)$, with the appropriate shift in dimensions, where C^* denotes the simplicial cochain groups. To do this, it is convenient to refine the grading in $\mathbf{k}[\mathcal{K}]$ as follows.

CONSTRUCTION 3.2.8 (multigraded structure in face rings and Tor-algebras). A *multigrading* (more precisely, an $\mathbb{N}^m$-grading) is defined in $\mathbf{k}[v_1,\ldots,v_m]$ by setting

$$\operatorname{mdeg} v_1^{i_1}\cdots v_m^{i_m}=(2i_1,\ldots,2i_m).$$

Since $\mathbf{k}[\mathcal{K}]$ is the quotient of the polynomial ring by a monomial ideal, it inherits the multigrading. We may assume that all free modules in the resolution (A.2) are multigraded and the differentials preserve the multidegree. Then the algebra $\operatorname{Tor}_{\mathbf{k}[m]}(\mathbf{k}[\mathcal{K}],\mathbf{k})$ acquires the canonical $\mathbb{Z}\oplus\mathbb{N}^m$-grading, i.e.

$$\operatorname{Tor}_{\mathbf{k}[v_1,\ldots,v_m]}\big(\mathbf{k}[\mathcal{K}],\mathbf{k}\big)=\bigoplus_{i\geqslant 0,\ \boldsymbol{a}\in\mathbb{N}^m}\operatorname{Tor}^{-i,2\boldsymbol{a}}_{\mathbf{k}[v_1,\ldots,v_m]}\big(\mathbf{k}[\mathcal{K}],\mathbf{k}\big).$$

The differential algebra $R^*(\mathcal{K})$ also acquires a $\mathbb{Z}\oplus\mathbb{N}^m$-grading, and Lemma 3.2.6 implies that

$$\operatorname{Tor}^{-i,2\boldsymbol{a}}_{\mathbf{k}[v_1,\ldots,v_m]}\big(\mathbf{k}[\mathcal{K}],\mathbf{k}\big)\cong H^{-i,2\boldsymbol{a}}[R^*(\mathcal{K}),d]. \tag{3.8}$$

We may view a subset $J\subset[m]$ as a $(0,1)$-vector in $\mathbb{N}^m$ whose jth coordinate is 1 if $j\in J$ and is 0 otherwise. Then there is the following multigraded version of Hochster's formula:

THEOREM 3.2.9. *For any subset $J \subset [m]$ we have*

$$\operatorname{Tor}^{-i,2J}_{\mathbf{k}[v_1,\ldots,v_m]}\bigl(\mathbf{k}[\mathcal{K}],\mathbf{k}\bigr) \cong \widetilde{H}^{|J|-i-1}(\mathcal{K}_J),$$

and $\operatorname{Tor}^{-i,2\boldsymbol{a}}_{\mathbf{k}[m]}(\mathbf{k}[\mathcal{K}],\mathbf{k}) = 0$ *if $\boldsymbol{a}$ is not a $(0,1)$-vector.*

PROOF OF THEOREM 3.2.4 AND THEOREM 3.2.9. Let $C^q(\mathcal{K}_J)$ denote the qth simplicial cochain group with coefficients in $\mathbf{k}$. Denote by $\alpha_L \in C^{p-1}(\mathcal{K}_J)$ the basis cochain corresponding to an oriented simplex $L = (l_1,\ldots,l_p) \in \mathcal{K}_J$; it takes value 1 on L and vanishes on all other simplices. Now we define a $\mathbf{k}$-linear map

$$\begin{aligned} f\colon C^{p-1}(\mathcal{K}_J) &\longrightarrow R^{p-|J|,2J}(\mathcal{K}), \\ \alpha_L &\longmapsto \varepsilon(L,J)\, u_{J\setminus L} v_L, \end{aligned} \tag{3.9}$$

where $\varepsilon(L,J)$ is the sign defined by

$$\varepsilon(L,J) = \prod_{j\in L} \varepsilon(j,J),$$

and $\varepsilon(j,J) = (-1)^{r-1}$ if j is the rth element of the set $J \subset [m]$, written in increasing order. Obviously, f is an isomorphism of $\mathbf{k}$-vector spaces, and a direct check shows that it commutes with the differentials. Indeed, we have

$$\begin{aligned} f(d\alpha_L) &= f\Bigl(\sum_{j\in J\setminus L,\, j\cup L\in\mathcal{K}_J} \varepsilon(j, j\cup L)\,\alpha_{j\cup L}\Bigr) \\ &= \sum_{j\in J\setminus L} \varepsilon(j\cup L, J)\varepsilon(j, j\cup L)\, u_{J\setminus(j\cup L)} v_{j\cup L} \end{aligned}$$

(note that $v_{j\cup L} \in \mathbf{k}[\mathcal{K}]$, and hence it is zero unless $j\cup L \in \mathcal{K}_J$). On the other hand,

$$df(\alpha_L) = \sum_{j\in J\setminus L} \varepsilon(L,J)\varepsilon(j, J\setminus L) u_{J\setminus(j\cup L)} v_{j\cup L}.$$

By the definition of $\varepsilon(L,J)$,

$$\varepsilon(j\cup L,J)\varepsilon(j,j\cup L) = \varepsilon(L,J)\varepsilon(j,J)\varepsilon(j,j\cup L) = \varepsilon(L,J)\varepsilon(j,J\setminus L),$$

which implies that $f(d\alpha_L) = df(\alpha_L)$. Therefore, f together with the map $\mathbf{k} \to R^{-|J|,2J}(\mathcal{K})$, $1 \mapsto u_J$, defines an isomorphism of cochain complexes

$$\begin{tikzcd}
0 \arrow[r] & \mathbf{k} \arrow[r, "d"] \arrow[d, "\cong"] & C^0(\mathcal{K}_J) \arrow[r, "d"] \arrow[d, "\cong"] & \cdots \arrow[r, "d"] & C^{p-1}(\mathcal{K}_J) \arrow[r, "d"] \arrow[d, "\cong"] & \cdots \\
0 \arrow[r] & R^{-|J|,2J}(\mathcal{K}) \arrow[r, "d"] & R^{1-|J|,2J}(\mathcal{K}) \arrow[r, "d"] & \cdots \arrow[r, "d"] & R^{p-|J|,2J}(\mathcal{K}) \arrow[r, "d"] & \cdots
\end{tikzcd}$$

Then it follows from (3.8) that

$$\widetilde{H}^{p-1}(K_J) \cong \operatorname{Tor}^{p-|J|,2J}_{\mathbf{k}[v_1,\ldots,v_m]}\bigl(\mathbf{k}[\mathcal{K}],\mathbf{k}\bigr),$$

which is equivalent to the first isomorphism of Theorem 3.2.9. Since $R^{-i,2\boldsymbol{a}}(\mathcal{K}) = 0$ if $\boldsymbol{a}$ is not a $(0,1)$-vector, $\operatorname{Tor}^{-i,2\boldsymbol{a}}_{\mathbf{k}[m]}(\mathbf{k}[\mathcal{K}],\mathbf{k})$ vanishes for such $\boldsymbol{a}$. □

Since $\operatorname{Tor}_{\mathbf{k}[m]}(\mathbf{k}[\mathcal{K}],\mathbf{k})$ is an algebra, the isomorphisms of Theorem 3.2.4 turn the direct sum

$$\bigoplus_{\substack{p\geqslant 0\\ J\subset[m]}} \widetilde{H}^{p-1}(\mathcal{K}_J) \tag{3.10}$$

into a (multigraded) **k**-algebra. Consider the product in the simplicial cochains of full subcomplexes given by

$$(3.11)\qquad \begin{aligned} \mu\colon C^{p-1}(\mathcal{K}_I)\otimes C^{q-1}(\mathcal{K}_J) &\longrightarrow C^{p+q-1}(\mathcal{K}_{I\sqcup J}),\\ \alpha_L\otimes\alpha_M \quad &\longmapsto \begin{cases} \alpha_{L\sqcup M}, & \text{if } I\cap J=\varnothing;\\ 0, & \text{otherwise.}\end{cases}\end{aligned}$$

Here $\alpha_{L\sqcup M}\in C^{p+q-1}(\mathcal{K}_{I\sqcup J})$ denotes the basis simplicial cochain corresponding to $L\sqcup M$ if the latter is a simplex of $\mathcal{K}_{I\sqcup J}$ and zero otherwise. If $I\cap J=\varnothing$, then $\mathcal{K}_{I\sqcup J}$ is a subcomplex in the join $\mathcal{K}_I*\mathcal{K}_J$, and the above product is the restriction to $\mathcal{K}_{I\sqcup J}$ of the standard exterior product

$$C^{p-1}(\mathcal{K}_I)\otimes C^{q-1}(\mathcal{K}_J)\longrightarrow C^{p+q-1}(\mathcal{K}_I*\mathcal{K}_J).$$

PROPOSITION 3.2.10. *The product in the direct sum* $\bigoplus_{p\geqslant 0,\ J\subset[m]}\widetilde{H}^{p-1}(\mathcal{K}_J)$ *induced by the isomorphisms from Hochster's theorem coincides up to a sign with the product given by* (3.11).

PROOF. This is a direct calculation. We use the isomorphism f given by (3.9):

$$\alpha_L\cdot\alpha_M=f^{-1}\bigl(f(\alpha_L)\cdot f(\alpha_M)\bigr)=f^{-1}\bigl(\varepsilon(L,I)\,u_{I\setminus L}v_L\,\varepsilon(M,J)\,u_{J\setminus M}v_M\bigr)$$

If $I\cap J\neq\varnothing$, then the product $u_{I\setminus L}v_Lu_{J\setminus M}v_M$ is zero in $R^*(\mathcal{K})$. Otherwise we have that $u_{I\setminus L}v_Lu_{J\setminus M}v_M=\zeta\,u_{(I\cup J)\setminus(L\cup M)}v_{L\cup M}$, where $\zeta=\prod_{k\in I\setminus L}\varepsilon(k,k\cup J\setminus M)$, and we can continue the above identity as

$$\alpha_L\cdot\alpha_M=\varepsilon(L,I)\,\varepsilon(M,J)\,\zeta\,\varepsilon(L\cup M,I\cup J)\,\alpha_{L\sqcup M}.$$

Note that this calculation also gives the explicit value for the correcting sign, but we shall not need this. □

Let P be a simple polytope. The multigraded components of $\mathrm{Tor}_{\mathbf{k}[m]}(\mathbf{k}[P],\mathbf{k})$ can be expressed directly in terms of P as follows. Let $\{F_1,\dots,F_m\}$ be the set of facets of P. Given $I\subset[m]$, we define the following subset of the boundary of P:

$$P_I=\bigcup_{i\in I}F_i.$$

PROPOSITION 3.2.11. *For any subset* $J\subset[m]$ *we have*

$$\mathrm{Tor}^{-i,2J}_{\mathbf{k}[v_1,\dots,v_m]}\bigl(\mathbf{k}[P],\mathbf{k}\bigr)\cong\widetilde{H}^{|J|-i-1}(P_J),$$

and $\mathrm{Tor}^{-i,2\boldsymbol{a}}_{\mathbf{k}[m]}(\mathbf{k}[P],\mathbf{k})=0$ *if* $\boldsymbol{a}$ *is not a* $(0,1)$*-vector.*

PROOF. Let $\mathcal{K}=\mathcal{K}_P$ be the nerve complex of P. Then the statement follows from Theorem 3.2.9 and the fact that $\mathcal{K}_J$ is a deformation retract of P_J. The latter is because P is simple, and therefore, $P_J=\bigcup_{i\in J}F_J=\bigcup_{i\in J}\mathrm{st}_{\mathcal{K}'}\{i\}$ (by Proposition 2.3.8), which is the combinatorial neighbourhood of $(\mathcal{K}_J)'$ in $\mathcal{K}'$. □

For a description of the multiplication in $\mathrm{Tor}_{\mathbf{k}[m]}(\mathbf{k}[P],\mathbf{k})$ in terms of P, see Exercise 3.2.14.

EXAMPLE 3.2.12.

1. Let P be a 4-gon, so that $\mathcal{K}_P =$ 

, as in Example 3.2.3. This time we calculate the Betti numbers $\beta^{-i,2j}(\mathbf{k}[P])$ using Hochster's formula. We have that

$$\begin{aligned} \beta^{0,0}\big(\mathbf{k}[P]\big) &= \dim \widetilde{H}^{-1}(\varnothing) = 1 && 1,\\ \beta^{-1,4}\big(\mathbf{k}[P]\big) &= \dim \widetilde{H}^{0}\big(P_{\{1,3\}}\big) \oplus \widetilde{H}^{0}\big(P_{\{2,4\}}\big) = 2 && u_1v_3, u_2v_4\\ \beta^{-2,8}\big(\mathbf{k}[P]\big) &= \dim \widetilde{H}^{1}\big(P_{\{1,2,3,4\}}\big) = 1 && u_1u_2v_3v_4, \end{aligned}$$

where in the right column we include cocycles in the Koszul algebra $\Lambda[u_1,\ldots,u_m]\otimes\mathbf{k}[P]$ representing generators of the corresponding cohomology groups. All other Betti numbers are zero. We have a nontrivial product $[u_1v_3]\cdot[u_2v_4] = [u_1u_2v_3v_4]$; all other products of positive-dimensional classes are zero. Note that in this example all Tor-groups have bases represented by monomials in the Koszul algebra. This is not the case in general, as is shown by the next example.

2. Now let $\mathcal{K} =$ 

be the union of two segments. Then the generator of

$$\mathrm{Tor}^{-3,8}_{\mathbf{k}[v_1,\ldots,v_4]}\big(\mathbf{k}[\mathcal{K}],\mathbf{k}\big) \cong \widetilde{H}^{0}\big(\mathcal{K}_{\{1,2,3,4\}};\mathbf{k}\big) \cong \mathbf{k}$$

is represented by the cocycle $u_1u_2u_3v_4 - u_1u_2u_4v_3$ in the Koszul algebra, and it cannot be represented by a monomial.

3. Let us calculate the Betti numbers (both bigraded and multigraded) of $\mathbf{k}[\mathcal{K}]$ for the complex shown in Fig. 3.1, using Hochster's formula. We have

$$\begin{aligned} \beta^{0,0} &= \dim \widetilde{H}^{0}(\varnothing) = 1,\\ \beta^{-1,4} &= \beta^{-1,(2,0,0,0,2)} + \beta^{-1,(0,0,2,2,0)} = \dim \widetilde{H}^{0}\big(\mathcal{K}_{\{1,5\}}\big) \oplus \widetilde{H}^{0}\big(\mathcal{K}_{\{3,4\}}\big) = 2,\\ \beta^{-1,6} &= \beta^{-1,(2,2,2,0,0)} + \beta^{-1,(0,2,0,2,2)} = \dim \widetilde{H}^{1}\big(\mathcal{K}_{\{1,2,3\}}\big) \oplus \widetilde{H}^{1}\big(\mathcal{K}_{\{2,4,5\}}\big) = 2,\\ \beta^{-2,8} &= \beta^{-2,(0,2,2,2,2)} + \beta^{-2,(2,0,2,2,2)} + \cdots + \beta^{-2,(2,2,2,2,0)}\\ &\qquad\qquad = \dim \widetilde{H}^{1}\big(\mathcal{K}_{\{2,3,4,5\}}\big) \oplus \cdots \oplus \widetilde{H}^{1}\big(\mathcal{K}_{\{1,2,3,4\}}\big) = 5,\\ \beta^{-3,10} &= \beta^{-3,(2,2,2,2,2)} = \dim \widetilde{H}^{1}\big(\mathcal{K}_{\{1,2,3,4,5\}}\big) = 2. \end{aligned}$$

All other Betti numbers are zero.

4. Let $\mathcal{K}$ be a triangulation of the real projective plane $\mathbb{R}P^2$ with m vertices (the minimal example has $m = 6$, see Fig. 3.2, where the vertices with the same labels are identified, and the boundary edges are identified according to the orientation shown). Then, by Hochster's formula,



FIGURE 3.2. 6-vertex triangulation of $\mathbb{R}P^2$.

$$\operatorname{Tor}^{3-m,2m}_{\mathbf{k}[v_1,\ldots,v_m]}\bigl(\mathbf{k}[\mathcal{K}],\mathbf{k}\bigr) = \widetilde{H}^2(\mathcal{K}_{[m]};\mathbf{k}) = \widetilde{H}^2(\mathbb{R}P^2;\mathbf{k}) = 0$$

if the characteristic of $\mathbf{k}$ is not 2. On the other hand,

$$\operatorname{Tor}^{3-m,2m}_{\mathbb{Z}_2[v_1,\ldots,v_m]}\bigl(\mathbb{Z}_2[\mathcal{K}],\mathbb{Z}_2\bigr) = \widetilde{H}^2(\mathcal{K}_{[m]};\mathbb{Z}_2) = \widetilde{H}^2(\mathbb{R}P^2;\mathbb{Z}_2) = \mathbb{Z}_2.$$

This example shows that the Tor-groups of $\mathbf{k}[\mathcal{K}]$, and even the algebraic Betti numbers, depend on $\mathbf{k}$. A similar example shows that $\operatorname{Tor}_{\mathbb{Z}[m]}(\mathbb{Z}[\mathcal{K}],\mathbb{Z})$ may have an arbitrary amount of additive torsion. (This is a well-known fact for the usual cohomology of spaces, and so we may take $\mathcal{K}$ to be a triangulation of a space with the appropriate torsion in cohomology.)

Exercises.

3.2.13. Let P be a pentagon. Calculate the bigraded Betti numbers of $\mathbf{k}[P]$ and the multiplication in $\operatorname{Tor}_{\mathbf{k}[m]}(\mathbf{k}[P],\mathbf{k})$

(a) using algebra $R^*(\mathcal{K}_P)$ and Lemma 3.2.6;
(b) using Hochster's theorem and Proposition 3.2.10,

and compare the results.

3.2.14. Use Proposition 3.2.11 and the isomorphism

$$\widetilde{H}^{|J|-i-1}(P_J) \cong H^{|J|-i}(P,P_J)$$

to show that the multiplication induced from $\operatorname{Tor}_{\mathbf{k}[m]}(\mathbf{k}[P],\mathbf{k})$ in the direct sum

$$\bigoplus_{\substack{p\geqslant 0\\ J\subset[m]}} H^p(P,P_J)$$

comes from the standard exterior multiplication

$$H^p(P,P_I)\otimes H^q(P,P_J) \longrightarrow H^{p+q}(P,P_I\cup P_J)$$

when $I\cap J=\varnothing$ and is zero otherwise.

3.2.15. Complete the details in the following algebraic proof of the Alexander duality (Theorem 2.4.5); this argument ascends to the original work of Hochster [**124**]:

1. Choose $J\notin\mathcal{K}$, that is, $\widehat{J}=[m]\setminus J\in\widehat{\mathcal{K}}$, and show that for any $L=\{l_1,\ldots,l_q\}\subset J$,

$$J\setminus L\notin\mathcal{K} \iff L\in\operatorname{lk}_{\widehat{\mathcal{K}}}\widehat{J}.$$

2. Consider the Koszul algebra

$$S(\mathcal{K}) = \bigl[\Lambda[u_1,\ldots,u_m]\otimes\mathcal{I}_{\mathcal{K}},d\bigr]$$

of the Stanley–Reisner ideal $\mathcal{I}_{\mathcal{K}}$ (see Lemma A.2.10 and the remark after it), and show that its multigraded component $S^{-q,2J}(\mathcal{K})$ has a $\mathbf{k}$-basis consisting of monomials $u_Lv_{J\setminus L}$ where $L\in\operatorname{lk}_{\widehat{\mathcal{K}}}\widehat{J}$.

3. Consider the $\mathbf{k}$-vector space isomorphism

$$\begin{aligned} g\colon C_{q-1}(\operatorname{lk}_{\widehat{\mathcal{K}}}\widehat{J}) &\longrightarrow S^{-q,2J}(\mathcal{K}),\\ [L] &\longmapsto u_Lv_{J\setminus L}, \end{aligned}$$

where $[L]\in C_{q-1}(\operatorname{lk}_{\widehat{\mathcal{K}}}\widehat{J})$ is the basis simplicial chain corresponding to L. Show that g commutes with the differentials, and therefore defines an isomorphism of chain complexes (in analogy with (3.9), but with no correction sign).

4. Deduce that

$$\widetilde{H}_{q-1}(\mathrm{lk}_{\widehat{\mathcal{K}}}\,\widehat{J}) \cong \mathrm{Tor}^{-q,2J}_{\mathbf{k}[m]}(\mathcal{I}_{\mathcal{K}},\mathbf{k}) \cong \mathrm{Tor}^{-q-1,2J}_{\mathbf{k}[m]}(\mathbf{k}[\mathcal{K}],\mathbf{k}) \cong \widetilde{H}^{|J|-q-2}(\mathcal{K}_J),$$

where the first isomorphism is obtained by passing to homology in step 3, the second follows from the long exact sequence of Theorem A.2.5 (e), and the third is Theorem 3.2.4. It remains to note that the resulting isomorphism is equivalent to that of Corollary 2.4.6.

3.3. Cohen–Macaulay complexes

It is usually quite difficult to determine whether a given ring is Cohen–Macaulay (see Appendix, Section A.3). One of the key results of combinatorial commutative algebra, the *Reisner Theorem*, gives an effective criterion for the Cohen–Macaulayness of face rings, in terms of simplicial cohomology of $\mathcal{K}$. A reformulation of Reisner's criterion, due to Munkres and Stanley, tells us that the Cohen–Macaulayness of $\mathbf{k}[\mathcal{K}]$ is a topological property of $\mathcal{K}$, i.e. it depends only on the topology of the realisation $|\mathcal{K}|$. These results have many important applications in both combinatorial commutative algebra and toric topology.

Here we assume that $\mathbf{k}$ is a field, unless otherwise stated. If $\mathcal{K}$ is of dimension $n-1$, then the Krull dimension of $\mathbf{k}[\mathcal{K}]$ is n (an exercise). We start with the following combinatorial description of homogeneous systems of parameters (hsop's) in $\mathbf{k}[\mathcal{K}]$ in terms of the restriction homomorphisms $s_I\colon \mathbf{k}[\mathcal{K}] \to \mathbf{k}[v_i\colon i\in I]$ (see Proposition 3.1.8).

Lemma 3.3.1. *Let $\mathcal{K}$ be a simplicial complex of dimension $n-1$. A sequence of homogeneous elements $\boldsymbol{t}=(t_1,\ldots,t_n)$ of $\mathbf{k}[\mathcal{K}]$ is a homogeneous system of parameters if and only if*

$$\dim_{\mathbf{k}}\bigl(\mathbf{k}[v_i\colon i\in I]/s_I(\boldsymbol{t})\bigr) < \infty$$

for each simplex $I\in\mathcal{K}$, where $s_I(\boldsymbol{t})$ is the image of the sequence $\boldsymbol{t}$ under the restriction map s_I.

Proof. Assume that $\boldsymbol{t}$ is an hsop. By applying the right exact functor $\otimes_{\mathbf{k}[\boldsymbol{t}]}$ to the epimorphism $s_I\colon \mathbf{k}[\mathcal{K}] \to \mathbf{k}[v_i\colon i\in I]$ we obtain that $\mathbf{k}[\mathcal{K}]/\boldsymbol{t} \to \mathbf{k}[v_i\colon i\in I]/s_I(\boldsymbol{t})$ is also an epimorphism. Hence,

$$\dim_{\mathbf{k}}\bigl(\mathbf{k}[v_i\colon i\in I]/s_I(\boldsymbol{t})\bigr) \leqslant \dim_{\mathbf{k}}\bigl(\mathbf{k}[\mathcal{K}]/\boldsymbol{t}\bigr) < \infty.$$

For the opposite statement, assume that

$$\dim_{\mathbf{k}} \bigoplus_{I\in\mathcal{K}} \mathbf{k}[v_i\colon i\in I]/s_I(\boldsymbol{t}) < \infty.$$

Consider the short exact sequence of $\mathbf{k}[\boldsymbol{t}]$-modules

$$0 \longrightarrow \mathbf{k}[\mathcal{K}] \xrightarrow{\oplus s_I} \bigoplus_{I\in\mathcal{K}} \mathbf{k}[v_i\colon i\in I] \longrightarrow Q \longrightarrow 0,$$

where Q is the quotient module, and the following fragment of the corresponding long exact sequence for Tor (Theorem A.2.5 (e)):

$$\cdots \longrightarrow \mathrm{Tor}^{-1}_{\mathbf{k}[\boldsymbol{t}]}(Q,\mathbf{k}) \longrightarrow \mathbf{k}[\mathcal{K}]/\boldsymbol{t} \longrightarrow \bigoplus_{I\in\mathcal{K}} \mathbf{k}[v_i\colon i\in I]/s_I(\boldsymbol{t}) \longrightarrow \cdots.$$

Since $\bigoplus_{I\in\mathcal{K}}\mathbf{k}[v_i\colon i\in I]$ is a finitely generated $\mathbf{k}[\boldsymbol{t}]$-module, its quotient Q is also finitely generated. Hence $\dim_{\mathbf{k}}\operatorname{Tor}^{-1}_{\mathbf{k}[\boldsymbol{t}]}(Q,\mathbf{k})<\infty$ (see Proposition A.2.6), and, by the exact sequence above, $\dim_{\mathbf{k}}\bigl(\mathbf{k}[\mathcal{K}]/\boldsymbol{t}\bigr)<\infty$. Therefore, $\boldsymbol{t}$ is an hsop in $\mathbf{k}[\mathcal{K}]$. $\square$

Recall that we refer to a sequence $\boldsymbol{t}=(t_1,\dots,t_n)\in\mathbf{k}[\mathcal{K}]$ as linear if $\deg t_i=2$ for all i. We may write a linear sequence as

$$t_i=\lambda_{i1}v_1+\cdots+\lambda_{im}v_m,\quad\text{for }1\leqslant i\leqslant n. \tag{3.12}$$

Lemma 3.3.2. *A linear sequence* $\boldsymbol{t}=(t_1,\dots,t_n)\subset\mathbf{k}[\mathcal{K}]$, $\dim\mathcal{K}=n-1$, *is an lsop if and only if the restriction* $s_I(\boldsymbol{t})$ *to each simplex of* $I\in\mathcal{K}$ *generates the polynomial algebra* $\mathbf{k}[v_i\colon i\in I]$ *(i.e. the rank of the* $n\times|I|$ *matrix* $\Lambda_I=(\lambda_{ij})$, $1\leqslant i\leqslant n$, $j\in I$, *is equal to* $|I|$*).*

Proof. Indeed, if $\boldsymbol{t}$ is linear, then the conditions $\dim_{\mathbf{k}}\mathbf{k}[v_i\colon i\in I]/s_I(\boldsymbol{t})<\infty$ and $\mathbf{k}[v_i\colon i\in I]/s_I(\boldsymbol{t})\cong\mathbf{k}$ are equivalent. The latter means that $s_I(\boldsymbol{t})$ generates $\mathbf{k}[v_i\colon i\in I]$ as a $\mathbf{k}$-algebra. $\square$

Note that it is enough to verify the condition of Lemmata 3.3.1 and 3.3.2 only for maximal simplices $I\in\mathcal{K}$.

Definition 3.3.3. A linear sequence $\boldsymbol{t}=(t_1,\dots,t_n)\subset\mathbb{Z}[\mathcal{K}]$ is referred to as a *integral lsop* if its reduction modulo p is an lsop in $\mathbb{Z}_p[\mathcal{K}]$ for any prime p. Equivalently, $\boldsymbol{t}$ is an integral lsop if $n=\dim\mathcal{K}+1$ and the restriction $s_I(\boldsymbol{t})$ to each simplex $I\in\mathcal{K}$ generates the polynomial ring $\mathbb{Z}[v_i\colon i\in I]$ (the equivalence of these two conditions is an exercise).

Although the rational face ring $\mathbb{Q}[\mathcal{K}]$ always admit an lsop by Theorem A.3.10, an lsop in $\mathbb{Z}_p[\mathcal{K}]$ for a prime p (or an integral lsop in $\mathbb{Z}[\mathcal{K}]$) may fail to exist, as is shown by the next example.

Example 3.3.4.

1. Let $\mathcal{K}$ be a simplicial complex of dimension $n-1$ on $m\geqslant 2^n$ vertices whose 1-skeleton is a complete graph. Then the face ring $\mathbb{Z}_2[\mathcal{K}]$ does not admit an lsop. Indeed, assume that (3.12) is an lsop. Then, by Corollary 3.3.2, each column vector of the $n\times m$-matrix (λ_{ij}) is nonzero, and all column vectors are pairwise different (since each pair of vertices of $\mathcal{K}$ spans an edge). This is a contradiction, since the number of different nonzero vectors in $\mathbb{Z}_2^n$ is 2^n-1. By considering the reduction modulo 2 we obtain that $\mathbb{Z}[\mathcal{K}]$ also does not admit an integral lsop.

2. There are also simple polytopes P whose face rings $\mathbf{k}[P]$ do not admit an lsop over $\mathbb{Z}_2$ or $\mathbb{Z}$. Indeed, let P be the dual of a 2-neighbourly simplicial n-polytope (e.g., a cyclic polytope of dimension $n\geqslant 4$, see Example 1.1.18) with $m\geqslant 2^n$ vertices. Then the 1-skeleton of $\mathcal{K}_P$ is a complete graph, and therefore $\mathbb{Z}[P]=\mathbb{Z}[\mathcal{K}_P]$ does not admit an lsop. This example is taken from **[70]**.

By considering the reduction modulo 2 we observe that the ring $\mathbb{Z}[\mathcal{K}]$ for $\mathcal{K}$ from the previous example also does not admit an integral lsop. Existence of integral lsop's in the face rings $\mathbb{Z}[\mathcal{K}]$ is a very subtle question of great importance for toric topology; it will be discussed in more detail in Section 4.7.

Definition 3.3.5. $\mathcal{K}$ is a *Cohen–Macaulay complex over a field* $\mathbf{k}$ if $\mathbf{k}[\mathcal{K}]$ is a Cohen–Macaulay algebra. We say that $\mathcal{K}$ is a *Cohen–Macaulay complex over* $\mathbb{Z}$, or simply a *Cohen–Macaulay complex*, if $\mathbf{k}[\mathcal{K}]$ is a Cohen–Macaulay algebra for $\mathbf{k}=\mathbb{Q}$ and any finite field.

REMARK. We shall often consider $\mathbf{k}[\mathcal{K}]$ as a $\mathbf{k}[m]$-module rather than a $\mathbf{k}$-algebra. However, this does not affect regular sequences and the Cohen–Macaulay property: it is an easy exercise to show that a sequence $\boldsymbol{t} \subset \mathbf{k}[m]$ is $\mathbf{k}[m]$-regular for $\mathbf{k}[\mathcal{K}]$ as a $\mathbf{k}[m]$-module if and only the image of $\boldsymbol{t}$ in $\mathbf{k}[\mathcal{K}]$ is $\mathbf{k}[\mathcal{K}]$-regular. In particular, $\mathbf{k}[\mathcal{K}]$ is a Cohen–Macaulay algebra if and only if it is a Cohen–Macaulay $\mathbf{k}[m]$-module. We shall therefore not distinguish between these two notions.

EXAMPLE 3.3.6. Let $\mathcal{K} = \partial\Delta^2$. Then $\mathbf{k}[\mathcal{K}] = \mathbf{k}[v_1, v_2, v_3]/(v_1v_2v_3)$ and the Krull dimension is $\dim \mathbf{k}[\mathcal{K}] = 2$. The elements $v_1, v_2 \in \mathbf{k}[\mathcal{K}]$ are algebraically independent, but do not form an lsop, since $\mathbf{k}[\mathcal{K}]/(v_1, v_2) \cong \mathbf{k}[v_3]$ and $\dim\bigl(\mathbf{k}[\mathcal{K}]/(v_1, v_2)\bigr) = 1$. On the other hand, the elements $t_1 = v_1 - v_3$, $t_2 = v_2 - v_3$ form an lsop, since $\mathbf{k}[\mathcal{K}]/(t_1, t_2) \cong \mathbf{k}[t]/t^3$. It is easy to see that $\mathbf{k}[\mathcal{K}]$ is a free $\mathbf{k}[t_1, t_2]$-module on the basis $\{1, v_1, v_1^2\}$. Therefore, $\mathbf{k}[\mathcal{K}]$ is a Cohen–Macaulay ring and (t_1, t_2) is a regular sequence.

Cohen–Macaulay complexes can be characterised homologically as follows:

PROPOSITION 3.3.7. *The following conditions are equivalent for a simplicial complex $\mathcal{K}$ of dimension $n-1$ with m vertices:*

(a) *$\mathcal{K}$ is Cohen–Macaulay over a field $\mathbf{k}$;*
(b) *$\beta^{-i}(\mathbf{k}[\mathcal{K}]) = 0$ for $i > m - n$ and $\beta^{-(m-n)}(\mathbf{k}[\mathcal{K}]) \neq 0$.*

PROOF. By definition, condition (a) is that $\operatorname{depth} \mathbf{k}[\mathcal{K}] = n$. Condition (b) is equivalent to the equality $\operatorname{pdim} \mathbf{k}[\mathcal{K}] = m - n$, by Corollary A.2.7. Since $\operatorname{depth} \mathbf{k}[m] = m$, the two conditions are equivalent by Theorem A.3.8. □

EXAMPLE 3.3.8. Let $\mathcal{K}$ be the 6-vertex triangulation of $\mathbb{R}P^2$, see Example 3.2.12.4 and Figure 3.2. Then $m - n = 3$, and, by Theorem 3.2.4,

$$\beta^{-4}\bigl(\mathbb{Z}_2[\mathcal{K}]\bigr) = \dim_{\mathbb{Z}_2} \widetilde{H}^1(\mathbb{R}P^2; \mathbb{Z}_2) = 1,$$

so $\mathcal{K}$ is not Cohen–Macaulay over $\mathbb{Z}_2$. On the other hand, a similar calculation shows that if the characteristic of $\mathbf{k}$ is not 2, then $\beta^{-i}(\mathbf{k}[\mathcal{K}]) = 0$ for $i > 3$ and $\beta^{-3}(\mathbf{k}[\mathcal{K}]) = 6$, i.e. $\mathcal{K}$ is Cohen–Macaulay over such fields.

PROPOSITION 3.3.9 (Stanley). *If $\mathcal{K}$ is a Cohen–Macaulay complex of dimension $n-1$, then $\boldsymbol{h}(\mathcal{K}) = (h_0, \ldots, h_n)$ is an M-vector (see Definition 1.4.13).*

PROOF. Let $\mathcal{K}$ be Cohen–Macaulay, and let $\boldsymbol{t} = (t_1, \ldots, t_n)$ be an lsop in $\mathbf{k}[\mathcal{K}]$, where $\mathbf{k}$ is a field of zero characteristic. Then, by Proposition A.3.14,

$$F\bigl(\mathbf{k}[\mathcal{K}], \lambda\bigr) = \frac{F\bigl(\mathbf{k}[\mathcal{K}]/\boldsymbol{t}; \lambda\bigr)}{(1-\lambda^2)^n}.$$

On the other hand, the Poincaré series of $\mathbf{k}[\mathcal{K}]$ is given by Theorem 3.1.10, which implies that

$$F\bigl(\mathbf{k}[\mathcal{K}]/\boldsymbol{t}; \lambda\bigr) = h_0 + h_1\lambda^2 + \cdots + h_n\lambda^{2n}.$$

Now, $A = \mathbf{k}[\mathcal{K}]/\boldsymbol{t}$ is a graded algebra generated by its degree-two elements and $\dim_{\mathbf{k}} A^{2i} = h_i$, so $(h_0, \ldots, h_n)$ is an M-vector by definition. □

REMARK. According to a result of Stanley [**208**, Theorem II.3.3], if $(h_0, \ldots, h_n)$ is an M-vector, then there exists an $(n-1)$-dimensional Cohen–Macaulay complex $\mathcal{K}$ such that $h_i(\mathcal{K}) = h_i$. Together with Proposition 3.3.9, this gives a complete characterisation of face vectors of Cohen–Macaulay complexes.

The following fundamental result gives a combinatorial characterisation of Cohen–Macaulay complexes:

THEOREM 3.3.10 (Reisner [**196**]). *Let* $\mathbf{k} = \mathbb{Z}$ *or a field. A simplicial complex* $\mathcal{K}$ *is Cohen–Macaulay over* $\mathbf{k}$ *if and only if for any simplex* $I \in \mathcal{K}$ *(including* $I = \varnothing$*) and* $i < \dim(\operatorname{lk} I)$*, we have* $\widetilde{H}_i(\operatorname{lk} I; \mathbf{k}) = 0$.

The proof can be found in many texts on combinatorial commutative algebra, see e.g. [**208**, §II.4] or [**160**, Chapter 13]. It is not very hard, but uses the notion of *local cohomology*, which is beyond the scope of this book.

The following reformulation of Reisner's Theorem shows that the Cohen–Macaulayness is a topological property of a simplicial complex.

PROPOSITION 3.3.11 (Munkres, Stanley). *A simplicial complex* $\mathcal{K}$ *is Cohen–Macaulay over* $\mathbf{k}$ *if and only if for any point* $x \in |\mathcal{K}|$*, we have*

$$H_i(|\mathcal{K}|, |\mathcal{K}| \setminus x; \mathbf{k}) = \widetilde{H}_i(\mathcal{K}; \mathbf{k}) = 0 \quad \text{for } i < \dim \mathcal{K}.$$

PROOF. Let $I \in \mathcal{K}$. If $I = \varnothing$, then $\widetilde{H}_i(\mathcal{K}; \mathbf{k}) = \widetilde{H}_i(\operatorname{lk} I; \mathbf{k})$. If $I \neq \varnothing$, then

$$H_i(|\mathcal{K}|, |\mathcal{K}| \setminus x; \mathbf{k}) = \widetilde{H}_{i-|I|}(\operatorname{lk} I; \mathbf{k}) \tag{3.13}$$

for any x in the interior of I, by Proposition 2.2.14.

If $\mathcal{K}$ is Cohen–Macaulay, then it is pure (Exercise 3.3.20) and therefore $\operatorname{lk} I$ is pure of dimension $\dim \mathcal{K} - |I|$ (Exercise 2.2.19). Hence, $i < \dim \mathcal{K}$ implies that $i - |I| < \dim \operatorname{lk} I$ and $H_i(|\mathcal{K}|, |\mathcal{K}| \setminus x; \mathbf{k}) = 0$ by (3.13) and Theorem 3.3.10.

On the other hand, if $H_i(|\mathcal{K}|, |\mathcal{K}| \setminus x; \mathbf{k}) = 0$ for $i < \dim \mathcal{K}$, then $\widetilde{H}_j(\operatorname{lk} I; \mathbf{k}) = H_{j+|I|}(|\mathcal{K}|, |\mathcal{K}| \setminus x; \mathbf{k}) = 0$ for $j < \dim \operatorname{lk} I$, since $j + |I| < \dim \operatorname{lk} I + |I| \leqslant \dim \mathcal{K}$. Thus, $\mathcal{K}$ is Cohen–Macaulay by Theorem 3.3.10. ☐

COROLLARY 3.3.12. *If a triangulation of a space* X *is a Cohen–Macaulay complex, then any other triangulation of* X *is Cohen–Macaulay as well.*

COROLLARY 3.3.13. *Any triangulated sphere is a Cohen–Macaulay complex.*

In particular, the h-vector of a triangulated sphere is an M-vector. This fact was used by Stanley in his generalisation of the UBT (Theorem 1.4.4) to arbitrary sphere triangulations:

THEOREM 3.3.14 (UBT for spheres). *For any triangulated* $(n-1)$*-dimensional sphere* $\mathcal{K}$ *with* m *vertices, the* h*-vector* $(h_0, h_1, \ldots, h_n)$ *satisfies the inequalities*

$$h_i(\mathcal{K}) \leqslant \tbinom{m-n+i-1}{i}.$$

Therefore, the UBT holds for triangulated spheres, that is,

$$f_i(\mathcal{K}) \leqslant f_i\bigl(C^n(m)\bigr) \quad \text{for } i = 1, \ldots, n-1.$$

PROOF. Let $A = \mathbf{k}[\mathcal{K}]/\boldsymbol{t}$ be the algebra from the proof of Proposition 3.3.9, so that $\dim_{\mathbf{k}} A^{2i} = h_i$. In particular, $\dim_{\mathbf{k}} A^2 = h_1 = m - n$. Since A is generated by A^2, the number h_i cannot exceed the number of monomials of degree i in $m-n$ generators, i.e. $h_i \leqslant \binom{m-n+i-1}{i}$. The rest follows from Lemma 1.4.6. ☐

Exercises.

3.3.15. If $\mathcal{K}$ is a simplicial complex of dimension $n-1$, then $\dim \mathbf{k}[\mathcal{K}] = n$.

3.3.16. Let $\boldsymbol{t}$ be an lsop in $\mathbf{k}[\mathcal{K}]$. Then the $\mathbf{k}$-vector space $\mathbf{k}[\mathcal{K}]/\boldsymbol{t}$ is generated by monomials v_I for $I \in \mathcal{K}$. (Hint: prove that $v_i v_I = 0$ in $\mathbf{k}[\mathcal{K}]/\boldsymbol{t}$ for any $i \in [m]$ and for any *maximal* simplex $I \in \mathcal{K}$, and then use Proposition 3.1.9).

3.3.17. A sequence $\boldsymbol{t} \subset \mathbf{k}[m]$ is $\mathbf{k}[m]$-regular for $\mathbf{k}[\mathcal{K}]$ as a $\mathbf{k}[m]$-module if and only the image of $\boldsymbol{t}$ in $\mathbf{k}[\mathcal{K}]$ is $\mathbf{k}[\mathcal{K}]$-regular.

3.3.18. Let $\boldsymbol{t} = (t_1, \dots, t_n) \subset \mathbb{Z}[\mathcal{K}]$ be a linear sequence, $\dim \mathcal{K} = n-1$. The following conditions are equivalent:

(a) the reduction of $\boldsymbol{t}$ modulo p is an lsop in $\mathbb{Z}_p[\mathcal{K}]$ for any prime p;
(b) the restriction $s_I(\boldsymbol{t})$ to each simplex $I \in \mathcal{K}$ generates the polynomial ring $\mathbb{Z}[v_i \colon i \in I]$;
(c) for each $I \in \mathcal{K}$ the columns of the $n \times |I|$ matrix (λ_{ij}), $1 \leqslant i \leqslant n$, $j \in I$, generate the integer lattice $\mathbb{Z}^{|I|}$.

3.3.19. A finitely generated commutative $\mathbf{k}$-algebra is called a *complete intersection algebra* if it is the quotient of a polynomial algebra by a regular sequence. Observe that a complete intersection algebra is Cohen–Macaulay. Show that a face ring $\mathbf{k}[\mathcal{K}]$ is a complete intersection algebra if and only if it is isomorphic to the quotient of the form

$$\mathbf{k}[v_1, \dots, v_m]/(v_1 v_2 \cdots v_{k_1}, v_{k_1+1} v_{k_1+2} \cdots v_{k_1+k_2}, \dots, v_{k_1+\dots+k_{p-1}+1} \cdots v_{k_1+\dots+k_p}).$$

This is equivalent to $\mathcal{K}$ being decomposable into a join of the form

$$\partial\Delta^{k_1-1} * \partial\Delta^{k_2-1} * \dots * \partial\Delta^{k_p-1} * \Delta^{m-s-1},$$

where $s = k_1 + \dots + k_p$ and the join factor Δ^{m-s-1} is void if $s = m$.

3.3.20. A Cohen–Macaulay complex is pure. (Hint: given a maximal simplex $J \in \mathcal{K}$, consider the ideal $\mathcal{I}_J = (v_i \colon i \notin J)$ in $\mathbf{k}[m]$, and show that

$$\operatorname{depth} \mathbf{k}[\mathcal{K}] \leqslant \dim(\mathbf{k}[m]/\mathcal{I}_J) = \dim \mathbf{k}[v_j \colon j \in J] = |J|.)$$

3.4. Gorenstein complexes and Dehn–Sommerville relations

Gorenstein rings are a class Cohen–Macaulay rings with a special duality property. As in the case of Cohen–Macaulayness, simplicial complexes whose face rings are Gorenstein play an important role in combinatorial commutative algebra. In a sense, non-acyclic Gorenstein complexes provide a 'best possible algebraic approximation' to sphere triangulations. We review here the most important aspects of Gorenstein complexes. The proofs of the main results of this section, in particular Theorems 3.4.2 and 3.4.4, require considerably more commutative algebraic techniques than those from the previous sections. We refer the reader to [**35**, Chapter 3] for the general theory of Gorenstein rings and the missing proofs.

We recall from Proposition 3.3.7 that nonzero Betti numbers of a Cohen–Macaulay complex $\mathcal{K}$ of dimension $n-1$ with m vertices appear up to homological degree $-(m-n)$, and $\beta^{-(m-n)}(\mathbf{k}[\mathcal{K}]) \neq 0$.

Definition 3.4.1. A Cohen–Macaulay complex $\mathcal{K}$ of dimension $n-1$ with m vertices is called *Gorenstein* (over a field $\mathbf{k}$) if $\beta^{-(m-n)}(\mathbf{k}[\mathcal{K}]) = 1$, that is, if

$\mathrm{Tor}^{-(m-n)}_{\mathbf{k}[m]}(\mathbf{k}[\mathcal{K}],\mathbf{k})\cong\mathbf{k}$. Furthermore, $\mathcal{K}$ is *Gorenstein** if $\mathcal{K}$ is Gorenstein and $\mathcal{K}=\operatorname{core}\mathcal{K}$ (see Definition 2.2.15).

Since $\mathcal{K}=\operatorname{core}(\mathcal{K})*\Delta^{s-1}$ for some s, we have $\mathbf{k}[\mathcal{K}]=\mathbf{k}[\operatorname{core}(\mathcal{K})]\otimes\mathbf{k}[s]$. Then Lemma A.3.5 implies that

$$\mathrm{Tor}^{-i}_{\mathbf{k}[m]}\bigl(\mathbf{k}[\mathcal{K}],\mathbf{k}\bigr)\cong\mathrm{Tor}^{-i}_{\mathbf{k}[m-s]}\bigl(\mathbf{k}[\operatorname{core}\mathcal{K}],\mathbf{k}\bigr).$$

Therefore, $\mathcal{K}$ is Gorenstein if and only if $\operatorname{core}\mathcal{K}$ is Gorenstein*.

As in the case of Cohen–Macaulay complexes, Gorenstein* complexes can be characterised topologically as follows.

Theorem 3.4.2 ([**208**, §II.5] or [**35**, Theorem 5.6.1]). *The following conditions are equivalent:*

(a) *$\mathcal{K}$ is a Gorenstein* complex over* $\mathbf{k}$*;*
(b) *for any simplex $I\in\mathcal{K}$ (including $I=\varnothing$) the subcomplex* $\operatorname{lk}I$ *has homology of a sphere of dimension* $\dim(\operatorname{lk}I)$*;*
(c) *for any $x\in|\mathcal{K}|$,*

$$H^i\bigl(|\mathcal{K}|,|\mathcal{K}|\setminus x;\mathbf{k}\bigr)=\widetilde{H}^i(\mathcal{K};\mathbf{k})=\begin{cases}\mathbf{k} & \text{if } i=\dim\mathcal{K};\\ 0 & \text{otherwise.}\end{cases}$$

In topology, polyhedra $|\mathcal{K}|$ satisfying the conditions of the previous theorem are sometimes called *generalised homology spheres* (‘generalised’ because a homology sphere is usually assumed to be a manifold). In particular, triangulated spheres are Gorenstein* complexes. Triangulated manifolds are not Gorenstein* or even Cohen–Macaulay in general (*Buchsbaum complexes* provide a proper algebraic approximation to triangulated manifolds, see [**208**, § II.8]). Nevertheless, the Tor-algebra of a Gorenstein* complex behaves like the cohomology algebra of a manifold: it satisfies *Poincaré duality.* This fundamental result was proved by Avramov and Golod for Noetherian local rings; here we state the graded version of their theorem in the case of face rings.

Definition 3.4.3. A graded commutative connected $\mathbf{k}$-algebra A is called a *Poincaré algebra* if it is finite dimensional over $\mathbf{k}$, i.e. $A=\bigoplus_{i=0}^{d}A^i$, and the $\mathbf{k}$-linear maps

$$\begin{aligned}A^i&\to\mathrm{Hom}_{\mathbf{k}}(A^{d-i},A^d),\\ a&\mapsto\varphi_a,\quad\text{where }\varphi_a(b)=ab\end{aligned}$$

are isomorphisms for $0\leqslant i\leqslant d$. The classical example of a Poincaré algebra is the cohomology algebra of a manifold.

Theorem 3.4.4 (Avramov–Golod, [**35**, Theorem 3.4.5]). *A simplicial complex $\mathcal{K}$ is Gorenstein* if and only if the algebra $T=\bigoplus_{i=0}^{d}T^i$, where $T^i=\mathrm{Tor}^{-i}_{\mathbf{k}[m]}(\mathbf{k}[\mathcal{K}],\mathbf{k})$ and $d=\max\{j\colon\ \mathrm{Tor}^{-j}_{\mathbf{k}[m]}(\mathbf{k}[\mathcal{K}],\mathbf{k})\neq0\}$, is a Poincaré algebra.*

Corollary 3.4.5. *Let $\mathcal{K}$ be a Gorensten* complex of dimension $n-1$ on the set $[m]$. Then the Betti numbers and the Poincaré series of the* Tor *groups satisfy*

$$\beta^{-i,2j}\bigl(\mathbf{k}[\mathcal{K}]\bigr)=\beta^{-(m-n)+i,\,2(m-j)}\bigl(\mathbf{k}[\mathcal{K}]\bigr),\quad 0\leqslant i\leqslant m-n,\ 0\leqslant j\leqslant m,$$

$$F\bigl(\mathrm{Tor}^{-i}_{\mathbf{k}[m]}(\mathbf{k}[\mathcal{K}],\mathbf{k});\ \lambda\bigr)=\lambda^{2m}F\bigl(\mathrm{Tor}^{-(m-n)+i}_{\mathbf{k}[m]}(\mathbf{k}[\mathcal{K}],\mathbf{k});\ \tfrac{1}{\lambda}\bigr).$$

Proof. Theorems 3.2.4 and 3.4.2 imply that

$$\beta^{-(m-n)}\big(\mathbf{k}[\mathcal{K}]\big)=\beta^{-(m-n),2m}\big(\mathbf{k}[\mathcal{K}]\big)=1.$$

We therefore have $d=m-n$ and $T^d=\mathrm{Tor}^{-(m-n),\,2m}_{\mathbf{k}[m]}(\mathbf{k}[\mathcal{K}],\mathbf{k})\cong\mathbf{k}$ in the notation of Theorem 3.4.4. Since the multiplication in the Tor-algebra preserves the bigrading, the isomorphisms $T^i\overset{\cong}{\to}\mathrm{Hom}_{\mathbf{k}}(T^{m-n-i},T^{m-n})$ from the definition of a Poincaré algebra can be refined to isomorphisms

$$T^{i,\,2j}\xrightarrow{\cong}\mathrm{Hom}_{\mathbf{k}}(T^{m-n-i,\,2(m-j)},T^{m-n,\,2m}),$$

where $T^{m-n,\,2m}\cong\mathbf{k}$. This implies the first identity, and the second is a direct corollary. □

As a further corollary we obtain the following symmetry property for the Poincaré series of the face ring:

Corollary 3.4.6. *If $\mathcal{K}$ is Gorenstein* of dimension $n-1$, then*

$$F\big(\mathbf{k}[\mathcal{K}],\lambda\big)=(-1)^nF\big(\mathbf{k}[\mathcal{K}],\tfrac{1}{\lambda}\big).$$

Proof. We apply Proposition A.2.1 to the minimal resolution of the $\mathbf{k}[m]$-module $\mathbf{k}[\mathcal{K}]$. Note that $F(\mathbf{k}[m];\lambda)=(1-\lambda^2)^{-m}$. It follows from the formula for the Poincaré series from Proposition A.2.1 and Proposition A.2.6 that

$$F\big(\mathbf{k}[\mathcal{K}];\lambda\big)=(1-\lambda^2)^{-m}\sum_{i=0}^{m-n}(-1)^iF\Big(\mathrm{Tor}^{-i}_{\mathbf{k}[m]}\big(\mathbf{k}[\mathcal{K}],\mathbf{k}\big);\lambda\Big).$$

Using Corollary 3.4.5, we calculate

$$\begin{aligned}F\big(\mathbf{k}[\mathcal{K}];\lambda\big)&=(1-\lambda^2)^{-m}\sum_{i=0}^{m-n}(-1)^i\lambda^{2m}F\Big(\mathrm{Tor}^{-(m-n)+i}_{\mathbf{k}[m]}\big(\mathbf{k}[\mathcal{K}],\mathbf{k}\big);\tfrac{1}{\lambda}\Big)=\\&=\big(1-(\tfrac{1}{\lambda})^2\big)^{-m}(-1)^m\sum_{j=0}^{m-n}(-1)^{m-n-j}F\Big(\mathrm{Tor}^{-j}_{\mathbf{k}[m]}\big(\mathbf{k}[\mathcal{K}],\mathbf{k}\big);\tfrac{1}{\lambda}\Big)=\\&=(-1)^nF\big(\mathbf{k}[\mathcal{K}];\tfrac{1}{\lambda}\big).\end{aligned}$$ □

Corollary 3.4.7. *The Dehn–Sommerville relations $h_i=h_{n-i}$ hold for any Gorenstein* complex of dimension $n-1$ (in particular, for any triangulation of an $(n-1)$-sphere).*

Proof. This follows from the explicit form of the Poincaré series for $\mathbf{k}[\mathcal{K}]$ (Theorem 3.1.10) and the previous corollary. □

The Dehn–Sommerville relations may be further generalised to wider classes of complexes and posets; we give some of these generalisations in Section 3.8 below.

Unlike the situation with Cohen–Macaulay complexes, a characterisation of h-vectors (or, equivalently, f-vectors) of Gorenstein complexes is not known:

Problem 3.4.8 (Stanley). Characterise the h-vectors of Gorenstein complexes.

If the *g-conjecture* (i.e. the inequalities of Theorem 1.4.14 (b) and (c)) holds for Gorenstein complexes, then this would imply a solution to the above problem.

Exercises.

3.4.9. A Gorenstein complex $\mathcal{K}$ is Gorenstein* if and only if it is *non-acyclic* (i.e. $\widetilde{H}^*(\mathcal{K};\mathbf{k}) \neq 0$).

3.4.10. Let $\mathcal{K}$ be a Gorenstein* complex of dimension $n-1$ on $[m]$. Show that

$$\widetilde{H}^k(\mathcal{K}_J) \cong \widetilde{H}^{n-2-k}(\mathcal{K}_{\widehat{J}})$$

for any $J \subset [m]$, where $\widehat{J} = [m] \setminus J$. This is known as *Alexander duality for non-acyclic Gorenstein complexes.*

3.5. Face rings of simplicial posets

The whole theory of face rings may be extended to simplicial posets (defined in Section 2.8), thereby leading to new important classes of rings in combinatorial commutative algebra and applications in toric topology.

The *face ring* $\mathbf{k}[\mathcal{S}]$ of a simplicial poset $\mathcal{S}$ was introduced by Stanley **[207]** as a quotient of certain graded polynomial ring by a homogeneous ideal determined by the poset relation in $\mathcal{S}$. The rings $\mathbf{k}[\mathcal{S}]$ have remarkable algebraic and homological properties, albeit they are much more complicated than the Stanley–Reisner face rings $\mathbf{k}[\mathcal{K}]$. Unlike $\mathbf{k}[\mathcal{K}]$, the ring $\mathbf{k}[\mathcal{S}]$ is not generated in the lowest positive degree. Face rings of simplicial posets were further studied by Duval **[80]** and Maeda–Masuda–Panov **[150]**, **[146]**, among others. *Cohen–Macaulay* and *Gorenstein** face rings are particularly important; both properties are topological, that is, depend only on the topological type of the geometric realisation $|\mathcal{S}|$.

As usual, we shall not distinguish between simplicial posets $\mathcal{S}$ and their geometric realisations (simplicial cell complexes) $|\mathcal{S}|$. Given two elements $\sigma, \tau \in \mathcal{S}$, we denote by $\sigma \vee \tau$ the set of their joins, and denote by $\sigma \wedge \tau$ the set of their *meets*. Whenever either of these sets consists of a single element, we use the same notation for this particular element of $\mathcal{S}$.

To make clear the idea behind the definition of the face ring of a simplicial poset, we first consider the case when $\mathcal{S}$ is a simplicial complex $\mathcal{K}$. Then $\sigma \wedge \tau$ consists of a single element (possibly $\varnothing$), and $\sigma \vee \tau$ is either empty or consists of a single element. We consider the graded polynomial ring $\mathbf{k}[v_\sigma \colon \sigma \in \mathcal{K}]$ with one generator v_σ of degree $\deg v_\sigma = 2|\sigma|$ for each simplex $\sigma \in \mathcal{K}$. The following proposition provides an alternative presentation of the face ring $\mathbf{k}[\mathcal{K}]$, with a larger set of generators:

Proposition 3.5.1. *There is a canonical isomorphism of graded rings*

$$\mathbf{k}[\mathcal{K}] \cong \mathbf{k}[v_\sigma \colon \sigma \in \mathcal{K}]/\mathcal{I}'_{\mathcal{K}},$$

where $\mathcal{I}'_{\mathcal{K}}$ is the ideal generated by the element $v_\varnothing - 1$ and all elements of the form

$$v_\sigma v_\tau - v_{\sigma\wedge\tau} v_{\sigma\vee\tau}.$$

Here we set $v_{\sigma\vee\tau} = 0$ whenever $\sigma \vee \tau$ is empty.

Proof. The isomorphism is established by the map taking v_σ to $\prod_{i\in\sigma} v_i$. The rest is left as an exercise. □

Now let $\mathcal{S}$ be an arbitrary simplicial poset with the vertex set $V(\mathcal{S}) = [m]$. In this case both $\sigma \vee \tau$ and $\sigma \wedge \tau$ may consist of more than one element, but $\sigma \wedge \tau$ consists of a single element whenever $\sigma \vee \tau$ is nonempty.



Figure 3.3. Simplicial cell complexes.

We consider the graded polynomial ring $\mathbf{k}[v_\sigma \colon \sigma \in \mathcal{S}]$ with one generator v_σ of degree $\deg v_\sigma = 2|\sigma|$ for every element $\sigma \in \mathcal{S}$.

Definition 3.5.2 ([**207**]). The *face ring* of a simplicial poset $\mathcal{S}$ is the quotient

$$\mathbf{k}[\mathcal{S}] = \mathbf{k}[v_\sigma \colon \sigma \in \mathcal{S}] \;/\; \mathcal{I}_{\mathcal{S}},$$

where $\mathcal{I}_{\mathcal{S}}$ is the ideal generated by the elements $v_{\hat{0}} - 1$ and

$$v_\sigma v_\tau - v_{\sigma\wedge\tau} \cdot \sum_{\eta \in \sigma\vee\tau} v_\eta. \tag{3.14}$$

The sum over the empty set is zero, so we have $v_\sigma v_\tau = 0$ in $\mathbf{k}[\mathcal{S}]$ if $\sigma \vee \tau$ is empty.

The grading may be refined to an $\mathbb{N}^m$-grading by setting $\operatorname{mdeg} v_\sigma = 2V(\sigma)$. Here $V(\sigma) \subset [m]$ is the vertex set of σ, and we identify subsets of $[m]$ with $(0,1)$-vectors in $\{0,1\}^m \subset \mathbb{N}^m$ as usual. In particular, $\operatorname{mdeg} v_i = 2\boldsymbol{e}_i$.

Example 3.5.3.

1. The simplicial cell complex shown in Fig. 3.3 (a) is obtained by gluing two segments along their boundaries and has rank 2. The vertices are $1, 2$ and we denote the 1-dimensional simplices by σ and τ. Then the face ring $\mathbf{k}[\mathcal{S}]$ is the quotient of the graded polynomial ring

$$\mathbf{k}[v_1, v_2, v_\sigma, v_\tau], \quad \deg v_1 = \deg v_2 = 2, \quad \deg v_\sigma = \deg v_\tau = 4$$

by the two relations

$$v_1 v_2 = v_\sigma + v_\tau, \quad v_\sigma v_\tau = 0.$$

2. The simplicial cell complex in Fig. 3.3 (b) is obtained by gluing two triangles along their boundaries and has rank 3. The vertices are $1, 2, 3$ and we denote the 1-dimensional simplices (edges) by e, f and g, and the 2-dimensional simplices by σ and τ. The face ring $\mathbf{k}[\mathcal{S}]$ is isomorphic to the quotient of the polynomial ring

$$\mathbf{k}[v_1, v_2, v_3, v_\sigma, v_\tau], \quad \deg v_1 = \deg v_2 = \deg v_3 = 2, \quad \deg v_\sigma = \deg v_\tau = 6$$

by the two relations

$$v_1 v_2 v_3 = v_\sigma + v_\tau, \quad v_\sigma v_\tau = 0.$$

The generators corresponding to the edges can be excluded because of the relations $v_e = v_1 v_2$, $v_f = v_2 v_3$ and $v_g = v_1 v_3$.

Remark. The ideal $\mathcal{I}_{\mathcal{S}}$ is generated by *straightening relations* (3.14); these relations allow us to express the product of any pair of generators via products of generators corresponding to pairs of ordered elements of the poset. This can be restated by saying that $\mathbf{k}[\mathcal{S}]$ is an example of an *algebra with straightening law* (ASL for short, also known as a *Hodge algebra*). Lemma 3.5.4 and Theorem 3.5.7

below reflect algebraic properties of ASL's, and may be restated in this generality. For more on the theory of ASL's see [**208**, § III.6] and [**35**, Chapter 7].

A monomial $v_{\sigma_1}^{i_1}v_{\sigma_2}^{i_2}\cdots v_{\sigma_k}^{i_k}\in\mathbf{k}[v_\sigma\colon\sigma\in\mathcal{S}]$ is *standard* if $\sigma_1<\sigma_2<\cdots<\sigma_k$.

LEMMA 3.5.4. *Any element of* $\mathbf{k}[\mathcal{S}]$ *can be written as a linear combination of standard monomials.*

PROOF. It is enough to prove the statement for elements of $\mathbf{k}[\mathcal{S}]$ represented by monomials in generators v_σ. We write such a monomial as $a=v_{\tau_1}v_{\tau_2}\cdots v_{\tau_k}$ where some of the τ_i may coincide. We need to show that any such monomial can be expressed as a sum of monomials $\sum v_{\sigma_1}\cdots v_{\sigma_l}$ with $\sigma_1\leqslant\cdots\leqslant\sigma_l$. We may assume by induction that $\tau_2\leqslant\cdots\leqslant\tau_k$. Using relation (3.14) we can replace a by

$$v_{\tau_1\wedge\tau_2}\Bigl(\sum_{\rho\in\tau_1\vee\tau_2}v_\rho\Bigr)v_{\tau_3}\cdots v_{\tau_k}.$$

Now the first two factors in each summand above correspond to ordered elements of $\mathcal{S}$. We proceed by replacing the products $v_\rho v_{\tau_3}$ by $v_{\rho\wedge\tau_3}(\sum_{\pi\in\rho\vee\tau_3}v_\pi)$. Since $\tau_1\wedge\tau_2\leqslant\rho\wedge\tau_3$, now the first three factors in each monomial are in order. Continuing this process, we obtain in the end a sum of monomials corresponding to totally ordered sets of elements of $\mathcal{S}$. □

We refer to the presentation from Lemma 3.5.4 as a *standard representation* of an element $a\in\mathbf{k}[\mathcal{S}]$.

Given $\sigma\in\mathcal{S}$, we define the corresponding *restriction homomorphism* as

$$s_\sigma\colon\mathbf{k}[\mathcal{S}]\to\mathbf{k}[\mathcal{S}]/(v_\tau\colon\tau\not\leqslant\sigma).$$

The following result is straightforward.

PROPOSITION 3.5.5. *Let* $|\sigma|=k$ *with* $V(\sigma)=\{i_1,\dots,i_k\}$. *Then the image of the homomorphism* s_σ *is the polynomial ring* $\mathbf{k}[v_{i_1},\dots,v_{i_k}]$.

The next result generalises Proposition 3.1.8 to simplicial posets.

THEOREM 3.5.6. *The direct sum*

$$s=\bigoplus\nolimits_{\sigma\in\mathcal{S}}s_\sigma\colon\mathbf{k}[\mathcal{S}]\longrightarrow\bigoplus_{\sigma\in\mathcal{S}}\mathbf{k}[v_i\colon i\in V(\sigma)]$$

of all restriction maps is a monomorphism.

PROOF. Take a nonzero element $a\in\mathbf{k}[\mathcal{S}]$ and write its standard representation. Fix a standard monomial $v_{\sigma_1}^{i_1}\cdots v_{\sigma_k}^{i_k}$ which enters into this decomposition with a nonzero coefficient. Allowing some of the exponents i_j to be zero, we may assume that σ_k is a maximal element in $\mathcal{S}$ and $|\sigma_j|=j$ for $1\leqslant j\leqslant k$. We shall prove that $s_{\sigma_k}(a)\neq0$. Identify $s_{\sigma_k}(\mathbf{k}[\mathcal{S}])$ with the polynomial ring $\mathbf{k}[t_1,\dots,t_k]$ (so that $t_j=v_{i_j}$ in the notation of Proposition 3.5.5). Then $s_{\sigma_k}(v_{\sigma_k})=t_1\cdots t_k$, and we may assume without loss of generality that $s_{\sigma_k}(v_{\sigma_j})=t_1\cdots t_j$ for $1\leqslant j\leqslant k$. Hence,

$$s_{\sigma_k}\bigl(v_{\sigma_1}^{i_1}\cdots v_{\sigma_k}^{i_k}\bigr)=t_1^{i_1}(t_1t_2)^{i_2}\cdots(t_1\cdots t_k)^{i_k}.$$

If we prove that no other monomial $v_{\tau_1}^{j_1}\cdots v_{\tau_m}^{j_m}$ is mapped by s_{σ_k} to the same element of $\mathbf{k}[t_1,\dots,t_k]$, then this would imply that $s_{\sigma_k}(a)\neq0$. Note that

$$s_{\sigma_k}(v_{\tau_1}^{j_1}\cdots v_{\tau_m}^{j_m})=0\qquad\text{if }\tau_i\not\leqslant\sigma_k\text{ for some }i\text{ with }j_i\neq0,$$

so that we may assume that $m = k$. Now suppose that

$$s_{\sigma_k}\big(v_{\sigma_1}^{i_1}\cdots v_{\sigma_k}^{i_k}\big) = s_{\sigma_k}\big(v_{\tau_1}^{j_1}\cdots v_{\tau_k}^{j_k}\big). \tag{3.15}$$

We shall prove that $v_{\sigma_1}^{i_1}\cdots v_{\sigma_k}^{i_k} = v_{\tau_1}^{j_1}\ldots v_{\tau_k}^{j_k}$. We may assume by induction that the 'tails' of these monomial coincide, that is, there is some q, $1 \leqslant q \leqslant k$, such that $i_p = j_p$ and $\sigma_p = \tau_p$ for $i_p \neq 0$ whenever $p > q$. We shall prove that $i_q = j_q$ and $\sigma_q = \tau_q$ if $i_q \neq 0$. We obtain from (3.15) that

$$\begin{aligned}s_{\sigma_k}\big(v_{\sigma_1}^{i_1}\cdots v_{\sigma_q}^{i_q}\big)(t_1\cdots t_{q+1})^{i_{q+1}}\cdots(t_1\cdots t_k)^{i_k} =\\ = s_{\sigma_k}\big(v_{\tau_1}^{j_1}\cdots v_{\tau_q}^{j_q}\big)(t_1\cdots t_{q+1})^{i_{q+1}}\cdots(t_1\cdots t_k)^{j_k},\end{aligned}$$

hence, $s_{\sigma_k}(v_{\sigma_1}^{i_1}\cdots v_{\sigma_q}^{i_q}) = s_{\sigma_k}(v_{\tau_1}^{j_1}\cdots v_{\tau_q}^{j_q})$. Let j_l be the last nonzero exponent in $v_{\tau_1}^{j_1}\cdots v_{\tau_q}^{j_q}$ (i.e. $j_{l+1} = \cdots = j_q = 0$). Then we also have $i_{l+1} = \cdots = i_q = 0$, as otherwise $s_{\sigma_k}(v_{\sigma_1}^{i_1}\cdots v_{\sigma_q}^{i_q})$ is divisible by $t_1\cdots t_{l+1}$, while $s_{\sigma_k}(v_{\tau_1}^{j_1}\cdots v_{\tau_q}^{j_q})$ is not. We also have $i_l = j_l$ and $\sigma_l = \tau_l$, since i_l is the maximal power of the monomial $t_1\cdots t_l$ which divides $s_{\sigma_k}(v_{\sigma_1}^{i_1}\cdots v_{\sigma_q}^{i_q})$. We conclude by induction that $v_{\sigma_1}^{i_1}\cdots v_{\sigma_k}^{i_k} = v_{\tau_1}^{j_1}\cdots v_{\tau_k}^{j_k}$, and $s_{\sigma_k}(a) \neq 0$. □

REMARK. The proof above also shows that the map $s = \bigoplus_\sigma s_\sigma$ in Theorem 3.5.6 can be defined as the sum over the maximal elements $\sigma \in \mathcal{S}$ only.

THEOREM 3.5.7. *The standard representation of an element* $a \in \mathbf{k}[\mathcal{S}]$ *is unique. In other words, the standard monomials* $v_{\sigma_1}^{i_1}\cdots v_{\sigma_k}^{i_k}$ *form a* $\mathbf{k}$*-basis of* $\mathbf{k}[\mathcal{S}]$.

PROOF. This follows directly from Lemma 3.5.4 and Theorem 3.5.6. □

Lemma 3.3.1, which describes hsop's in the face rings of simplicial complexes, can be readily extended to simplicial posets (the same proof based on the properties of the restriction map s works):

LEMMA 3.5.8. *Let* $\mathcal{S}$ *be a simplicial poset of rank* n. *A sequence of homogeneous elements* $\boldsymbol{t} = (t_1,\ldots,t_n)$ *of* $\mathbf{k}[\mathcal{S}]$ *is a homogeneous system of parameters if and only*

$$\dim_{\mathbf{k}}\big(\mathbf{k}[v_i\colon i\in V(\sigma)]/s_\sigma(\boldsymbol{t})\big) < \infty$$

for each element $\sigma \in \mathcal{S}$.

The *f-vector* of a simplicial poset $\mathcal{S}$ is $\boldsymbol{f}(\mathcal{S}) = (f_0,\ldots,f_{n-1})$, where $n-1 = \dim\mathcal{S}$ and f_i is the number of elements of rank $i+1$ (i.e. number of faces of dimension i in the simplicial cell complex). We also set $f_{-1} = 1$. The *h-vector* $\boldsymbol{h}(\mathcal{S}) = (h_0,\ldots,h_n)$ is then defined by (2.3).

The Poincaré series of the face ring $\mathbf{k}[\mathcal{S}]$ has exactly the same form as in the case of simplicial complexes:

THEOREM 3.5.9. *We have*

$$F\big(\mathbf{k}[\mathcal{S}];\lambda\big) = \sum_{k=0}^{n}\frac{f_{k-1}\lambda^{2k}}{(1-\lambda^2)^k} = \frac{h_0 + h_1\lambda^2 + \cdots + h_n\lambda^{2n}}{(1-\lambda^2)^n}.$$

PROOF. By Theorem 3.5.7, we need to calculate the Poincaré series of the $\mathbf{k}$-vector space generated by the monomials $v_{\sigma_1}^{i_1}\cdots v_{\sigma_k}^{i_k}$ with $\sigma_1 < \cdots < \sigma_k$. For every $\sigma \in \mathcal{S}$ denote by $\mathcal{M}_\sigma$ the set of such monomials with $\sigma_k = \sigma$ and $i_k > 0$. Let $|\sigma| = k$; consider the restriction homomorphism s_σ to the polynomial ring $\mathbf{k}[t_1,\ldots,t_k]$. Then $s_\sigma(\mathcal{M}_\sigma)$ is the set of monomials in $\mathbf{k}[t_1,\ldots,t_k]$ which are divisible by $t_1\cdots t_k$.

Therefore, the Poincaré series of the subspace generated by the set $\mathcal{M}_\sigma$ is $\frac{\lambda^{2k}}{(1-\lambda^2)^k}$. Now, to finish the proof of the first identity we note that $\mathcal{S}$ is the union $\bigcup_{\sigma\in\mathcal{S}}\mathcal{M}_\sigma$ of the nonintersecting subsets $\mathcal{M}_\sigma$. The second identity follows from (2.3). □

As we have seen in Exercise 3.1.15, the face ring $\mathbf{k}[\mathcal{K}]$ of a simplicial complex can be realised as the limit of a diagram of polynomial algebras over $\mathrm{CAT}^{op}(\mathcal{K})$. A similar description exists for the face ring $\mathbf{k}[\mathcal{S}]$:

Construction 3.5.10 ($\mathbf{k}[\mathcal{S}]$ as a limit). We consider the diagram $\mathbf{k}[\,\cdot\,]_\mathcal{S}$ similar to that of Exercise 3.1.15:

$$\begin{aligned}\mathbf{k}[\,\cdot\,]_\mathcal{S}\colon \mathrm{CAT}^{op}(\mathcal{S}) &\longrightarrow \mathrm{CGA},\\ \sigma &\longmapsto \mathbf{k}[v_i\colon i\in V(\sigma)],\end{aligned}$$

whose value on a morphism $\sigma\leqslant\tau$ is the surjection

$$\mathbf{k}[v_i\colon i\in V(\tau)]\to\mathbf{k}[v_i\colon i\in V(\sigma)]$$

sending each v_i with $i\notin V(\sigma)$ to zero.

Lemma 3.5.11. *We have*

$$\mathbf{k}[\mathcal{S}]=\lim\mathbf{k}[\,\cdot\,]_\mathcal{S}$$

where the limit is taken in the category CGA.

Proof. We set up a total order on the elements of $\mathcal{S}$ so that the rank function does not decrease, and proceed by induction. We therefore may assume the statement is proved for a simplicial poset $\mathcal{T}$, and need to prove it for $\mathcal{S}$ which is obtained from $\mathcal{T}$ by adding one element σ. Then $\mathcal{S}_{<\sigma}=\{\tau\in\mathcal{S}\colon\tau<\sigma\}$ is the face poset of the boundary of the simplex Δ^σ. Geometrically, we may think of $|\mathcal{S}|$ as obtained from $|\mathcal{T}|$ by attaching one simplex Δ^σ along its boundary (if $|\sigma|=1$, then Δ^σ is a single point, so $|\mathcal{S}|$ is a disjoint union of $|\mathcal{T}|$ and a point). We therefore need to prove that the following is a pullback diagram:

$$\begin{tikzcd} \mathbf{k}[\mathcal{S}] \arrow[r] \arrow[d] & \mathbf{k}[\mathcal{S}_{\leqslant\sigma}] \arrow[d] \\ \mathbf{k}[\mathcal{T}] \arrow[r] & \mathbf{k}[\mathcal{S}_{<\sigma}]. \end{tikzcd} \tag{3.16}$$

Here the vertical arrows map v_σ to 0, while the horizontal ones map v_τ to 0 for $\tau\nleqslant\sigma$. Denote by A the pullback of (3.16) with $\mathbf{k}[\mathcal{S}]$ dropped. We need to show that the natural map $\mathbf{k}[\mathcal{S}]\to A$ is an isomorphism.

Since the limits in CGA are created in the underlying category of graded $\mathbf{k}$-vector spaces, the space of A is the direct sum of $\mathbf{k}[\mathcal{T}]$ and $\mathbf{k}[\mathcal{S}_{\leqslant\sigma}]$ with the pieces $\mathbf{k}[\mathcal{S}_{<\sigma}]$ identified in both spaces. In other words,

$$A=T\oplus\mathbf{k}[\mathcal{S}_{<\sigma}]\oplus S, \tag{3.17}$$

where T is the complement to $\mathbf{k}[\mathcal{S}_{<\sigma}]$ in $\mathbf{k}[\mathcal{T}]$, and S is the complement to $\mathbf{k}[\mathcal{S}_{<\sigma}]$ in $\mathbf{k}[\mathcal{S}_{\leqslant\sigma}]$. By Theorem 3.5.7, the space $\mathbf{k}[\mathcal{S}_{<\sigma}]$ has basis of standard monomials $v_{\tau_1}^{j_1}v_{\tau_2}^{j_2}\cdots v_{\tau_k}^{j_k}$ with $\tau_k<\sigma$. Similarly, S has basis of those monomials with $\tau_k=\sigma$ and $j_k>0$, while T has basis of those monomials with $\tau_k\nleqslant\sigma$ and $j_k>0$. Yet another application of Theorem 3.5.7 gives a decomposition of $\mathbf{k}[\mathcal{S}]$ identical to (3.17): a standard basis monomial $v_{\tau_1}^{j_1}v_{\tau_2}^{j_2}\cdots v_{\tau_k}^{j_k}$ with $j_k>0$ has either $\tau_k\nleqslant\sigma$, or $\tau_k<\sigma$, or $\tau_k=\sigma$. These three possibilities map to T, $\mathbf{k}[\mathcal{S}_{<\sigma}]$ and S respectively. It follows that $\mathbf{k}[\mathcal{S}] \to A$ is an isomorphism of $\mathbf{k}$-vector spaces. Since it is an algebra map, it is also an isomorphism of algebras, thus finishing the proof. $\square$

The description of $\mathbf{k}[\mathcal{S}]$ as a limit has the following important corollary, describing the functorial properties of the face rings and generalising Proposition 3.1.5.

PROPOSITION 3.5.12. *Let $f\colon \mathcal{S} \to \mathcal{T}$ be a rank-preserving map of simplicial posets. Define a homomorphism*

$$f^*\colon \mathbf{k}[w_\tau\colon \tau \in \mathcal{T}] \to \mathbf{k}[v_\sigma\colon \sigma \in \mathcal{S}], \quad f^*(w_\tau) = \sum_{\sigma \in f^{-1}(\tau),\ |\sigma|=|\tau|} v_\sigma.$$

Then f^* descends to a ring homomorphism $\mathbf{k}[\mathcal{T}] \to \mathbf{k}[\mathcal{S}]$, which we continue to denote by f^*.

PROOF. The poset map f gives rise to a functor $f\colon \textsc{cat}^{op}(\mathcal{S}) \to \textsc{cat}^{op}(\mathcal{T})$ and therefore to a natural transformation

$$f^*\colon [\textsc{cat}^{op}(\mathcal{T}), \textsc{cga}] \to [\textsc{cat}^{op}(\mathcal{S}), \textsc{cga}],$$

where $[\textsc{cat}^{op}(\mathcal{S}), \textsc{cga}]$ denotes the functors from $\textsc{cat}^{op}(\mathcal{S})$ to $\textsc{cga}$. It is easy to see that $f^*\,\mathbf{k}[\,\cdot\,]_{\mathcal{T}} = \mathbf{k}[\,\cdot\,]_{\mathcal{S}}$ in the notation of Construction 3.5.10, so we have the induced map of limits $f^*\colon \mathbf{k}[\mathcal{T}] \to \mathbf{k}[\mathcal{S}]$. We also have that $f^*(w_\tau) = \sum_{\sigma \in f^{-1}(\tau)} v_\sigma$ by the construction of lim in $\textsc{cga}$. $\square$

EXAMPLE 3.5.13. The folding map (2.9) induces a monomorphism $\mathbf{k}[\mathcal{K}_{\mathcal{S}}] \to \mathbf{k}[\mathcal{S}]$, which embeds $\mathbf{k}[\mathcal{K}_{\mathcal{S}}]$ in $\mathbf{k}[\mathcal{S}]$ as the subring generated by the elements v_i.

REMARK. An attempt to prove Proposition 3.5.12 directly from the definition, by showing that $f^*(\mathcal{I}_{\mathcal{T}}) \subset \mathcal{I}_{\mathcal{S}}$, runs into a complicated combinatorial analysis of the poset structure. This is an example of a situation where the use of an abstract categorical description of $\mathbf{k}[\mathcal{S}]$ proves to be beneficial.

Let $\mathbf{k}[m] = \mathbf{k}[v_1, \ldots, v_m]$ be the polynomial algebra on m generators of degree 2 corresponding to the vertices of $\mathcal{S}$. The face ring $\mathbf{k}[\mathcal{S}]$ acquires a $\mathbf{k}[m]$-algebra structure via the map $\mathbf{k}[m] \to \mathbf{k}[\mathcal{S}]$ sending each v_i identically. (Unlike the case of simplicial complexes, this map is generally not surjective.) We thereby obtain a $\mathbb{Z} \oplus \mathbb{N}^m$-graded Tor-algebra of $\mathbf{k}[\mathcal{S}]$:

$$\operatorname{Tor}_{\mathbf{k}[v_1,\ldots,v_m]}\bigl(\mathbf{k}[\mathcal{S}], \mathbf{k}\bigr) = \bigoplus_{i \geqslant 0,\, \boldsymbol{a} \in \mathbb{N}^m} \operatorname{Tor}^{-i,\,2\boldsymbol{a}}_{\mathbf{k}[v_1,\ldots,v_m]}\bigl(\mathbf{k}[\mathcal{S}], \mathbf{k}\bigr),$$

by analogy with Construction 3.2.8 for simplicial complexes.

We finish this section by stating a generalisation of Hochster's theorem to simplicial posets, and deriving some of its corollaries.

THEOREM 3.5.14 (Duval **[80]**, see also **[142]**). *For any subset $J \subset [m]$ we have*

$$\operatorname{Tor}^{-i,2J}_{\mathbf{k}[v_1,\ldots,v_m]}\bigl(\mathbf{k}[\mathcal{S}], \mathbf{k}\bigr) \cong \widetilde{H}^{|J|-i-1}(|\mathcal{S}_J|; \mathbf{k}),$$

where $\mathcal{S}_J$ the subposet of $\mathcal{S}$ consisting of those σ for which $V(\sigma) \subset J$. Also, $\operatorname{Tor}^{-i,2\boldsymbol{a}}_{\mathbf{k}[m]}(\mathbf{k}[\mathcal{S}], \mathbf{k}) = 0$ if $\boldsymbol{a}$ is not a $(0,1)$-vector.

PROOF. The argument follows the lines of the proof of Theorem 3.2.9. We define the quotient differential graded algebra

$$R^*(\mathcal{S}) = \Lambda[u_1, \ldots, u_m] \otimes \mathbf{k}[\mathcal{S}]/\mathcal{I}_R$$

where $\mathcal{I}_R$ is the ideal generated by the elements

$$u_i v_\sigma \quad \text{with } i \in V(\sigma), \quad \text{and} \quad v_\sigma v_\tau \quad \text{with } \sigma \wedge \tau \neq \hat{0}.$$

Note that the latter condition is equivalent to $V(\sigma) \cap V(\tau) \neq \varnothing$.

Then we need to prove the analogue of Lemma 3.2.6, that is, to show that the quotient projection

$$\varrho\colon \Lambda[u_1, \ldots, u_m] \otimes \mathbf{k}[\mathcal{S}] \to R^*(\mathcal{S})$$

induces an isomorphism in cohomology. This can be done by providing the appropriate chain homotopy, as in the proof of Lemma 3.2.6, but the formulae will be more complicated. Alternatively, we can use a topological argument, see the proof of Theorem 4.9.6 below.

Let $\mathcal{C}^{p-1}(|\mathcal{S}_J|)$ denote the $(p-1)$th cellular cochain group of $|\mathcal{S}_J|$ with coefficients in $\mathbf{k}$. It has a basis of cochains α_σ corresponding to elements $\sigma \in \mathcal{S}_J$ with $|\sigma| = p$. We define a $\mathbf{k}$-linear map

$$f\colon \mathcal{C}^{p-1}(|\mathcal{S}_J|) \longrightarrow R^{p-|J|,2J}(\mathcal{S}), \quad \alpha_\sigma \longmapsto \varepsilon(V(\sigma), J)\, u_{J\setminus V(\sigma)} v_\sigma,$$

where $\varepsilon(V(\sigma), J)$ is the sign from the proof of Theorem 3.2.9. This map is an isomorphism of cochain complexes; the details are left to the reader. Therefore,

$$\widetilde{H}^{p-1}(|\mathcal{S}_J|) \cong \operatorname{Tor}^{p-|J|,2J}_{\mathbf{k}[v_1,\ldots,v_m]}\bigl(\mathbf{k}[\mathcal{S}], \mathbf{k}\bigr),$$

which is equivalent to the first required isomorphism. Since $R^{-i,2\boldsymbol{a}}(\mathcal{S}) = 0$ if $\boldsymbol{a}$ is not a $(0,1)$-vector, $\operatorname{Tor}^{-i,2\boldsymbol{a}}_{\mathbf{k}[m]}(\mathbf{k}[\mathcal{K}], \mathbf{k})$ vanishes for such $\boldsymbol{a}$. □

We define the *multigraded algebraic Betti numbers* of $\mathbf{k}[\mathcal{S}]$ as

$$\beta^{-i,\,2\boldsymbol{a}}\bigl(\mathbf{k}[\mathcal{S}]\bigr) = \dim_{\mathbf{k}} \operatorname{Tor}^{-i,\,2\boldsymbol{a}}_{\mathbf{k}[v_1,\ldots,v_m]}\bigl(\mathbf{k}[\mathcal{S}], \mathbf{k}\bigr),$$

for $0 \leqslant i \leqslant m$, $\boldsymbol{a} \in \mathbb{N}^m$. We also set

$$\beta^{-i}\bigl(\mathbf{k}[\mathcal{S}]\bigr) = \dim_{\mathbf{k}} \operatorname{Tor}^{-i}_{\mathbf{k}[v_1,\ldots,v_m]}\bigl(\mathbf{k}[\mathcal{S}], \mathbf{k}\bigr) = \sum_{\boldsymbol{a}\in\mathbb{N}^m} \beta^{-i,\,2\boldsymbol{a}}\bigl(\mathbf{k}[\mathcal{S}]\bigr).$$

Example 3.5.15. Let $\mathcal{S}$ be the simplicial poset of Example 3.5.3.1. By Theorem 3.5.14, $\beta^{0,(0,0)}(\mathbf{k}[\mathcal{S}]) = \beta^{0,(2,2)}(\mathbf{k}[\mathcal{S}]) = 1$, and the other Betti numbers are zero. This implies that $\mathbf{k}[\mathcal{S}]$ is a free $\mathbf{k}[v_1, v_2]$-module with two generators, 1 and v_σ, of degree 0 and 4 respectively.

Note that unlike the case of simplicial complexes, $\beta^0(\mathbf{k}[\mathcal{S}])$ may be larger than 1. In fact, the following proposition follows easily from Theorem 3.5.14.

Proposition 3.5.16. *The number of generators of* $\mathbf{k}[\mathcal{S}]$ *as a* $\mathbf{k}[m]$*-module equals*

$$\beta^0(\mathbf{k}[\mathcal{S}]) = \sum_{J\subset[m]} \dim \widetilde{H}^{|J|-1}\bigl(|\mathcal{S}_J|\bigr).$$

Exercises.

3.5.17. Finish the proof of Proposition 3.5.1.

3.5.18. Fill in the details in the proof of Theorem 3.5.14.

3.5.19. Calculate the multigraded Betti numbers for the simplicial poset of Example 3.5.3.2.

Face rings: additional topics

3.6. Cohen–Macaulay simplicial posets

Assume given a property A of simplicial complexes. Then we can extend this property to posets by postulating that a poset $\mathcal{P}$ has the property A if the order complex $\operatorname{ord}(\mathcal{P})$ (see Definition 2.3.6) has the property A. In particular, Cohen–Macaulay and Gorenstein posets can be defined in this way. Simplicial posets $\mathcal{S}$ are of particular interest to us; in this case the order complex is identified with the barycentric subdivision $\mathcal{S}'$ (to be precise, with the cone over the barycentric subdivision, as we include the empty simplex, but this difference is inessential for the definitions to follow).

Definition 3.6.1. A simplicial poset $\mathcal{S}$ is *Cohen–Macaulay* (over $\mathbf{k}$) if its barycentric subdivision $\mathcal{S}'$ is a Cohen–Macaulay simplicial complex.

By definition, $\mathcal{S}$ is a Cohen–Macaulay simplicial poset if and only if the face ring $\mathbf{k}[\mathcal{S}']$ is Cohen–Macaulay. Since the face ring is also defined for the face poset $\mathcal{S}$ itself (and not only for its barycentric subdivision), it is perfectly natural to ask whether the class of Cohen–Macaulay simplicial posets admits an intrinsic description in terms of their face rings $\mathbf{k}[\mathcal{S}]$. One would achieve such a description by proving that the ring $\mathbf{k}[\mathcal{S}']$ is Cohen–Macaulay if and only if the ring $\mathbf{k}[\mathcal{S}]$ is Cohen–Macaulay. The 'if' part follows from the general theory of ASL's, see [**207**, Corollary 3.7]. The 'only if' part was proved in [**146**]; the proof uses the decomposition of the barycentric subdivision into a sequence of stellar subdivisions and then goes on to show that the Cohen–Macaulay property is preserved under stellar subdivisions. We include this characterisation of Cohen–Macaulay simplicial posets in terms of their face rings in Theorem 3.6.7 below.

Since many of the constructions in this section are geometric, we often talk about simplicial cell complexes rather than simplicial posets. We say that a simplicial subdivision of a simplicial cell complex $\mathcal{S}$ is *regular* if it is a simplicial complex. For instance, the barycentric subdivision is regular. Since the Cohen–Macaulayness of a simplicial complex is a topological property (see Proposition 3.3.11), we have the following statement.

Proposition 3.6.2. *The following conditions are equivalent:*

(a) *the barycentric subdivision of a simplicial cell complex $\mathcal{S}$ is a Cohen–Macaulay complex;*
(b) *any regular subdivision of $\mathcal{S}$ is a Cohen–Macaulay complex;*
(c) *a regular subdivision of $\mathcal{S}$ is a Cohen–Macaulay complex.*

As a further corollary we obtain that Proposition 3.3.11 itself extends to simplicial cell complexes, i.e. the property of a simplicial cell complex to be Cohen–Macaulay is also topological.

By analogy with Definition 2.2.13, we define the *star* and the *link* of $\sigma \in \mathcal{S}$ as the following subcomplexes:

$$\begin{aligned}
&\operatorname{st}_{\mathcal{S}} \sigma = \{\tau \in \mathcal{S} \colon \sigma \vee \tau \text{ is nonempty}\};\\
&\operatorname{lk}_{\mathcal{S}} \sigma = \{\tau \in \mathcal{S} \colon \sigma \vee \tau \text{ is nonempty, and } \tau \wedge \sigma = \hat{0}\}.
\end{aligned}$$

Remark. If $\mathcal{S}$ is a simplicial complex, then the poset $\operatorname{lk}_{\mathcal{S}} \sigma$ is isomorphic to the open semi-interval

$$\mathcal{S}_{>\sigma} = \{\rho \in \mathcal{S} \colon \rho > \sigma\},$$

and $|\operatorname{st}_{\mathcal{S}} \sigma| \cong \Delta^{\sigma} * |\operatorname{lk}_{\mathcal{S}} \sigma|$, where $*$ denotes the join. However, none of these isomorphisms holds for general $\mathcal{S}$, see Example 3.6.5 below.

Because of this remark, we cannot simply extend the definition of stellar subdivisions (Definition 2.7.1) to simplicial cell complexes. Instead, we define the *stellar subdivision* $\operatorname{ss}_{\sigma} \mathcal{S}$ of $\mathcal{S}$ at σ as the simplicial cell complex obtained by stellarly subdividing each face containing σ in a compatible way.

Proposition 3.6.3. *The barycentric subdivision $\mathcal{S}'$ can be obtained as a sequence of stellar subdivisions, one at each face $\sigma \in \mathcal{S}$, starting from the maximal faces. Moreover, each stellar subdivision in the sequence is applied to a face whose star is a simplicial complex.*

Proof. Assume $\dim \mathcal{S} = n-1$. We start by applying to $\mathcal{S}$ stellar subdivisions at all $(n-1)$-dimensional faces. Denote the resulting complex by $\mathcal{S}_1$. The $(n-2)$-faces of $\mathcal{S}_1$ are of two types: the ‘old’ ones, remaining from $\mathcal{S}$, and the ’new’ ones, appearing as the result of the stellar subdivisions. Then we take stellar subdivisions of $\mathcal{S}_1$ at all ‘old’ $(n-2)$-faces, and denote the result by $\mathcal{S}_2$. Next we apply to $\mathcal{S}_2$ stellar subdivisions at all $(n-3)$-faces remaining from $\mathcal{S}$. Proceeding in this way, at the end we get $\mathcal{S}_{n-1} = \mathcal{S}'$. To prove the second statement, consider two subsequent complexes $\mathcal{R}$ and $\widetilde{\mathcal{R}}$ in the sequence, so that $\widetilde{\mathcal{R}}$ is obtained from $\mathcal{R}$ by a single stellar subdivision at some $\sigma \in \mathcal{S}$. Then $\operatorname{st}_{\mathcal{R}} \sigma$ is isomorphic to $\Delta^{\sigma} * (\mathcal{S}_{>\sigma})'$ and therefore it is a simplicial complex. □

We proceed with two lemmata necessary to prove our main result.

Lemma 3.6.4. *Let $\mathcal{S}$ be a simplicial poset of rank n with vertex set $V(\mathcal{S}) = [m]$, and assume that the first k vertices span a face σ. Assume further that $\operatorname{st}_{\mathcal{S}} \sigma$ is a simplicial complex, and let $\widetilde{\mathcal{S}}$ be the stellar subdivision of $\mathcal{S}$ at σ. Let v denote the degree-two generator of $\mathbf{k}[\widetilde{\mathcal{S}}]$ corresponding to the added vertex. Then there exists a unique homomorphism $\beta \colon \mathbf{k}[\mathcal{S}] \to \mathbf{k}[\widetilde{\mathcal{S}}]$ such that*

$$\begin{aligned}
&v_\tau \mapsto v_\tau && \text{for } \tau \notin \operatorname{st}_{\mathcal{S}} \sigma;\\
&v_i \mapsto v + v_i, && \text{for } i = 1, \dots, k;\\
&v_i \mapsto v_i, && \text{for } i = k+1 \dots, m.
\end{aligned}$$

Moreover, β is injective, and if $\boldsymbol{t}$ is an hsop in $\mathbf{k}[\mathcal{S}]$, then $\beta(\boldsymbol{t})$ is an hsop in $\mathbf{k}[\widetilde{\mathcal{S}}]$.

Proof. In order to define the map β we first need to specify the images of v_τ for all $\tau \in \operatorname{st}_{\mathcal{S}} \sigma$. Choose such a v_τ and let $V(\tau) = \{i_1, \dots, i_\ell\}$ be its vertex set. Then we have the following identity in the ring $\mathbf{k}[\mathcal{S}] = \mathbf{k}[v_\tau \colon \tau \in \mathcal{S}]/\mathcal{I}_{\mathcal{S}}$:

$$v_{i_1} \cdots v_{i_\ell} = v_\tau + \sum_{\eta \colon V(\eta) = V(\tau),\, \eta \neq \tau} v_\eta. \tag{3.18}$$

For any v_η in the latter sum we have $\eta \notin \mathrm{st}_{\mathcal{S}}\,\sigma$, since $\mathrm{st}_{\mathcal{S}}\,\sigma$ is a simplicial complex, in which any set of vertices spans at most one face. Since β is already defined on the product on the left hand side and on the sum on the right hand side above, this determines $\beta(v_\tau)$ uniquely.

We therefore obtain a map of polynomial algebras $\mathbf{k}[v_\tau\colon \tau\in\mathcal{S}]\to\mathbf{k}[v_\tau\colon\tau\in\widetilde{\mathcal{S}}]$ (which we denote by the same letter β for a moment), and need to check that it descends to a map of face rings, $\mathbf{k}[\mathcal{S}]\to\mathbf{k}[\widetilde{\mathcal{S}}]$. In other words, we need to verify that $\beta(\mathcal{I}_{\mathcal{S}})\subset\mathcal{I}_{\widetilde{\mathcal{S}}}$.

It is clear from the definition of β that we have the commutative diagram

$$\begin{tikzcd}
\mathbf{k}[v_\tau\colon \tau\in\mathcal{S}] \arrow[r, "p"] \arrow[d, "\beta"] & \mathbf{k}[\mathcal{S}] \arrow[r, "s"] \arrow[d, dashed, "\beta"] & \bigoplus_{\tau\in\mathcal{S}}\mathbf{k}[v_i\colon i\in V(\tau)] \arrow[d, "s(\beta)"] \\
\mathbf{k}[v_\tau\colon \tau\in\widetilde{\mathcal{S}}] \arrow[r, "\widetilde{p}"] & \mathbf{k}[\widetilde{\mathcal{S}}] \arrow[r, "\widetilde{s}"] & \bigoplus_{\tau\in\widetilde{\mathcal{S}}}\mathbf{k}[v_i\colon i\in V(\tau)],
\end{tikzcd}$$

in which the middle vertical map is not yet defined. Here by s and $\widetilde{s}$ we denote the restriction maps from Theorem 3.5.6, and $s(\beta)$ is the map induced by β on the direct sum of polynomial algebras. Now let $x\in\mathcal{I}_{\mathcal{S}}$, i.e. $p(x)=0$. Then, by the commutativity of the diagram, $\widetilde{s}\widetilde{p}\beta(x)=0$. Since $\widetilde{s}$ is injective, we have $\widetilde{p}\beta(x)=0$. Hence, $\beta(x)\in\mathcal{I}_{\widetilde{\mathcal{S}}}$, which implies that the middle vertical map is well defined.

The last statement also follows from the commutative diagram above. The map $s(\beta)$ sends each direct summand of its domain isomorphically to at least one summand of its range, and therefore it is injective. Thus, $\beta\colon\mathbf{k}[\mathcal{S}]\to\mathbf{k}[\mathcal{S}']$ is also injective. The statement about hsop's then follows Lemma 3.5.8. □

Remark. If we defined the map β by sending each v_i identically, then it would still give rise to a ring homomorphism $\mathbf{k}[\mathcal{S}]\to\mathbf{k}[\widetilde{\mathcal{S}}]$, but the latter would not be injective (for example, it would map $v_\sigma\in\mathbf{k}[\mathcal{S}]$ to zero).

Example 3.6.5. The assumption on $\mathrm{st}_{\mathcal{S}}\,\sigma$ in Lemma 3.6.4 is not always satisfied. For example, if $\mathcal{S}$ is obtained by identifying two 2-simplices along their boundaries, and σ is any edge, then $\mathrm{st}_{\mathcal{S}}\,\sigma=\mathcal{S}$, which is not a simplicial complex.

Note also that if $\mathrm{st}_{\mathcal{S}}\,\sigma$ is not a simplicial complex, then the map $\beta\colon\mathbf{k}[\mathcal{S}]\to\mathbf{k}[\mathcal{S}']$ is not determined uniquely by the conditions specified in Lemma 3.6.4 (we cannot determine the images of v_τ with $\tau\in\mathrm{st}_{\mathcal{S}}\,\sigma$). Nevertheless, it is still possible to define the map $\beta\colon\mathbf{k}[\mathcal{S}]\to\mathbf{k}[\mathcal{S}']$ for an arbitrary simplicial poset $\mathcal{S}$, see Section 7.7.

Lemma 3.6.6. *Assume that* $\mathbf{k}[\mathcal{S}]$ *is a Cohen–Macaulay ring, and let* $\widetilde{\mathcal{S}}$ *be a stellar subdivision of* $\mathcal{S}$ *at* σ *such that* $\mathrm{st}_{\mathcal{S}}\,\sigma$ *is a simplicial complex. Then* $\mathbf{k}[\widetilde{\mathcal{S}}]$ *is a Cohen–Macaulay ring.*

Proof. We first prove that $\mathrm{st}_{\mathcal{S}}\,\sigma$ is a Cohen–Macaulay complex. Since $\mathrm{st}_{\mathcal{S}}\,\sigma=\Delta^\sigma*\mathrm{lk}_{\mathcal{S}}\,\sigma$, it is enough to verify that $\mathrm{lk}_{\mathcal{S}}\,\sigma$ is Cohen–Macaulay. This follows from Reisner's Theorem (Theorem 3.3.10) and the fact that simplicial cohomology of $\mathrm{lk}_{\mathcal{S}}\,\sigma$ is a direct summand in local cohomology of $\mathbf{k}[\mathcal{S}]$ (see [**208**, Theorem II.4.1] or [**35**, Theorem 5.3.8]).

Choose an hsop $\boldsymbol{t}=(t_1,\dots,t_n)$ in $\mathbf{k}[\mathcal{S}]$ and set $\widetilde{\boldsymbol{t}}=\beta(\boldsymbol{t})$. Let

$$p\colon\mathbf{k}[\mathcal{S}]\to\mathbf{k}[\mathcal{S}]/(v_\tau\colon\tau\notin\mathrm{st}_{\mathcal{S}}\,\sigma)=\mathbf{k}[\mathrm{st}_{\mathcal{S}}\,\sigma]$$

be the quotient projection. Set $R = \ker p$. Similarly, set

$$\widetilde{R} = \ker\bigl(\widetilde{p}\colon \mathbf{k}[\widetilde{\mathcal{S}}] \to \mathbf{k}[\mathrm{st}_{\widetilde{\mathcal{S}}}\, v]\bigr),$$

where v is the new vertex added in the process of stellar subdivision. Since the simplicial cell complexes $\mathcal{S}$ and $\widetilde{\mathcal{S}}$ do not differ on the complements of $\mathrm{st}_{\mathcal{S}}\,\sigma$ and $\mathrm{st}_{\widetilde{\mathcal{S}}}\, v$ respectively, the map β restricts to the identity isomorphism $R \to \widetilde{R}$. We therefore have the following commutative diagram with exact rows:

$$\begin{tikzcd}
0 \arrow[r] & R \arrow[r] \arrow[d, "\cong"] & \mathbf{k}[\mathcal{S}] \arrow[r, "p"] \arrow[d, "\beta"] & \mathbf{k}[\mathrm{st}_{\mathcal{S}}\,\sigma] \arrow[r] \arrow[d] & 0 \\
0 \arrow[r] & \widetilde{R} \arrow[r] & \mathbf{k}[\widetilde{\mathcal{S}}] \arrow[r, "\widetilde{p}"] & \mathbf{k}[\mathrm{st}_{\widetilde{\mathcal{S}}}\, v] \arrow[r] & 0,
\end{tikzcd}$$

Applying the functors $\otimes_{\mathbf{k}[\boldsymbol{t}]}\mathbf{k}$ and $\otimes_{\mathbf{k}[\widetilde{\boldsymbol{t}}]}\mathbf{k}$ to the diagram above, we get a map between the long exact sequences for Tor. Consider the following fragment:

$$\begin{tikzcd}
\mathrm{Tor}^{-2}_{\mathbf{k}[\boldsymbol{t}]}(\mathbf{k}[\mathrm{st}\,\sigma],\mathbf{k}) \arrow[r, "f"] \arrow[d] & \mathrm{Tor}^{-1}_{\mathbf{k}[\boldsymbol{t}]}(R,\mathbf{k}) \arrow[r] \arrow[d, "\cong"] & \mathrm{Tor}^{-1}_{\mathbf{k}[\boldsymbol{t}]}\,\mathbf{k}[\mathcal{S}],\mathbf{k}) \arrow[r] \arrow[d] & \mathrm{Tor}^{-1}_{\mathbf{k}[\boldsymbol{t}]}(\mathbf{k}[\mathrm{st}\,\sigma],\mathbf{k}) \arrow[d] \\
\mathrm{Tor}^{-2}_{\mathbf{k}[\widetilde{\boldsymbol{t}}]}(\mathbf{k}[\mathrm{st}\,v],\mathbf{k}) \arrow[r, "\widetilde{f}"] & \mathrm{Tor}^{-1}_{\mathbf{k}[\widetilde{\boldsymbol{t}}]}(\widetilde{R},\mathbf{k}) \arrow[r] & \mathrm{Tor}^{-1}_{\mathbf{k}[\widetilde{\boldsymbol{t}}]}(\mathbf{k}[\widetilde{\mathcal{S}}],\mathbf{k}) \arrow[r] & \mathrm{Tor}^{-1}_{\mathbf{k}[\widetilde{\boldsymbol{t}}]}(\mathbf{k}[\mathrm{st}\,v],\mathbf{k}).
\end{tikzcd}$$

Since $\mathbf{k}[\mathcal{S}]$ is Cohen–Macaulay, $\mathrm{Tor}^{-1}_{\mathbf{k}[\boldsymbol{t}]}(\mathbf{k}[\mathcal{S}],\mathbf{k}) = 0$ and the map f is surjective. Then $\widetilde{f}$ is also surjective. Since $\mathrm{st}_{\mathcal{S}}\,\sigma$ is a Cohen–Macaulay simplicial complex and $|\,\mathrm{st}_{\mathcal{S}}\,\sigma| \cong |\,\mathrm{st}_{\widetilde{\mathcal{S}}}\, v|$, Proposition 3.3.11 implies that $\mathbf{k}[\mathrm{st}\,v]$ is Cohen–Macaulay. Therefore, $\mathrm{Tor}^{-1}_{\mathbf{k}[\widetilde{\boldsymbol{t}}]}(\mathbf{k}[\mathrm{st}\,v],\mathbf{k}) = 0$. Since $\widetilde{f}$ is surjective, we also have $\mathrm{Tor}^{-1}_{\mathbf{k}[\widetilde{\boldsymbol{t}}]}(\mathbf{k}[\widetilde{\mathcal{S}}],\mathbf{k}) = 0$. Hence $\mathbf{k}[\widetilde{\mathcal{S}}]$ is free as a $\mathbf{k}[\widetilde{\boldsymbol{t}}]$-module (see [**145**, Lemma VII.6.2]) and thereby is Cohen–Macaulay. □

Now we can prove the main result of this section:

Theorem 3.6.7. *A simplicial poset $\mathcal{S}$ is Cohen–Macaulay if and only if the face ring $\mathbf{k}[\mathcal{S}]$ is Cohen–Macaulay.*

Proof. The fact that the face ring of a Cohen–Macaulay simplicial poset $\mathcal{S}$ is Cohen–Macaulay is proved in [**207**, Corollary 3.7] (see also [**208**, § III.6]) using the theory of ASL's.

Assume now that $\mathbf{k}[\mathcal{S}]$ is a Cohen–Macaulay ring. Since the barycentric subdivision $\mathcal{S}'$ is obtained by a sequence of stellar subdivisions, subsequent application of Lemma 3.6.6 shows that $\mathbf{k}[\mathcal{S}']$ is also Cohen–Macaulay. Thus, $\mathcal{S}'$ is a Cohen–Macaulay poset. □

We end this section by giving Stanley's characterisation of h-vectors of Cohen–Macaulay simplicial posets.

Theorem 3.6.8 (Stanley). *The integer vector $\boldsymbol{h} = (h_0, h_1, \dots, h_n)$ is the h-vector of a Cohen–Macaulay simplicial poset if and only if $h_0 = 1$ and $h_i \geqslant 0$.*

Proof. Let $\boldsymbol{h} = \boldsymbol{h}(\mathcal{S})$ for a Cohen–Macaulay simplicial poset $\mathcal{S}$. The condition $h_0 = 1$ follows from the definition of the h-vector, see (2.3). Let $\mathbf{k}$ be a field of zero characteristic, and $\boldsymbol{t} = (t_1, \dots, t_n)$ an lsop in $\mathbf{k}[\mathcal{S}]$ (since $\mathbf{k}[\mathcal{S}]$ is not generated by linear elements, the existence of an lsop is not automatic and is left as an exercise;

alternatively, see [**207**, Lemma 3.9]). Comparing the formula for the Poincaré series from Proposition A.3.14 with that of Theorem 3.5.9, we obtain

$$F\big(\mathbf{k}[\mathcal{S}]/\boldsymbol{t};\lambda\big)=h_0+h_1\lambda^2+\cdots+h_n\lambda^{2n}.$$

Hence, $h_i\geqslant 0$, as needed.

Now we construct a Cohen–Macaulay simplicial cell complex $\mathcal{S}$ with any given h-vector such that $h_0=1$ and $h_i\geqslant 0$. First note that $\boldsymbol{h}(\Delta^{n-1})=(1,0,\dots,0)$ and Δ^{n-1} is a Cohen–Macaulay simplicial (cell) complex. Now, given an $(n-1)$-dimensional Cohen–Macaulay simplicial cell complex $\mathcal{S}$ with the h-vector $(h_0,\dots,h_n)$, it suffices to construct, for any $k=1,\dots,n$, a new Cohen–Macaulay simplicial cell complex $\mathcal{S}_k$ with the h-vector given by

$$\boldsymbol{h}(\mathcal{S}_k)=(h_0,\dots,h_{k-1},h_k+1,h_{k+1},\dots,h_n). \tag{3.19}$$

To do this, we choose an $(n-1)$-face of $\mathcal{S}$, and in this face choose some k faces of dimension $n-2$. Then add to $\mathcal{S}$ a new $(n-1)$-simplex by attaching it along some k faces of dimension $n-2$ to the chosen k faces of $\mathcal{S}$. A direct check shows that the h-vector of the resulting simplicial cell complex $\mathcal{S}_k$ is given by (3.19). The fact that $\mathcal{S}_k$ is Cohen–Macaulay follows directly from Proposition 3.3.11. □

Note that this characterisation is substantially simpler than that for simplicial complexes (see Propositions 3.3.9 and the remark after it).

Exercises.

3.6.9. The map of face rings $\mathbf{k}[\mathcal{S}]\to\mathbf{k}[\widetilde{\mathcal{S}}]$ of Lemma 3.6.4 is not induced by any poset map $\widetilde{\mathcal{S}}\to\mathcal{S}$.

3.6.10. Let $\widetilde{\mathcal{S}}$ be a stellar subdivision of $\mathcal{S}$ at σ such that $\operatorname{st}_{\mathcal{S}}\sigma$ is a simplicial complex. Show that the ring $\mathbf{k}[\mathcal{S}]$ is Cohen–Macaulay if and only if $\mathbf{k}[\widetilde{\mathcal{S}}]$ is Cohen–Macaulay, i.e. the converse of Lemma 3.6.6 holds.

3.6.11. If $\mathbf{k}$ is of characteristic zero, then $\mathbf{k}[\mathcal{S}]$ admits an lsop.

3.7. Gorenstein simplicial posets

Gorenstein simplicial posets arise in toric topology as the combinatorial structures associated to the orbit quotients of *torus manifolds*, which are the subject of Chapter 7. It was exactly this particular feature of Gorenstein simplicial posets which allowed Masuda [**149**] to complete the characterisation of their h-vectors, conjectured by Stanley in [**207**]. We include Masuda's result here as Theorem 3.7.4.

Definition 3.7.1. A simplicial poset $\mathcal{S}$ is *Gorenstein* (respectively, *Gorenstein**) if its barycentric subdivision $\mathcal{S}'$ is a Gorenstein (respectively, Gorenstein*) simplicial complex.

Like Cohen–Macaulayness, the property of a simplicial poset $\mathcal{S}$ being Gorenstein* depends only on the topology of the realisation $|\mathcal{S}|$ (this follows from Theorem 3.4.2). In particular, simplicial cell subdivisions of spheres are Gorenstein*.

The problem of characterisation of h-vectors of Gorenstein* simplicial posets is more subtle than the corresponding question in the Cohen–Macaulay case. (Although this problem is much easier for simplicial posets than for simplicial complexes, see the discussion of the g-conjecture in Sections 2.5 and 3.4.)

Theorem 3.7.2. *Let* $\boldsymbol{h}(\mathcal{S}) = (h_0, h_1, \ldots, h_n)$ *be the* h*-vector of a Gorenstein* simplicial poset of rank* n*. Then* $h_0 = 1$*,* $h_i \geqslant 0$ *and* $h_i = h_{n-i}$ *for any* i*.*

Proof. The inequalities $h_i \geqslant 0$ follow from the fact that $\mathcal{S}$ is Cohen–Macaulay (Theorem 3.6.8). The identities $h_i = h_{n-i}$ will follow from the expression of the h-vector of the barycentric subdivision $\mathcal{S}'$ via $\boldsymbol{h}(\mathcal{S})$ and from the Dehn–Sommerville relations for the Gorenstein* simplicial complex $\mathcal{S}'$. Indeed, repeating the argument from Lemmata 2.3.4 and 2.3.5 we obtain the identity $\boldsymbol{h}(\mathcal{S}') = D\boldsymbol{h}(\mathcal{S})$, in which the vector $\boldsymbol{h}(\mathcal{S}')$ is symmetric, i.e. satisfies the Dehn–Sommerville relations. It can be checked directly using some identities for binomial coefficients that the operator D (and its inverse) takes symmetric vectors to symmetric ones (which is equivalent to the identity $d_{pq} = d_{n+1-p,n+1-q}$). This calculation can be avoided by using the following argument. The Dehn–Sommerville relations specify a linear subspace W of dimension $k = \left[\frac{n}{2}\right] + 1$ in the space $\mathbb{R}^{n+1}$ with coordinates $h_0, \ldots, h_n$. We need to check that this subspace is D-invariant. To do this it suffices to choose a basis $\boldsymbol{e}_1, \ldots, \boldsymbol{e}_k$ in W and check that $D\boldsymbol{e}_i \in W$ for all i. There is a basis in W consisting of h-vectors of simplicial spheres (and even simplicial polytopes, see the proof of Proposition 1.4.1). Since the barycentric subdivision of a simplicial sphere is a simplicial sphere, the vectors $D\boldsymbol{e}_i$, $1 \leqslant i \leqslant k$, are also symmetric, and W is a D-invariant subspace. Thus, the vector $\boldsymbol{h}(\mathcal{S}) = D^{-1}\boldsymbol{h}(\mathcal{S}')$ satisfies the Dehn–Sommerville relations. □

Theorem 3.7.3 ([**207**, Theorem 4.3]). *Let* $\boldsymbol{h} = (h_0, h_1, \ldots, h_n)$ *be an integer vector with* $h_0 = 1$*,* $h_i \geqslant 0$ *and* $h_i = h_{n-i}$*. Any of the following (mutually exclusive) conditions are sufficient for the existence of a Gorenstein* simplicial poset of rank* n *and* h*-vector* $h(\mathcal{S}) = \boldsymbol{h}$*:*

(a) *n is odd;*
(b) *n is even and $h_{n/2}$ is even;*
(c) *n is even, $h_{n/2}$ is odd, and $h_i > 0$ for all i.*

Proof. We start with the following two basic examples of $(n-1)$-dimensional simplicial cell complexes of dimension: $\partial\Delta^n$, with h-vector $\boldsymbol{h}(\partial\Delta^n) = (1, 1, \ldots, 1)$; and $\mathcal{S}_n$, the simplicial cell complex obtained by identifying two $(n-1)$-simplices along their boundaries, with $\boldsymbol{h}(\mathcal{S}_n) = (1, 0, \ldots, 0, 1)$. By applying the standard operations of join and connected sum (Constructions 2.2.8 and 2.2.11) to these two complexes we can obtain a simplicial cell complex with any prescribed h-vector satisfying the conditions of the theorem. Indeed, for $k \neq n-k$ we have

$$\boldsymbol{h}(\mathcal{S}_k * \mathcal{S}_{n-k}) = (1, 0, \ldots, 0, 1, 0, \ldots, 0, 1, 0, \ldots, 0, 1),$$

where $h_k = h_{n-k} = 1$, and the other entries are zero. Also, for $n = 2k$ we have

$$\boldsymbol{h}(\mathcal{S}_k * \mathcal{S}_k) = (1, 0, \ldots, 0, 2, 0, \ldots, 0, 1),$$

where $h_k = 2$. Now, by taking connected sum of the appropriate number of complexes $\partial\Delta^n$, $\mathcal{S}_n$ and $\mathcal{S}_k * \mathcal{S}_{n-k}$ and using the identity

$$h_i(\mathcal{S} \mathbin{\#} \widetilde{\mathcal{S}}) = h_i(\mathcal{S}) + h_i(\widetilde{\mathcal{S}}) \quad \text{for } 1 \leqslant i \leqslant n-1,$$

(see Example 1.3.8, which is valid for any two pure $(n-1)$-dimensional simplicial cell complexes), we obtain any required h-vector. □

The subtlest part of the characterisation of h-vectors of Gorenstein* simplicial posets is the following result, which was proved by Masuda [**149**].

THEOREM 3.7.4 (Masuda). *Let* $\boldsymbol{h}(\mathcal{S}) = (h_0, h_1, \ldots, h_n)$ *be the* h*-vector of a Gorenstein* simplicial poset* $\mathcal{S}$ *of even rank* n*, and let* $h_i = 0$ *for some* i*. Then the number* $h_{n/2}$ *is even.*

Note that the evenness of $h_{n/2}$ is equivalent to the evenness of the number of facets $f_{n-1} = \sum_{i=0}^{n} h_i$. The idea behind Masuda's proof of Theorem 3.7.4 lies within the topological theory of torus manifolds, see Section 7.2.3.

We combine the results of Theorems 3.7.2, 3.7.3 and 3.7.4 in the following characterisation result for the h-vectors of Gorenstein* simplicial posets.

THEOREM 3.7.5. *An integer vector* $\boldsymbol{h} = (h_0, h_1, \ldots, h_n)$ *is the* h*-vector of a Gorenstein* simplicial poset of rank* n *(or a simplicial cell subdivision of an* $(n-1)$*-sphere) if and only if the following conditions are satisfied:*

(a) $h_0 = 1$ *and* $h_i \geqslant 0$*;*
(b) $h_i = h_{n-i}$ *for all* i*;*
(c) *either* $h_i > 0$ *for all* i *or* $\sum_{i=0}^{n} h_i$ *is even.*

3.8. Generalised Dehn–Sommerville relations

In this section we obtain some further generalisations of the Dehn–Sommerville relations, in particular, to arbitrary triangulated manifolds.

Let $\mathcal{S}$ be a simplicial poset of rank n. Given $\sigma \in \mathcal{S}$, consider the closed upper semi-interval $\mathcal{S}_{\geqslant\sigma} = \{\tau \in \mathcal{S} \colon \tau \geqslant \sigma\}$ with the induced rank function, and set

$$\chi(\mathcal{S}_{\geqslant\sigma}) = \sum_{\tau\geqslant\sigma} (-1)^{|\tau|-1}. \tag{3.20}$$

A simplicial poset $\mathcal{S}$ of rank n satisfying $\chi(\mathcal{S}_{\geqslant\sigma}) = (-1)^{n-1}$ for all $\sigma \in \mathcal{S}$ is called *Eulerian*. According to a result of [**205**, (3.40)], the Dehn–Sommerville relations $h_i = h_{n-i}$ hold for Eulerian posets. This can be generalised as follows.

THEOREM 3.8.1 (see [**146**, Theorem 9.1]). *The following identity holds for the* h*-vector* $\boldsymbol{h}(\mathcal{S}) = (h_0, \ldots, h_n)$ *of a simplicial poset* S *of rank* n*:*

$$\sum_{i=0}^{n} (h_{n-i} - h_i) t^i = \sum_{\sigma\in\mathcal{S}} \Bigl(1 + (-1)^n \chi(\mathcal{S}_{\geqslant\sigma})\Bigr)(t-1)^{n-|\sigma|}.$$

In particular, if $\mathcal{S}$ *is Eulerian, then* $h_i = h_{n-i}$*.*

PROOF. We have

$$\begin{aligned}
\sum_{i=0}^{n} h_i t^i &= t^n \sum_{i=0}^{n} h_i \bigl(\tfrac{1}{t}\bigr)^{n-i} = t^n \sum_{i=0}^{n} f_{i-1} \bigl(\tfrac{1-t}{t}\bigr)^{n-i} \\
&= \sum_{i=0}^{n} f_{i-1} t^i (1-t)^{n-i} = \sum_{\tau\in\mathcal{S}} t^{|\tau|}(1-t)^{n-|\tau|} \\
&= \sum_{\tau\in\mathcal{S}} \sum_{\sigma\leqslant\tau} (t-1)^{|\tau|-|\sigma|}(1-t)^{n-|\tau|} = \sum_{\tau\in\mathcal{S}} \sum_{\sigma\leqslant\tau} (-1)^{n-|\tau|}(t-1)^{n-|\sigma|} \\
&= \sum_{\sigma\in\mathcal{S}} (t-1)^{n-|\sigma|} \sum_{\tau\geqslant\sigma} (-1)^{n-|\tau|} = \sum_{\sigma\in\mathcal{S}} (t-1)^{n-|\sigma|}(-1)^{n-1}\chi(\mathcal{S}_{\geqslant\sigma}),
\end{aligned} \tag{3.21}$$

where the fifth identity follows from the binomial expansion of the right hand side of the identity $t^{|\tau|} = ((t-1)+1)^{|\tau|}$ and the fact that $[\hat{0}, \tau] = \{\sigma \in \mathcal{S} \colon \sigma \leqslant \tau\}$ is a Boolean lattice of rank $|\tau|$.

On the other hand, we have

$$\sum_{i=0}^{n} h_{n-i}t^i = \sum_{i=0}^{n} h_i t^{n-i} = \sum_{i=0}^{n} f_{i-1}(t-1)^{n-i} = \sum_{\sigma\in\mathcal{S}}(t-1)^{n-|\sigma|}. \tag{3.22}$$

Subtracting (3.21) from (3.22) we obtain the required identity. □

As a corollary we obtain a generalisation of the Dehn–Sommerville relations to triangulated manifolds (which first appeared in [**44**, Corollary 4.5.4] as a corollary of bigraded Poincaré duality for moment-angle complexes):

Theorem 3.8.2. *Let $\mathcal{K}$ be a triangulation of a closed $(n-1)$-dimensional manifold. Then the h-vector $\boldsymbol{h}(\mathcal{K}) = (h_0,\dots,h_n)$ satisfies the identities*

$$h_{n-i} - h_i = (-1)^i\binom{n}{i}\bigl(\chi(\mathcal{K}) - \chi(S^{n-1})\bigr), \quad 0 \leqslant i \leqslant n.$$

Here $\chi(\mathcal{K}) = f_0 - f_1 + \dots + (-1)^{n-1}f_{n-1} = 1 + (-1)^{n-1}h_n$ is the Euler characteristic of $\mathcal{K}$ and $\chi(S^{n-1}) = 1 + (-1)^{n-1}$.

Proof. Viewing $\mathcal{K}$ as a simplicial poset, we calculate

$$\begin{aligned}\chi(\mathcal{K}_{\geqslant\sigma}) &= \sum_{\tau>\sigma}(-1)^{|\tau|-1} + (-1)^{|\sigma|-1} = (-1)^{|\sigma|}\Bigl(\sum_{\tau>\sigma}(-1)^{|\tau|-|\sigma|-1} - 1\Bigr)\\ &= (-1)^{|\sigma|}\Bigl(\sum_{\varnothing\neq\rho\in\operatorname{lk}_{\mathcal{K}}\sigma}(-1)^{|\rho|-1} - 1\Bigr) = (-1)^{|\sigma|}\bigl(\chi(\operatorname{lk}_{\mathcal{K}}\sigma) - 1\bigr).\end{aligned}$$

Here we have used the fact that the poset of nonempty faces of $\operatorname{lk}_{\mathcal{K}}\sigma$ is isomorphic to $\mathcal{S}_{>\sigma}$, with the rank function shifted by $|\sigma|$. Now since $\mathcal{K}$ is a triangulated $(n-1)$-dimensional manifold, the link of a nonempty face $\sigma \in \mathcal{K}$ has the homology of a sphere of dimension $(n-|\sigma|-1)$. Hence, $\chi(\operatorname{lk}_{\mathcal{K}}\sigma) = 1 + (-1)^{n-|\sigma|-1}$, and therefore $\chi(\mathcal{K}_{\geqslant\sigma}) = (-1)^{n-1}$ for $\sigma \neq \varnothing$. Also, $\operatorname{lk}_{\mathcal{K}}\varnothing = \mathcal{K}$. Now using the identity of Theorem 3.8.1 we calculate

$$\sum_{i=0}^{n}(h_{n-i}-h_i)t^i = \bigl(1+(-1)^n(\chi(\mathcal{K})-1)\bigr)(t-1)^n = (-1)^n\bigl(\chi(\mathcal{K})-\chi(S^{n-1})\bigr)(t-1)^n.$$

The required identity follows by comparing the coefficients of t^i. □

Exercises.

3.8.3. The identity of Theorem 3.8.2 holds for simplicial posets.

CHAPTER 4

Moment-angle complexes

This is the first genuinely 'toric' chapter of this book; it links the combinatorial and algebraic constructions of the previous chapters to the world of toric spaces.

The term 'moment-angle complex' refers to a decomposition of a certain toric space $\mathcal{Z}_{\mathcal{K}}$ into products of polydiscs and tori parametrised by simplices in a given simplicial complex $\mathcal{K}$. The underlying space $\mathcal{Z}_{\mathcal{K}}$ features in several important algebraic, symplectic and topological constructions of torus actions, which are the subject of the next three chapters. The decomposition of $\mathcal{Z}_{\mathcal{K}}$ as the 'moment-angle complex', which first appeared in [**42**], [**43**], provided an effective topological instrument for studying these spaces.

The basic building block in the 'moment-angle' decomposition of $\mathcal{Z}_{\mathcal{K}}$ is the pair (D^2, S^1) of a unit disc and circle, and the whole construction can be extended naturally to arbitrary pairs of spaces (X, A). The resulting complex $(X, A)^{\mathcal{K}}$ is now known as the 'polyhedral product space' over a simplicial complex $\mathcal{K}$; this terminology was suggested by William Browder, cf. [**11**]. Many spaces important for toric topology admit polyhedral product decompositions.

It has soon become clear that the construction of the moment-angle complex $\mathcal{Z}_{\mathcal{K}}$ and its generalisation $(X, A)^{\mathcal{K}}$ is of truly universal nature, and has remarkable functorial properties. The most basic of these is that the construction of $\mathcal{Z}_{\mathcal{K}}$ establishes a functor from simplicial complexes and simplicial maps to spaces with torus actions and equivariant maps. If $\mathcal{K}$ is a simplicial subdivision of a sphere (a triangulated sphere), then $\mathcal{Z}_{\mathcal{K}}$ is a manifold, and most important geometric examples of $\mathcal{Z}_{\mathcal{K}}$ arise in this way. In the case when A is a point, the polyhedral product $(X, pt)^{\mathcal{K}}$ interpolates between the m-fold wedge, or bouquet, of X (corresponding to m discrete points as $\mathcal{K}$) and the m-fold product of X (corresponding to the full simplex as $\mathcal{K}$). Parallel to the topological and geometric study of moment-angle complexes $\mathcal{Z}_{\mathcal{K}}$ and related toric spaces, a homotopy-theoretic study of polyhedral products $(X, A)^{\mathcal{K}}$ has now gained its own momentum. Basic homotopy properties of moment-angle complexes are given in Section 4.2, while more advanced homotopy-theoretic aspects of toric topology are the subject of Chapter 8.

The key result of this chapter is the calculation of the integral cohomology ring of $\mathcal{Z}_{\mathcal{K}}$, carried out in Section 4.4. The ring $H^*(\mathcal{Z}_{\mathcal{K}})$ is shown to be isomorphic to the Tor-algebra $\mathrm{Tor}_{\mathbb{Z}[v_1,\ldots,v_m]}(\mathbb{Z}[\mathcal{K}], \mathbb{Z})$, where $\mathbb{Z}[\mathcal{K}]$ is the face ring of $\mathcal{K}$. The canonical bigraded structure in the Tor groups thereby acquires a geometric interpretation in terms of the bigraded cell decomposition of $\mathcal{Z}_{\mathcal{K}}$. The calculation of $H^*(\mathcal{Z}_{\mathcal{K}})$ builds upon a construction of a ring model for *cellular* cochains of $\mathcal{Z}_{\mathcal{K}}$ and the corresponding cellular diagonal approximation, which is functorial with respect to maps of moment-angle complexes induced by simplicial maps of $\mathcal{K}$. This functorial property of the cellular diagonal approximation for $\mathcal{Z}_{\mathcal{K}}$ is quite special, due to the lack of such a construction for general cell complexes.

The construction of the moment-angle complex therefore brings the methods of equivariant topology to bear on the study of combinatorics of simplicial complexes, and gives a new geometric dimension to combinatorial commutative algebra. In particular, homological invariants of face rings, such as Tor-algebras or algebraic Betti numbers, can be now interpreted geometrically in terms of cohomology of moment-angle complexes. This link is explored further in Sections 4.5 and 4.9.

Another important aspect of the theory of moment-angle complexes is their connection to coordinate subspace arrangements and their complements. As we have already seen in Proposition 3.1.12, coordinate subspace arrangements arise as affine varieties corresponding to face rings. Their complements have played an important role in toric geometry and singularity theory, and, more recently, in the theory of linkages and robotic motion planning. Arrangements of coordinate subspaces in $\mathbb{C}^m$ correspond bijectively to simplicial complexes $\mathcal{K}$ on the set $[m]$, and the complement of such an arrangement deformation retracts onto the corresponding moment-angle complex $\mathcal{Z}_{\mathcal{K}}$. In particular, the moment-angle complex and the complement of the arrangement have the same homotopy type. We therefore may use the results on moment-angle complexes to obtain a description of the cohomology groups and cup product structure of a coordinate subspace arrangement complement. The formula obtained for the cohomology groups is related to the general formula of Goresky–MacPherson [**102**] by means of Alexander duality.

The material of this chapter, with the exception of 'Additional topics', is mainly a freshened and modernised exposition of the results obtained by the authors and their collaborators in [**43**], [**45**], [**20**], [**179**].

Spaces with torus actions, or *toric spaces*, will be the main players throughout the rest of this book. (See Appendix B.3 for the key concepts of the theory of group actions on topological spaces.) The most basic example of a toric space is the complex m-dimensional space $\mathbb{C}^m$, on which the *standard torus*

$$\mathbb{T}^m = \bigl\{\boldsymbol{t} = (t_1, \ldots, t_m) \in \mathbb{C}^m \colon |t_i| = 1 \quad \text{for } i = 1, \ldots, m\bigr\}$$

acts coordinatewise. That is, the action is given by

$$\begin{aligned}&\mathbb{T}^m \times \mathbb{C}^m \longrightarrow \mathbb{C}^m,\\&(t_1, \ldots, t_m) \cdot (z_1, \ldots, z_m) = (t_1 z_1, \ldots, t_m z_m).\end{aligned}$$

The quotient $\mathbb{C}^m/\mathbb{T}^m$ of this action is the *positive orthant*

$$\mathbb{R}^m_{\geqslant} = \bigl\{(y_1, \ldots, y_m) \in \mathbb{R}^m \colon y_i \geqslant 0 \quad \text{for } i = 1, \ldots, m\bigr\},$$

with the quotient projection given by

$$\begin{aligned}\mu\colon \mathbb{C}^m &\longrightarrow \mathbb{R}^m_{\geqslant},\\(z_1, \ldots, z_m) &\longmapsto (|z_1|^2, \ldots, |z_m|^2)\end{aligned}$$

(or by $(z_1, \ldots, z_m) \longmapsto (|z_1|, \ldots, |z_m|)$, but the former is usually more preferable).

We shall use the blackboard bold capital $\mathbb{T}$ in the notation for the standard torus $\mathbb{T}^m$ only, and use italic T^m to denote an abstract m-torus, i.e. a compact abelian Lie group isomorphic to a product of m circles. We shall also denote the standard unit circle by $\mathbb{S}$ or $\mathbb{T}$ occasionally, to distinguish it from an abstract circle S^1.

All homology and cohomology groups in this chapter are with integer coefficients, unless another coefficient group is explicitly specified.

4.1. Definitions and functorial properties

Moment-angle complex $\mathcal{Z}_{\mathcal{K}}$. We consider the unit polydisc in the m-dimensional complex space $\mathbb{C}^m$:

$$\mathbb{D}^m = \bigl\{(z_1,\dots,z_m)\in\mathbb{C}^m\colon |z_i|^2\leqslant 1 \quad \text{for } i=1,\dots,m\bigr\}.$$

The polydisc $\mathbb{D}^m$ is a $\mathbb{T}^m$-invariant subspace of $\mathbb{C}^m$, and the quotient $\mathbb{D}^m/\mathbb{T}^m$ is identified with the standard unit cube $\mathbb{I}^m\subset\mathbb{R}^m_{\geqslant}$.

Construction 4.1.1 (moment-angle complex). Let $\mathcal{K}$ be a simplicial complex on the set $[m]$. We recall the cubical subcomplex $\mathrm{cc}(\mathcal{K})$ in $\mathbb{I}^m$ from Construction 2.9.11, which subdivides $\operatorname{cone}\mathcal{K}$. The *moment-angle complex* $\mathcal{Z}_{\mathcal{K}}$ corresponding to $\mathcal{K}$ is defined from the pullback square

$$\begin{tikzcd}
\mathcal{Z}_{\mathcal{K}} \arrow[r, hook] \arrow[d] & \mathbb{D}^m \arrow[d, "\mu"] \\
\mathrm{cc}(\mathcal{K}) \arrow[r, hook] & \mathbb{I}^m
\end{tikzcd}$$

Explicitly, $\mathcal{Z}_{\mathcal{K}}=\mu^{-1}(\mathrm{cc}(\mathcal{K}))$. By construction, $\mathcal{Z}_{\mathcal{K}}$ is a $\mathbb{T}^m$-invariant subspace in the polydisc $\mathbb{D}^m$, and the quotient $\mathcal{Z}_{\mathcal{K}}/\mathbb{T}^m$ is homeomorphic to $|\operatorname{cone}\mathcal{K}|$.

Using the decomposition $\mathrm{cc}(\mathcal{K})=\bigcup_{I\in\mathcal{K}}C_I$ into faces, see (2.10), it follows that

$$\mathcal{Z}_{\mathcal{K}}=\bigcup_{I\in\mathcal{K}}B_I, \tag{4.1}$$

where

$$B_I=\mu^{-1}(C_I)=\bigl\{(z_1,\dots,z_m)\in\mathbb{D}^m\colon |z_j|^2=1 \text{ for } j\notin I\bigr\},$$

and the union in (4.1) is understood as the union of subsets inside the polydisc $\mathbb{D}^m$. Note that B_I is a product of $|I|$ discs and $m-|I|$ circles. Following our notational tradition, we denote a topological 2-disc (the underlying space of $\mathbb{D}$) by D^2. Then we may rewrite (4.1) as the following decomposition of $\mathcal{Z}_{\mathcal{K}}$ into products of discs and circles:

$$\mathcal{Z}_{\mathcal{K}}=\bigcup_{I\in\mathcal{K}}\Bigl(\prod_{i\in I}D^2\times\prod_{i\notin I}S^1\Bigr), \tag{4.2}$$

From now on we shall denote the space B_I by $(D^2,S^1)^I$. Obviously, the union in (4.1) or (4.2) can be taken over the maximal simplices $I\in\mathcal{K}$ only.

Using the abstract categorical language, we may consider the face category $\mathrm{CAT}(\mathcal{K})$ (faces and inclusions), and define the functor (or diagram)

$$\begin{aligned}\mathcal{D}_{\mathcal{K}}(D^2,S^1)\colon \mathrm{CAT}(\mathcal{K}) &\longrightarrow \mathrm{TOP},\\ I &\longmapsto B_I=(D^2,S^1)^I,\end{aligned}$$

which maps the morphism $I\subset J$ of $\mathrm{CAT}(\mathcal{K})$ to the inclusion of spaces $(D^2,S^1)^I\subset(D^2,S^1)^J$. Then we have

$$\mathcal{Z}_{\mathcal{K}}=\operatorname{colim}\mathcal{D}_{\mathcal{K}}(D^2,S^1)=\operatorname*{colim}_{I\in\mathcal{K}}(D^2,S^1)^I.$$

Example 4.1.2.
1. Let $\mathcal{K}=\Delta^{m-1}$ be the full simplex. Then $\mathrm{cc}(\mathcal{K})=\mathbb{I}^m$ and $\mathcal{Z}_{\mathcal{K}}=\mathbb{D}^m$.

2. Let $\mathcal{K}$ be a simplicial complex on $[m]$, and let $\mathcal{K}^\circ$ be the complex on $[m+1]$ obtained by adding one ghost vertex $\circ = \{m+1\}$ to $\mathcal{K}$. Then the cubical complex $\mathrm{cc}(\mathcal{K}^\circ)$ is contained in the facet $\{y_{m+1} = 1\}$ of the cube $\mathbb{I}^{m+1}$, and

$$\mathcal{Z}_{\mathcal{K}^\circ} = \mathcal{Z}_{\mathcal{K}} \times S^1.$$

In particular, if $\mathcal{K}$ is the 'empty' simplicial complex on $[m]$, consisting of the empty simplex $\varnothing$ only, then $\mathrm{cc}(\mathcal{K})$ is the vertex $(1,\ldots,1) \in \mathbb{I}^m$ and $\mathcal{Z}_{\mathcal{K}} = \mu^{-1}(1,\ldots,1) = \mathbb{T}^m$ is the standard m-torus.

For arbitrary $\mathcal{K}$ on $[m]$, the moment-angle complex $\mathcal{Z}_{\mathcal{K}}$ contains the m-torus $\mathbb{T}^m$ (corresponding to $\mathcal{K} = \varnothing$) and is contained in the polydisc $\mathbb{D}^m$ (corresponding to $\mathcal{K} = \Delta^{m-1}$).

3. Let $\mathcal{K}$ be the complex consisting of two disjoint points. Then

$$\mathcal{Z}_{\mathcal{K}} = (D^2 \times S^1) \cup (S^1 \times D^2) = \partial(D^2 \times D^2) \cong S^3,$$

the standard decomposition of a 3-sphere into the union of two solid tori.

4. More generally, if $\mathcal{K} = \partial\Delta^{m-1}$ (the boundary of a simplex), then

$$\begin{aligned}\mathcal{Z}_{\mathcal{K}} &= (D^2 \times \cdots \times D^2 \times S^1) \cup (D^2 \times \cdots \times S^1 \times D^2) \cup \cdots \cup (S^1 \times \cdots \times D^2 \times D^2)\\ &= \partial\big((D^2)^m\big) \cong S^{2m-1}.\end{aligned}$$

5. Let $\mathcal{K} =$ , the boundary of a 4-gon. Then we have four maximal simplices $\{1,3\}$, $\{2,3\}$, $\{1,4\}$ and $\{2,4\}$, and

$$\begin{aligned}\mathcal{Z}_{\mathcal{K}} &= (D^2 \times S^1 \times D^2 \times S^1) \cup (S^1 \times D^2 \times D^2 \times S^1)\\ &\qquad \cup (D^2 \times S^1 \times S^1 \times D^2) \cup (S^1 \times D^2 \times S^1 \times D^2)\\ &= \big((D^2 \times S^1) \cup (S^1 \times D^2)\big) \times D^2 \times S^1 \cup \big((D^2 \times S^1) \cup (S^1 \times D^2)\big) \times S^1 \times D^2\\ &= \big((D^2 \times S^1) \cup (S^1 \times D^2)\big) \times \big((D^2 \times S^1) \cup (S^1 \times D^2)\big) \cong S^3 \times S^3.\end{aligned}$$

In the last example, $\mathcal{K}$ is the join of $\{1,2\}$ and $\{3,4\}$. More generally,

Proposition 4.1.3. *Let $\mathcal{K} = \mathcal{K}_1 * \mathcal{K}_2$; then*

$$\mathcal{Z}_{\mathcal{K}} = \mathcal{Z}_{\mathcal{K}_1} \times \mathcal{Z}_{\mathcal{K}_2}.$$

Proof. By the definition of join (Construction 2.2.8), we have

$$\begin{aligned}\mathcal{Z}_{\mathcal{K}_1 * \mathcal{K}_2} &= \bigcup_{I_1 \in \mathcal{K}_1,\, I_2 \in \mathcal{K}_2} (D^2, S^1)^{I_1 \sqcup I_2} = \bigcup_{I_1 \in \mathcal{K}_1,\, I_2 \in \mathcal{K}_2} (D^2, S^1)^{I_1} \times (D^2, S^1)^{I_2}\\ &= \Big(\bigcup_{I_1 \in \mathcal{K}_1} (D^2, S^1)^{I_1}\Big) \times \Big(\bigcup_{I_2 \in \mathcal{K}_2} (D^2, S^1)^{I_2}\Big) = \mathcal{Z}_{\mathcal{K}_1} \times \mathcal{Z}_{\mathcal{K}_2}. \quad \square\end{aligned}$$

The topological structure of $\mathcal{Z}_{\mathcal{K}}$ is quite complicated even for simplicial complexes $\mathcal{K}$ with few vertices. Several different techniques will be developed to describe the topology of $\mathcal{Z}_{\mathcal{K}}$; this is one of the main subjects of the book. To get an idea on how $\mathcal{Z}_{\mathcal{K}}$ may look like we included Exercises 4.1.17 and 4.1.18, in which the topological structure of $\mathcal{Z}_{\mathcal{K}}$ is more complicated than in the examples above, but which are still accessible by relatively elementary topological methods. As we shall see below, the cohomology of $\mathcal{Z}_{\mathcal{K}}$ may have an arbitrary torsion (Corollary 4.4.9), as well as Massey products (Section 4.8).

Moment-angle complexes corresponding to triangulated spheres and manifolds are of particular interest:

Theorem 4.1.4. *Let $\mathcal{K}$ be a triangulation of an $(n-1)$-dimensional sphere with m vertices. Then $\mathcal{Z}_\mathcal{K}$ is a (closed) topological manifold of dimension $m+n$.*

If $\mathcal{K}$ is a triangulated manifold, then $\mathcal{Z}_\mathcal{K} \setminus \mu^{-1}(1,\dots,1)$ is an (open) non-compact manifold. Here $(1,\dots,1) \in \mathbb{I}^m$ is the cone vertex, and $\mu^{-1}(1,\dots,1) = \mathbb{T}^m$.

Proof. We first construct a decomposition of the polyhedron $|\operatorname{cone}(\mathcal{K}')|$ into 'faces' similar to the faces of a simple polytope (in fact, in the case of the nerve complex $\mathcal{K} = \mathcal{K}_P$, our faces will be exactly the faces of P). The vertices $i \in \mathcal{K}$ are also vertices of the barycentric subdivision $\mathcal{K}'$, and we set

$$F_i = \operatorname{st}_{\mathcal{K}'}\{i\}, \quad 1 \leqslant i \leqslant m, \tag{4.3}$$

(i.e. F_i is the star of the ith vertex of $\mathcal{K}$ in the barycentric subdivision $\mathcal{K}'$). We refer to F_i as *facets* of our face decomposition, and define a face of codimension k as a nonempty intersection of a set of k facets. In particular, the vertices of our face decomposition are the barycentres of $(n-1)$-dimensional simplices of $\mathcal{K}$. Such a barycentre b corresponds to a maximal simplex $I = \{i_1,\dots,i_n\}$ of $\mathcal{K}$, and we denote by U_I the open subset in $|\operatorname{cone}(\mathcal{K}')|$ obtained by removing all faces not containing b. We observe that any point of $|\operatorname{cone}\mathcal{K}'|$ is contained in U_I for some $I \in \mathcal{K}$.

Under the map $\operatorname{cone}(i_c)\colon |\operatorname{cone}(\mathcal{K}')| \to \mathbb{I}^m$ of Construction 2.9.11 the facet F_i is mapped to the intersection of $\operatorname{cc}(\mathcal{K})$ with the ith coordinate plane $y_i = 0$. Therefore, the image of U_I under the map $\operatorname{cone}(i_c)$ is given by

$$W_I = \operatorname{cone}(i_c)(U_I) = \{(y_1,\dots,y_m) \in \operatorname{cc}(\mathcal{K})\colon y_i \neq 0 \quad \text{for } i \notin I\}. \tag{4.4}$$

Assume now that $\mathcal{K}$ is a triangulated sphere. Then $|\operatorname{cone}\mathcal{K}'|$ is homeomorphic to an n-dimensional disc D^n, and each U_I is homeomorphic to an open subset in $\mathbb{R}^n_\geqslant$ preserving the dimension of faces. (This means that $|\operatorname{cone}\mathcal{K}'|$ is a *manifold with corners*, see Section D.1). By identifying $|\operatorname{cone}\mathcal{K}'|$ with $\operatorname{cc}(\mathcal{K})$ and further identifying $\operatorname{cc}(\mathcal{K})$ with the quotient $\mathcal{Z}_\mathcal{K}/\mathbb{T}^m$, we obtain that each point of $\mathcal{Z}_\mathcal{K}$ has a neighbourhood of the form $\mu^{-1}(W_I)$. It follows from (4.4) that the latter is homeomorphic to an open subset in $\mu_n^{-1}(\mathbb{R}^n_\geqslant) \times T^{m-n} = \mathbb{C}^n \times T^{m-n}$, where $\mathbb{R}^n$ is the coordinate n-plane corresponding to $i_1,\dots,i_n$, the map $\mu_n\colon \mathbb{C}^n \to \mathbb{R}^n_\geqslant$ is the restriction of $\mu\colon \mathbb{C}^m \to \mathbb{R}^m_\geqslant$ to the corresponding coordinate plane in $\mathbb{C}^m$, and the torus T^{m-n} sits in the complementary coordinate $(m-n)$-plane in $\mathbb{C}^m$. An open subset in $\mathbb{C}^n \times T^{m-n}$ with $n \geqslant 1$ can be regarded as an open subset in $\mathbb{R}^{m+n}$, and therefore $\mathcal{Z}_\mathcal{K}$ is an $(m+n)$-dimensional manifold.

If $\mathcal{K}$ is a triangulated manifold, then $|\operatorname{cone}\mathcal{K}'|$ is not a manifold because of the singularity at the cone vertex v. However, by removing this vertex we obtain a (non-compact) manifold whose boundary is $|\mathcal{K}|$. Using the face decomposition defined by (4.3) we obtain that $|\operatorname{cone}\mathcal{K}'| \setminus v$ is locally homeomorphic to $\mathbb{R}^n_\geqslant$ preserving the dimension of faces (i.e. $|\operatorname{cone}\mathcal{K}'| \setminus v$ is a non-compact manifold with corners). Under the identification of $|\operatorname{cone}\mathcal{K}'|$ with $\operatorname{cc}(\mathcal{K})$ the vertex of the cone is mapped to the vertex $(1,\dots,1) \in \mathbb{I}^m$, and $\mu^{-1}(1,\dots,1) = \mathbb{T}^m$. Therefore,

$$\mu^{-1}\bigl(\operatorname{cc}(\mathcal{K})\backslash(1,\dots,1)\bigr) = \mathcal{Z}_\mathcal{K} \setminus \mu^{-1}(1,\dots,1)$$

is an $(m+n)$-dimensional non-compact manifold. □

Remark. A pair of spaces (X,A) where A is a compact subset in X is called a *Lefschetz pair* if $X \setminus A$ is an open (non-compact) manifold. We therefore obtain that $(\mathcal{Z}_\mathcal{K}, \mu^{-1}(1,\dots,1))$ is a Lefschetz pair whenever $\mathcal{K}$ is a triangulated manifold.

We therefore refer to moment-angle complexes $\mathcal{Z}_\mathcal{K}$ corresponding to triangulated spheres as *moment-angle manifolds*. *Polytopal* moment-angle manifolds $\mathcal{Z}_{\mathcal{K}_P}$, corresponding to the nerve complexes $\mathcal{K}_P$ of simple polytopes (see Example 2.2.4), are particularly important. As we shall see in Chapter 6, polytopal moment-angle manifolds are smooth. A smooth structure also exists on moment-angle manifolds $\mathcal{Z}_\mathcal{K}$ corresponding to star-shaped spheres $\mathcal{K}$ (i.e. underlying complexes of complete simplicial fans, see Section 2.5). In general, the smoothness of $\mathcal{Z}_\mathcal{K}$ is open.

The geometry of moment-angle manifolds is nice and rich; it is the subject of Chapter 6.

Real moment-angle complex $\mathcal{R}_\mathcal{K}$. The construction of the moment-angle complex $\mathcal{Z}_\mathcal{K}$ has a real analogue, in which the complex space $\mathbb{C}^m$ is replaced by the real space $\mathbb{R}^m$, the complex polydisc $\mathbb{D}^m$ is replaced by the 'big' cube

$$[-1,1]^m = \bigl\{(u_1,\dots,u_m)\in\mathbb{R}^m\colon |u_i|^2\leqslant 1 \quad \text{for } i=1,\dots,m\bigr\},$$

the standard torus $\mathbb{T}^m$ is replaced by the 'real torus' $\mathbb{Z}_2^m$ (the product of m copies of the group $\mathbb{Z}_2=\{-1,1\}$), and the pair (D^2,S^1) is replaced by (D^1,S^0), where S^0 (a pair of points) is the boundary of the segment D^1. The group $(\mathbb{Z}_2)^m$ acts on the big cube $[-1,1]^m$ coordinatewise, with quotient the standard 'small' cube $\mathbb{I}^m$. The quotient projection $[-1,1]^m\to\mathbb{I}^m$ may be described by the map

$$\rho\colon (u_1,\dots,u_m)\mapsto(u_1^2,\dots,u_m^2).$$

Construction 4.1.5 (real moment-angle complex). Given a simplicial complex $\mathcal{K}$ on $[m]$, define the *real moment-angle complex* $\mathcal{R}_\mathcal{K}$ from the pullback square

$$\begin{tikzcd}
\mathcal{R}_\mathcal{K} \arrow[r, hook] \arrow[d] & {[-1,1]^m} \arrow[d, "\rho"] \\
\operatorname{cc}(\mathcal{K}) \arrow[r, hook] & \mathbb{I}^m
\end{tikzcd}$$

Explicitly, $\mathcal{R}_\mathcal{K}=\rho^{-1}(\operatorname{cc}(\mathcal{K}))$. By construction, $\mathcal{R}_\mathcal{K}$ is a $\mathbb{Z}_2^m$-invariant subspace in the 'big' cube $[-1,1]^m$, and the quotient $\mathcal{R}_\mathcal{K}/\mathbb{Z}_2^m$ is homeomorphic to $|\operatorname{cone}\mathcal{K}|$.

$\mathcal{R}_\mathcal{K}$ is a cubical subcomplex in $[-1,1]^m$ obtained by reflecting the subcomplex $\operatorname{cc}(\mathcal{K})\subset\mathbb{I}^m=[0,1]^m$ at all m coordinate hyperplanes of $\mathbb{R}^m$. If $\mathcal{K}=\mathcal{K}_P$ is the nerve complex of a simple polytope P, then $\operatorname{cc}(\mathcal{K}_P)$ can be viewed as a cubical subdivision of P embedded piecewise linearly into $\mathbb{R}^m_\geqslant$ (see Construction 2.9.7). In this case $\mathcal{R}_{\mathcal{K}_P}$ is obtained by reflecting the image of P at all coordinate planes.

By analogy with (4.2), we have

$$\mathcal{R}_\mathcal{K}=\bigcup_{I\in\mathcal{K}}\Bigl(\prod_{i\in I}D^1\times\prod_{i\notin I}S^0\Bigr).$$

Example 4.1.6.

1. Let $\mathcal{K}=\partial\Delta^{m-1}$ be the boundary of the standard simplex. Then $\mathcal{R}_\mathcal{K}\cong S^{m-1}$ is the boundary of the cube $[-1,1]^m$. For $m=3$ this complex is obtained by reflecting the complex shown in Fig. 2.14 (b) at all 3 coordinate planes of $\mathbb{R}^3$.

2. Let $\mathcal{K}$ consist of m disjoint points. Then $\mathcal{R}_\mathcal{K}$ is the 1-dimensional skeleton (graph) of the cube $[-1,1]^m$. For $m=3$ this complex is obtained by reflecting the complex shown in Fig. 2.14 (a) at all 3 coordinate planes of $\mathbb{R}^3$.

3. More generally, let $\mathcal{K} = \mathrm{sk}^i \Delta^{m-1}$ be the *i-dimensional skeleton* of Δ^{m-1} (i.e. the set of all faces of Δ^{m-1} of dimension $\leqslant i$). Then $\mathcal{R}_\mathcal{K}$ is the $(i+1)$-dimensional skeleton of the cube $[-1,1]^m$.

The following analogue of Theorem 4.1.4 holds, and is proved similarly:

THEOREM 4.1.7. *Let $\mathcal{K}$ be a triangulation of an $(n-1)$-dimensional sphere with m vertices. Then $\mathcal{R}_\mathcal{K}$ is a (closed) topological manifold of dimension n.*

If $\mathcal{K}$ is a triangulated manifold, then $\mathcal{R}_\mathcal{K} \setminus \rho^{-1}(1,\ldots,1)$ is an (open) non-compact manifold, where $\rho^{-1}(1,\ldots,1) = \{-1,1\}^m$.

The real moment-angle complexes corresponding to polygons can be identified easily (compare Exercise 4.1.18):

PROPOSITION 4.1.8. *Let $\mathcal{K}$ be the boundary of an m-gon. Then $\mathcal{R}_\mathcal{K}$ is homeomorphic to an oriented surface S_g of genus $g = 1 + (m-4)2^{m-3}$.*

PROOF. We observe that the manifold $\mathcal{R}_\mathcal{K}$ is orientable (an exercise). Since it is 2-dimensional, its topological type is determined by the Euler characteristic. Now $\mathcal{R}_\mathcal{K}$ is obtained by reflecting the m-gon embedded into $\mathbb{R}^m_\geqslant$ at m coordinate hyperplanes, so that $\mathcal{R}_\mathcal{K}$ is patched from 2^m polygons, meeting by 4 at each vertex. Therefore, the number of vertices is $m2^{m-2}$, the number of edges is $m2^{m-1}$, and the Euler characteristic is

$$\chi(\mathcal{R}_\mathcal{K}) = 2^{m-2}(4-m) = 2 - 2g.$$

The result follows. □

Polyhedral products $(\boldsymbol{X}, \boldsymbol{A})^\mathcal{K}$. Decomposition (4.2) of $\mathcal{Z}_\mathcal{K}$ which uses the disc and circle (D^2, S^1) is readily generalised to arbitrary pairs of spaces:

CONSTRUCTION 4.1.9 (polyhedral product). Let $\mathcal{K}$ be a simplicial complex on $[m]$ and

$$(\boldsymbol{X}, \boldsymbol{A}) = \{(X_1, A_1), \ldots, (X_m, A_m)\}$$

be a set of m pairs of spaces, $A_i \subset X_i$. For each simplex $I \in \mathcal{K}$ we set

$$(\boldsymbol{X}, \boldsymbol{A})^I = \bigl\{(x_1, \ldots, x_m) \in \prod_{j=1}^m X_j \colon\ x_j \in A_j \quad \text{for } j \notin I\bigr\}$$

and define the *polyhedral product* of $(\boldsymbol{X}, \boldsymbol{A})$ corresponding to $\mathcal{K}$ by

$$\mathcal{D}_\mathcal{K}(\boldsymbol{X}, \boldsymbol{A}) = \bigcup_{I \in \mathcal{K}} (\boldsymbol{X}, \boldsymbol{A})^I = \bigcup_{I \in \mathcal{K}} \Bigl(\prod_{i \in I} X_i \times \prod_{i \notin I} A_i\Bigr).$$

Using the categorical language, we can define the diagram

$$\begin{aligned} \mathcal{D}_\mathcal{K}(\boldsymbol{X}, \boldsymbol{A}) \colon \mathrm{CAT}(\mathcal{K}) &\longrightarrow \mathrm{TOP}, \\ I &\longmapsto (\boldsymbol{X}, \boldsymbol{A})^I, \end{aligned}$$

which maps the morphism $I \subset J$ of $\mathrm{CAT}(\mathcal{K})$ to the inclusion of spaces $(\boldsymbol{X}, \boldsymbol{A})^I \subset (\boldsymbol{X}, \boldsymbol{A})^J$. Then we have

$$(\boldsymbol{X}, \boldsymbol{A})^\mathcal{K} = \operatorname{colim} \mathcal{D}_\mathcal{K}(\boldsymbol{X}, \boldsymbol{A}) = \operatorname*{colim}_{I \in \mathcal{K}} (\boldsymbol{X}, \boldsymbol{A})^I.$$

In the case when all the pairs (X_i, A_i) are the same, i.e. $X_i = X$ and $A_i = A$ for $i = 1, \ldots, m$, we use the notation $(X, A)^\mathcal{K}$ for $(\boldsymbol{X}, \boldsymbol{A})^\mathcal{K}$. Also, if each X_i is a pointed space and $A_i = pt$, then we use the abbreviated notation $(\boldsymbol{X})^\mathcal{K}$ for $(\boldsymbol{X}, pt)^\mathcal{K}$, and $X^\mathcal{K}$ for $(X, pt)^\mathcal{K}$.

Remark. The decomposition of $\mathcal{Z}_\mathcal{K}$ into a union of products of discs and circles first appeared in [**42**] (in the polytopal case) and in [**44**] (in general). The term 'moment-angle complex' for $\mathcal{Z}_\mathcal{K} = (D^2, S^1)^\mathcal{K}$ was also introduced in [**44**], where several other examples of polyhedral products $(X, A)^\mathcal{K}$ were considered. The definition of $(X, A)^\mathcal{K}$ for an arbitrary pair of spaces (X, A) was suggested to the authors by N. Strickland (in a private communication, and also in an unpublished note) as a general framework for the constructions of [**44**]; it was also included in the final version of [**44**] and in [**45**]. Further generalisations of $(X, A)^\mathcal{K}$ to a set of pairs of spaces $(\boldsymbol{X}, \boldsymbol{A})$ were studied in the work of Grbić and Theriault [**104**], as well as Bahri, Bendersky, Cohen and Gitler [**11**], where the term 'polyhedral product' was introduced (following a suggestion of W. Browder). Since 2000, the terms 'generalised moment-angle complex', '$\mathcal{K}$-product' and 'partial product space' have been also used to refer to the spaces $(X, A)^\mathcal{K}$.

Recall that a *map of pairs* $(X, A) \to (X', A')$ is a commutative diagram

$$\begin{tikzcd} A \arrow[r, hook] \arrow[d] & X \arrow[d] \\ A' \arrow[r, hook] & X'. \end{tikzcd} \tag{4.5}$$

We refer to (X, A) as a *monoid pair* if X is a topological monoid (a space with a continuous associative multiplication and unit), and A is a submonoid. A *map of monoid pairs* is a map of pairs in which all maps in (4.5) are homomorphisms.

If (X, A) is a monoid pair, then a set map $\varphi\colon [l] \to [m]$ induces a map

$$\psi\colon \prod_{i=1}^{l} X \to \prod_{i=1}^{m} X, \quad (x_1, \ldots, x_l) \mapsto (y_1, \ldots, y_m), \tag{4.6}$$

where

$$y_j = \prod_{i \in \varphi^{-1}(j)} x_i, \qquad \text{for } j = 1, \ldots, m,$$

and we set $y_j = 1$ if $\varphi^{-1}(j) = \varnothing$.

Proposition 4.1.10. *If (X, A) is a monoid pair, then $(X, A)^\mathcal{K}$ is an invariant subspace of $\prod_{i=1}^m X$ with respect to the coordinatewise action of $\prod_{i=1}^m A$ on $\prod_{i=1}^m X$.*

Proof. Indeed, $(X, A)^I \subset \prod_{i=1}^m X$ is an invariant subset for each $I \in \mathcal{K}$. □

The following proposition describes the functorial properties of the polyhedral product in its $\mathcal{K}$ and $(\boldsymbol{X}, \boldsymbol{A})$ arguments.

Proposition 4.1.11.

(a) *A set of maps of pairs $(\boldsymbol{X}, \boldsymbol{A}) \to (\boldsymbol{X}', \boldsymbol{A}')$ induces a map of polyhedral products $(\boldsymbol{X}, \boldsymbol{A})^\mathcal{K} \to (\boldsymbol{X}', \boldsymbol{A}')^\mathcal{K}$. If two sets of maps $(\boldsymbol{X}, \boldsymbol{A}) \to (\boldsymbol{X}', \boldsymbol{A}')$ are componentwise homotopic, then the induced maps $(\boldsymbol{X}, \boldsymbol{A})^\mathcal{K} \to (\boldsymbol{X}', \boldsymbol{A}')^\mathcal{K}$ are also homotopic.*

(b) *An inclusion of a simplicial subcomplex $\mathcal{L} \hookrightarrow \mathcal{K}$ induces an inclusion of polyhedral products $(\boldsymbol{X}, \boldsymbol{A})^\mathcal{L} \hookrightarrow (\boldsymbol{X}, \boldsymbol{A})^\mathcal{K}$.*

(c) *If (X, A) is a monoid pair, then for any simplicial map $\varphi\colon \mathcal{L} \to \mathcal{K}$ of simplicial complexes on the sets $[l]$ and $[m]$ respectively, the map (4.6) restricts to a map of polyhedral products $\varphi_\mathcal{Z}\colon (X, A)^\mathcal{L} \to (X, A)^\mathcal{K}$.*

(d) *If* (X,A) *is a commutative monoid pair, then the restriction*

$$\psi|_A\colon \prod_{i=1}^{l} A \to \prod_{i=1}^{m} A$$

of (4.6) *is a homomorphism, and the induced map* $\varphi_{\mathcal Z}\colon (X,A)^{\mathcal L} \to (X,A)^{\mathcal K}$ *is* $\psi|_A$*-equivariant, i.e.*

$$\varphi_{\mathcal Z}(\boldsymbol a\cdot \boldsymbol x)=\psi|_A(\boldsymbol a)\cdot\varphi_{\mathcal Z}(\boldsymbol x)$$

for all $\boldsymbol a=(a_1,\ldots,a_l)\in\prod_{i=1}^{l}A$ *and* $\boldsymbol x=(x_1,\ldots,x_l)\in(X,A)^{\mathcal L}$.

PROOF. For (a), we observe that a set of maps $(\boldsymbol X,\boldsymbol A)\to(\boldsymbol X',\boldsymbol A')$ induces a map $(\boldsymbol X,\boldsymbol A)^I\to(\boldsymbol X',\boldsymbol A')^I$ for each $I\in\mathcal K$, and these maps corresponding to different $I,J\in\mathcal K$ are compatible on the intersection $(\boldsymbol X,\boldsymbol A)^I\cap(\boldsymbol X,\boldsymbol A)^J=(\boldsymbol X,\boldsymbol A)^{I\cap J}$. We therefore obtain a map $(\boldsymbol X,\boldsymbol A)^{\mathcal K}\to(\boldsymbol X',\boldsymbol A')^{\mathcal K}$. A componentwise homotopy between two maps $(\boldsymbol X,\boldsymbol A)\to(\boldsymbol X',\boldsymbol A')$ can be thought of as a map of pairs $(\boldsymbol X\times\mathbb I,\boldsymbol A\times\mathbb I)\to(\boldsymbol X',\boldsymbol A')$, where $\boldsymbol X\times\mathbb I$ consists of spaces $X_i\times\mathbb I$. It therefore induces a map of polyhedral products

$$(\boldsymbol X\times\mathbb I,\boldsymbol A\times\mathbb I)^{\mathcal K}\to(\boldsymbol X',\boldsymbol A')^{\mathcal K}$$

where $(\boldsymbol X\times\mathbb I,\boldsymbol A\times\mathbb I)^{\mathcal K}\cong(\boldsymbol X,\boldsymbol A)^{\mathcal K}\times(\mathbb I,\mathbb I)^{\mathcal K}=(\boldsymbol X,\boldsymbol A)^{\mathcal K}\times\mathbb I^m$. By restricting the resulting map $(\boldsymbol X,\boldsymbol A)^{\mathcal K}\times\mathbb I^m\to(\boldsymbol X',\boldsymbol A')^{\mathcal K}$ to the diagonal of the cube $\mathbb I^m$ we obtain a homotopy between the two induced maps $(\boldsymbol X,\boldsymbol A)^{\mathcal K}\to(\boldsymbol X',\boldsymbol A')^{\mathcal K}$.

To prove (b) we just observe that if $\mathcal L$ is a subcomplex of $\mathcal K$, then for each $I\in\mathcal L$ we have $(\boldsymbol X,\boldsymbol A)^I\subset\mathcal Z_{\mathcal K}$.

To prove (c) we observe that for any subset $I\subset[m]$ we have $\psi\bigl((X,A)^I\bigr)\subset(X,A)^{\varphi(I)}$. Let $I\in\mathcal L$, so that $(X,A)^I\subset(X,A)^{\mathcal L}$. Since φ is a simplicial map, we have $\varphi(I)\in\mathcal K$ and $(X,A)^{\varphi(I)}\subset(X,A)^{\mathcal K}$. Therefore, the map ψ restricts to a map of polyhedral products $(X,A)^{\mathcal L}\to(X,A)^{\mathcal K}$.

Statement (d) is proved by a direct calculation:

$$\begin{aligned}
\varphi_{\mathcal Z}(\boldsymbol a\cdot\boldsymbol x)=\varphi_{\mathcal Z}(a_1x_1,\ldots,a_lx_l)&=\Bigl(\prod_{i\in\varphi^{-1}(1)}a_ix_i,\ldots,\prod_{i\in\varphi^{-1}(m)}a_ix_i\Bigr)\\
&=\Bigl(\prod_{i\in\varphi^{-1}(1)}a_i\prod_{i\in\varphi^{-1}(1)}x_i,\ldots,\prod_{i\in\varphi^{-1}(m)}a_i\prod_{i\in\varphi^{-1}(m)}x_i\Bigr)\\
&=\Bigl(\prod_{i\in\varphi^{-1}(1)}a_i,\ldots,\prod_{i\in\varphi^{-1}(m)}a_i\Bigr)\cdot\Bigl(\prod_{i\in\varphi^{-1}(1)}x_i,\ldots,\prod_{i\in\varphi^{-1}(m)}x_i\Bigr)\\
&=\psi|_A(\boldsymbol a)\cdot\varphi_{\mathcal Z}(\boldsymbol x).
\end{aligned}$$

Note that we have used the commutativity of X in the third identity. □

We state the most important particular case of Proposition 4.1.11 separately:

PROPOSITION 4.1.12. *A simplicial map* $\mathcal L\to\mathcal K$ *of simplicial complexes on the sets* $[l]$ *and* $[m]$ *gives rise to a map of moment-angle complexes* $\mathcal Z_{\mathcal L}\to\mathcal Z_{\mathcal K}$ *which is equivariant with respect to the induced homomorphism of tori* $\mathbb T^l\to\mathbb T^m$.

The polyhedral product construction behaves nicely with respect to the join operation (the proof is exactly the same as that of Proposition 4.1.3):

Proposition 4.1.13. *We have that*

$$(\boldsymbol{X},\boldsymbol{A})^{\mathcal{K}_1*\mathcal{K}_2}=(\boldsymbol{X},\boldsymbol{A})^{\mathcal{K}_1}\times(\boldsymbol{X},\boldsymbol{A})^{\mathcal{K}_2}.$$

Example 4.1.14.

1. The moment-angle complex $\mathcal{Z}_{\mathcal{K}}$ is the polyhedral product $(D^2,S^1)^{\mathcal{K}}$ (when considered abstractly) or $(\mathbb{D},\mathbb{S})^{\mathcal{K}}$ (when viewed as a subcomplex in $\mathbb{D}^m$).

2. For the cubical complex of (2.13) we have

$$\operatorname{cc}(\mathcal{K})=(I^1,1)^{\mathcal{K}},$$

where $I^1=[0,1]$ is the unit segment and 1 is its edge. The quotient map $\mathcal{Z}_{\mathcal{K}}\to|\operatorname{cone}\mathcal{K}|$ is the map of polyhedral products $(D^2,S^1)^{\mathcal{K}}\to(I^1,1)^{\mathcal{K}}$ induced by the map of pairs $(D^2,S^1)\to(I^1,1)$, which is the quotient map by the S^1-action.

3. For the real moment-angle complex we have

$$\mathcal{R}_{\mathcal{K}}=(D^1,S^0)^{\mathcal{K}},$$

Where $D^1=[-1,1]$ is a 1-disc, and $S^0=\{-1,1\}$ is its boundary.

4. If $\mathcal{K}$ consists of m disjoint points and $A_i=pt$, then

$$(\boldsymbol{X},pt)^{\mathcal{K}}=\boldsymbol{X}^{\mathcal{K}}=X_1\vee X_2\vee\cdots\vee X_m$$

is the *wedge* (or *bouquet*) of the X_i's.

5. More generally, consider the sequence of inclusions of skeleta

$$\operatorname{sk}^0\Delta^{m-1}\subset\cdots\subset\operatorname{sk}^{m-2}\Delta^{m-1}\subset\operatorname{sk}^{m-1}\Delta^{m-1}=\Delta^{m-1}.$$

It gives rise to a filtration in the product $X_1\times X_2\times\cdots\times X_m$:

$$\begin{tikzcd}
\boldsymbol{X}^{\operatorname{sk}^0\Delta^{m-1}} \arrow[r, phantom, "\subset"] \arrow[d, equal] & \cdots \arrow[r, phantom, "\subset"] & \boldsymbol{X}^{\operatorname{sk}^{m-2}\Delta^{m-1}} \arrow[r, phantom, "\subset"] \arrow[d, equal] & \boldsymbol{X}^{\operatorname{sk}^{m-1}\Delta^{m-1}} \arrow[d, equal] \\
X_1\vee X_2\vee\cdots\vee X_m & & \boldsymbol{X}^{\partial\Delta^{m-1}} & X_1\times X_2\times\cdots\times X_m
\end{tikzcd}$$

Its second-to-last term, $\boldsymbol{X}^{\partial\Delta^{m-1}}$ is known to topologists as the *fat wedge* of the X_i's. Explicitly, the fat wedge of a sequence of pointed spaces $X_1,\ldots,X_m$ is

$$(X_1\times\cdots\times X_{m-1}\times pt)\cup(X_1\times\cdots\times pt\times X_m)\cup\cdots\cup(pt\times\cdots\times X_{m-1}\times X_m),$$

where the union is taken inside the product $X_1\times X_2\times\cdots\times X_m$.

The filtration above was considered by G. Porter **[186]**, who obtained a decomposition of its loop spaces in the case when each X_i is a suspension, generalising the *Hilton–Milnor Theorem*. We shall consider this decomposition in more detail in Chapter 8.

Exercises.

4.1.15. Show that if $\mathcal{K}$ is a triangulated sphere, then the manifold $\mathcal{R}_{\mathcal{K}}$ is orientable. (Hint: use the fact that $\mathcal{R}_{\mathcal{K}}$ is obtained by reflecting the n-ball $|\operatorname{cone}\mathcal{K}|$ at all m coordinate hyperplanes of $\mathbb{R}^m$ to extend the orientation from $|\operatorname{cone}\mathcal{K}|$ to the whole of $\mathcal{R}_{\mathcal{K}}$.)

4.1.16. Show that if $\mathcal{K}=\operatorname{sk}^i\Delta^{m-1}$, then $\mathcal{R}_{\mathcal{K}}$ is homotopy equivalent to a wedge of $(i+1)$-dimensional spheres (see Example 4.1.6.3). The number of spheres is given by $\sum_{k=i+2}^{m}\binom{m}{k}\binom{k-1}{i+1}$.

4.1.17. Let $\mathcal{K}$ be the complex consisting of three disjoint points. Show that $\mathcal{Z}_{\mathcal{K}}$ is homotopy equivalent to the following wedge (bouquet) of spheres:

$$\mathcal{Z}_{\mathcal{K}} \cong S^3 \vee S^3 \vee S^3 \vee S^4 \vee S^4.$$

(Hint: compare the case $m = 3, i = 0$ of the previous exercise; it may also help to look at the realisation of $\mathcal{Z}_{\mathcal{K}}$ as the complement of a coordinate subspace arrangement (up to homotopy), see Section 4.6.)

4.1.18. Let $\mathcal{K}$ be the boundary of a 5-gon. Show that the manifold $\mathcal{Z}_{\mathcal{K}}$ is homeomorphic to $(S^3 \times S^4)^{\# 5}$, a connected sum of 5 copies of $S^3 \times S^4$. (This may be a difficult one; see Theorem 4.5.12 for the general statement.)

4.1.19. If $\mathcal{K}$ is a triangulation of an $(n-1)$-sphere with m vertices, then for any $k > 0$ the polyhedral product $(D^k, S^{k-1})^{\mathcal{K}}$ is a manifold of dimension $m(k-1)+n$.

4.1.20. Let $\mathcal{K}_I$ be the full subcomplex corresponding to $I \subset [m]$. Then $\mathcal{Z}_{\mathcal{K}_I}$ is a retract of $\mathcal{Z}_{\mathcal{K}}$.

4.2. Basic homotopical properties

Two key observations of this section constitute the basis for the subsequent applications of the commutative algebra apparatus of Chapter 3 to toric topology. First, the cohomology of the polyhedral product space of the form $(\mathbb{C}P^\infty)^{\mathcal{K}} = (\mathbb{C}P^\infty, pt)^{\mathcal{K}}$ is isomorphic to the face ring $\mathbb{Z}[\mathcal{K}]$ (Proposition 4.2.1). Second, the moment-angle complex $\mathcal{Z}_{\mathcal{K}} = (D^2, S^1)^{\mathcal{K}}$ is the homotopy fibre of the canonical inclusion $(\mathbb{C}P^\infty)^{\mathcal{K}} \hookrightarrow (\mathbb{C}P^\infty, \mathbb{C}P^\infty)^{\mathcal{K}} = (\mathbb{C}P^\infty)^m$ (Theorem 4.2.2).

The classifying space BS^1 of the circle S^1 is the infinite-dimensional complex projective space $\mathbb{C}P^\infty$. The classifying space BT^m of the m-torus T^m is the product $(\mathbb{C}P^\infty)^m$ of m copies of $\mathbb{C}P^\infty$. The universal principal S^1-bundle is the infinite Hopf bundle $S^\infty \to \mathbb{C}P^\infty$ (the direct limit of Hopf bundles $S^{2k+1} \to \mathbb{C}P^k$), so the total space ET^m of the universal principal T^m-bundle over BT^m can be identified with the m-fold product of the infinite-dimensional sphere S^∞.

The integral cohomology ring of BT^m is isomorphic to the polynomial ring $\mathbb{Z}[v_1, \ldots, v_m]$, $\deg v_i = 2$ (this explains our choice of grading). The space BT^m has the canonical cell decomposition, in which each factor $\mathbb{C}P^\infty$ has one cell in every even dimension. The polyhedral product

$$(\mathbb{C}P^\infty)^{\mathcal{K}} = \bigcup_{I \in \mathcal{K}} (\mathbb{C}P^\infty, pt)^I$$

is a cellular subcomplex in $BT^m = (\mathbb{C}P^\infty)^m$.

Proposition 4.2.1. *The cohomology ring of* $(\mathbb{C}P^\infty)^{\mathcal{K}}$ *is isomorphic to the face ring* $\mathbb{Z}[\mathcal{K}]$*. The inclusion of a cellular subcomplex*

$$i\colon (\mathbb{C}P^\infty)^{\mathcal{K}} \hookrightarrow (\mathbb{C}P^\infty)^m$$

induces the quotient projection in cohomology:

$$i^*\colon \mathbb{Z}[v_1, \ldots, v_m] \to \mathbb{Z}[v_1, \ldots, v_m]/\mathcal{I}_{\mathcal{K}} = \mathbb{Z}[\mathcal{K}].$$

Proof. Since $(\mathbb{C}P^\infty)^m$ has only even-dimensional cells and $(\mathbb{C}P^\infty)^{\mathcal{K}}$ is a cellular subcomplex, the cohomology of both spaces coincides with their cellular cochains. Let D_j^{2k} denote the $2k$-dimensional cell in the jth factor of $(\mathbb{C}P^\infty)^m$.

The cellular cochain group $\mathcal{C}^*((\mathbb{C}P^\infty)^m)$ has basis of cochains $\bigl(D^{2k_1}_{j_1}\cdots D^{2k_p}_{j_p}\bigr)^*$ dual to the products of cells $D^{2k_1}_{j_1}\times\cdots\times D^{2k_p}_{j_p}$. The cochain map

$$\mathcal{C}^*\bigl((\mathbb{C}P^\infty)^m\bigr)\to\mathcal{C}^*\bigl((\mathbb{C}P^\infty)^\mathcal{K}\bigr)$$

induced by the inclusion $(\mathbb{C}P^\infty)^\mathcal{K}\hookrightarrow(\mathbb{C}P^\infty)^m$ is an epimorphism with kernel generated by those cochains $\bigl(D^{2k_1}_{j_1}\cdots D^{2k_p}_{j_p}\bigr)^*$ for which $\{j_1,\dots,j_p\}\notin\mathcal{K}$. Under the identification of $\mathcal{C}^*((\mathbb{C}P^\infty)^m)$ with $\mathbb{Z}[v_1,\dots,v_m]$, a cochain $\bigl(D^{2k_1}_{j_1}\cdots D^{2k_p}_{j_p}\bigr)^*$ is mapped to the monomial $v^{k_1}_{j_1}\cdots v^{k_p}_{j_p}$. Therefore, $\mathcal{C}^*((\mathbb{C}P^\infty)^\mathcal{K})$ is identified with the quotient of $\mathbb{Z}[v_1,\dots,v_m]$ by the subgroup generated by all monomials $v^{k_1}_{j_1}\cdots v^{k_p}_{j_p}$ with $\{j_1,\dots,j_p\}\notin\mathcal{K}$. By Proposition 3.1.9, this quotient is exactly $\mathbb{Z}[\mathcal{K}]$. □

We consider the Borel construction $ET^m\times_{T^m}\mathcal{Z}_\mathcal{K}$ for the T^m-space $\mathcal{Z}_\mathcal{K}$ (see Appendix B.3).

THEOREM 4.2.2. *The inclusion* $i\colon(\mathbb{C}P^\infty)^\mathcal{K}\hookrightarrow(\mathbb{C}P^\infty)^m$ *is decomposed into a composition of a homotopy equivalence*

$$h\colon(\mathbb{C}P^\infty)^\mathcal{K}\xrightarrow{\simeq}ET^m\times_{T^m}\mathcal{Z}_\mathcal{K}$$

and the fibration $p\colon ET^m\times_{T^m}\mathcal{Z}_\mathcal{K}\to BT^m=(\mathbb{C}P^\infty)^m$ *with fibre* $\mathcal{Z}_\mathcal{K}$.

In particular, the moment-angle complex $\mathcal{Z}_\mathcal{K}$ *is the homotopy fibre of the canonical inclusion* $i\colon(\mathbb{C}P^\infty)^\mathcal{K}\hookrightarrow(\mathbb{C}P^\infty)^m$.

PROOF. We use functoriality and homotopy invariance of the polyhedral product construction (Proposition 4.1.11). We have $\mathcal{Z}_\mathcal{K}=\bigcup_{I\in\mathcal{K}}(D^2,S^1)^I$, see (4.1), and each subset $(D^2,S^1)^I$ is T^m-invariant. We therefore obtain the following decomposition of the Borel construction as a polyhedral product:

$$\begin{aligned}ET^m\times_{T^m}\mathcal{Z}_\mathcal{K}=\bigcup_{I\in\mathcal{K}}\bigl(ET^m\times_{T^m}(D^2,S^1)^I\bigr)&=\bigcup_{I\in\mathcal{K}}\bigl(S^\infty\times_{S^1}D^2,S^\infty\times_{S^1}S^1\bigr)^I\\&=\bigl(S^\infty\times_{S^1}D^2,S^\infty\times_{S^1}S^1\bigr)^\mathcal{K},\end{aligned}$$

where $S^\infty=ET^1=ES^1$.

Now consider the commutative diagram

$$\begin{tikzcd} pt \arrow[r] \arrow[d] & S^\infty\times_{S^1}S^1 \arrow[r] \arrow[d] & pt \arrow[d] \\ \mathbb{C}P^\infty \arrow[r, "j"] & S^\infty\times_{S^1}D^2 \arrow[r, "f"] & \mathbb{C}P^\infty, \end{tikzcd} \tag{4.7}$$

where j is the inclusion of the zero section in a disc bundle (note that $\mathbb{C}P^\infty=S^\infty/S^1$), and f is the projection map from the bundle to its Thom space,

$$f\colon S^\infty\times_{S^1}D^2\to(S^\infty\times_{S^1}D^2)/(S^\infty\times_{S^1}S^1)\cong\mathbb{C}P^\infty.$$

Since $S^\infty\times_{S^1}S^1=S^\infty$ and D^2 are contractible, the composite maps $f\circ j$ and $j\circ f$ are homotopic to the identity. It follows that we have a homotopy equivalence of pairs $(\mathbb{C}P^\infty,pt)\to(S^\infty\times_{S^1}D^2,S^\infty\times_{S^1}S^1)$ which induces a homotopy equivalence of polyhedral products

$$h\colon(\mathbb{C}P^\infty)^\mathcal{K}\to(S^\infty\times_{S^1}D^2,S^\infty\times_{S^1}S^1)^\mathcal{K}=ET^m\times_{T^m}\mathcal{Z}_\mathcal{K}.$$

In order to establish the factorisation $i = p \circ h$ we consider the diagram

$$\begin{tikzcd}
pt \arrow[r] \arrow[d] & S^\infty \times_{S^1} S^1 \arrow[r] \arrow[d] & \mathbb{C}P^\infty \arrow[d, equal] \\
\mathbb{C}P^\infty \arrow[r, "j"] & S^\infty \times_{S^1} D^2 \arrow[r, "g"] & \mathbb{C}P^\infty,
\end{tikzcd}$$

where g is the projection of the disc bundle onto its base (note that his map is different from the map f above). By passing to the induced maps of polyhedral products we obtain the factorisation of

$$i\colon (\mathbb{C}P^\infty)^{\mathcal{K}} \xrightarrow{h} (S^\infty \times_{S^1} D^2, S^\infty \times_{S^1} S^1)^{\mathcal{K}} \xrightarrow{p} (\mathbb{C}P^\infty, \mathbb{C}P^\infty)^{\mathcal{K}}$$

into the composition of h and p. □

The following statement is originally due to Davis and Januszkiewicz [**70**, Theorem 4.8] (they used a different model for $\mathcal{Z}_{\mathcal{K}}$, which will be discussed in Section 6.2).

COROLLARY 4.2.3. *The equivariant cohomology ring of the moment-angle complex $\mathcal{Z}_{\mathcal{K}}$ is isomorphic to the face ring of $\mathcal{K}$:*

$$H^*_{T^m}(\mathcal{Z}_{\mathcal{K}}) \cong \mathbb{Z}[\mathcal{K}].$$

In equivariant cohomology, the projection $p\colon ET^m \times_{T^m} \mathcal{Z}_{\mathcal{K}} \to BT^m$ induces the quotient projection

$$p^*\colon \mathbb{Z}[v_1, \ldots, v_m] \to \mathbb{Z}[\mathcal{K}] = \mathbb{Z}[v_1, \ldots, v_m]/\mathcal{I}_{\mathcal{K}}.$$

PROOF. This follows from Theorem 4.2.2 and Proposition 4.2.1. □

We have the following basic information about the homotopy groups of $\mathcal{Z}_{\mathcal{K}}$:

PROPOSITION 4.2.4.

(a) *If $\mathcal{K}$ is a simplicial complex on the vertex set $[m]$ (i.e. there are no ghost vertices), then the moment-angle complex $\mathcal{Z}_{\mathcal{K}}$ is 2-connected (i.e. $\pi_1(\mathcal{Z}_{\mathcal{K}}) = \pi_2(\mathcal{Z}_{\mathcal{K}}) = 0$), and*

$$\pi_i(\mathcal{Z}_{\mathcal{K}}) = \pi_i\bigl((\mathbb{C}P^\infty)^{\mathcal{K}}\bigr) \quad \text{for } i \geqslant 3.$$

(b) *If $\mathcal{K}$ is a q-neighbourly simplicial complex, then $\pi_i(\mathcal{Z}_{\mathcal{K}}) = 0$ for $i < 2q+1$. Furthermore, $\pi_{2q+1}(\mathcal{Z}_{\mathcal{K}})$ is a free abelian group of rank equal to the number of $(q+1)$-element missing faces of $\mathcal{K}$.*

PROOF. We observe that $(\mathbb{C}P^\infty)^m$ is the Eilenberg–Mac Lane space $K(\mathbb{Z}^m, 2)$, and the 3-dimensional skeleton of $(\mathbb{C}P^\infty)^{\mathcal{K}}$ coincides with the 3-skeleton of $(\mathbb{C}P^\infty)^m$. If $\mathcal{K}$ is q-neighbourly, then the $(2q+1)$-skeletons of $(\mathbb{C}P^\infty)^{\mathcal{K}}$ and $(\mathbb{C}P^\infty)^m$ agree. Now both statements follow from the exact homotopy sequence of the map $(\mathbb{C}P^\infty)^{\mathcal{K}} \to (\mathbb{C}P^\infty)^m$ with homotopy fibre $\mathcal{Z}_{\mathcal{K}}$. □

EXAMPLE 4.2.5. Let $\mathcal{K} = \mathrm{sk}_1\Delta^3$ (a complete graph on 4 vertices). Then $\mathcal{K}$ is 2-neighbourly and has 4 missing triangles, so $\mathcal{Z}_{\mathcal{K}}$ is 4-connected and $\pi_5(\mathcal{Z}_{\mathcal{K}}) = \mathbb{Z}^4$.

4.3. Cell decomposition

We consider the following decomposition of the disc $\mathbb{D}$ into 3 cells: the point $1 \in \mathbb{D}$ is the 0-cell; the complement to 1 in the boundary circle is the 1-cell, which we denote by T; and the interior of $\mathbb{D}$ is the 2-cell, which we denote by D. By taking product we obtain a cellular decomposition of $\mathbb{D}^m$ whose cells are parametrised by pairs of subsets $J, I \subset [m]$ with $J \cap I = \varnothing$: the set J parametrises the T-cells in the product and I parametrises the D-cells. We denote the cell of $\mathbb{D}^m$ corresponding to a pair J, I by $\varkappa(J,I)$; it is a product of $|J|$ cells of T-type and $|I|$ cells of D-type (the positions in $[m] \setminus I \cup J$ are filled by 0-cells). Then $\mathcal{Z}_\mathcal{K}$ embeds as a cellular subcomplex in $\mathbb{D}^m$; we have $\varkappa(J,I) \subset \mathcal{Z}_\mathcal{K}$ whenever $I \in \mathcal{K}$.



Let $\mathcal{C}^*(\mathcal{Z}_\mathcal{K})$ be the cellular cochains of $\mathcal{Z}_\mathcal{K}$. It has a basis of cochains $\varkappa(J,I)^*$ dual to the corresponding cells. We introduce the bigrading by setting

$$\operatorname{bideg} \varkappa(J,I)^* = (-|J|, 2|I| + 2|J|),$$

so that $\operatorname{bideg} D = (0,2)$, $\operatorname{bideg} T = (-1,2)$ and $\operatorname{bideg} 1 = (0,0)$. Since the cellular differential preserves the second grading, the complex $\mathcal{C}^*(\mathcal{Z}_\mathcal{K})$ splits into the sum of its components with fixed second degree:

$$\mathcal{C}^*(\mathcal{Z}_\mathcal{K}) = \bigoplus_{q=0}^{m} \mathcal{C}^{*,2q}(\mathcal{Z}_\mathcal{K}). \tag{4.8}$$

The cohomology of the moment-angle complex therefore acquires an additional grading, and we define the *bigraded Betti numbers* of $\mathcal{Z}_\mathcal{K}$ by

$$b^{-p,2q}(\mathcal{Z}_\mathcal{K}) = \operatorname{rank} H^{-p,2q}(\mathcal{Z}_\mathcal{K}), \quad \text{for } 1 \leqslant p, q \leqslant m. \tag{4.9}$$

The ordinary Betti numbers of $\mathcal{Z}_\mathcal{K}$ therefore satisfy

$$b^k(\mathcal{Z}_\mathcal{K}) = \sum_{-p+2q=k} b^{-p,2q}(\mathcal{Z}_\mathcal{K}). \tag{4.10}$$

The map of the moment-angle complexes $\mathcal{Z}_\mathcal{K} \to \mathcal{Z}_\mathcal{L}$ induced by a simplicial map $\mathcal{K} \to \mathcal{L}$ (see Proposition 4.1.12) is clearly a cellular map. We therefore obtain

PROPOSITION 4.3.1. *The correspondence $\mathcal{K} \mapsto \mathcal{Z}_\mathcal{K}$ gives rise to a functor from the category of simplicial complexes and simplicial maps to the category of cell complexes with torus actions and equivariant maps. It also induces a natural transformation between the functor of simplicial cochains of $\mathcal{K}$ and the functor of cellular cochains of $\mathcal{Z}_\mathcal{K}$.*

The map $\mathcal{Z}_\mathcal{K} \to \mathcal{Z}_\mathcal{L}$ induced by a simplicial map $\mathcal{K} \to \mathcal{L}$ preserves the cellular bigrading, so that the bigraded cohomology groups are also functorial.

4.4. Cohomology ring

The main result of this section, Theorem 4.4.4, establishes an isomorphism between the integral cohomology ring of the moment-angle complex $\mathcal{Z}_\mathcal{K}$ and the Tor-algebra of the simplicial complex $\mathcal{K}$. This result was first proved in [**43**] for field coefficients using the Eilenberg–Moore spectral sequence of the fibration $\mathcal{Z}_\mathcal{K} \to (\mathbb{C}P^\infty)^\mathcal{K} \to (\mathbb{C}P^\infty)^m$. The proof given here is taken from [**20**] and [**47**]; it establishes the isomorphism over the integers and works with the cellular cochains.

One of the key steps in the proof is the construction of a cellular approximation of the diagonal map $\varDelta\colon \mathcal{Z}_{\mathcal{K}} \to \mathcal{Z}_{\mathcal{K}} \times \mathcal{Z}_{\mathcal{K}}$ which is functorial with respect to maps of moment-angle complexes induced by simplicial maps. The resulting cellular cochain algebra is isomorphic to the algebra $R^*(\mathcal{K})$ from Construction 3.2.5 (obtained by factorising the Koszul algebra of the face ring $\mathbb{Z}[\mathcal{K}]$ by an acyclic ideal); its cohomology is isomorphic to the Tor-algebra of $\mathcal{K}$.

Another proof of Theorem 4.4.4 was given by Franz [**93**].

Algebraic model for cellular cochains. We recall the algebra $R^*(\mathcal{K})$ from Construction 3.2.5:

$$R^*(\mathcal{K}) = \Lambda[u_1, \dots, u_m] \otimes \mathbb{Z}[\mathcal{K}]/(v_i^2 = u_i v_i = 0,\ 1 \leqslant i \leqslant m),$$

with the bigrading and differential given by

$$\operatorname{bideg} u_i = (-1, 2), \quad \operatorname{bideg} v_i = (0, 2), \quad du_i = v_i, \quad dv_i = 0.$$

The algebra $R^*(\mathcal{K})$ has finite rank as an abelian group, with a basis of monomials $u_J v_I$ where $J \subset [m]$, $I \in \mathcal{K}$ and $J \cap I = \varnothing$.

Comparing the differential graded module structures in $R^*(\mathcal{K})$ and $\mathcal{C}^*(\mathcal{Z}_{\mathcal{K}})$ we observe that they coincide, as described in the following statement:

LEMMA 4.4.1. *The map*

$$g\colon R^*(\mathcal{K}) \to \mathcal{C}^*(\mathcal{Z}_{\mathcal{K}}), \quad u_J v_I \mapsto \varkappa(J, I)^*,$$

is an isomorphism of cochain complexes. Hence, there is an additive isomorphism

$$H[R^*(\mathcal{K})] \cong H^*(\mathcal{Z}_{\mathcal{K}}).$$

PROOF. Since g arises from a bijective correspondence between bases of $R^*(\mathcal{K})$ and $\mathcal{C}^*(\mathcal{Z}_{\mathcal{K}})$, it is an isomorphism of bigraded modules (or groups). It also clearly commutes with the differentials:

$$\delta g(u_i) = \delta(T_i^*) = D_i^* = g(v_i) = g(du_i), \quad \delta g(v_i) = \delta(D_i^*) = 0 = g(dv_i),$$

where T_i denotes the cell $\varkappa(\{i\}, \varnothing)$, and $D_i = \varkappa(\varnothing, \{i\})$. □

Having identified the algebra $R^*(\mathcal{K})$ with the cellular cochains of the moment-angle complex, we can give a topological interpretation to the quasi-isomorphism of Lemma 3.2.6. To do this we shall identify the Koszul algebra $\Lambda[u_1, \dots, u_m] \otimes \mathbb{Z}[\mathcal{K}]$ with the cellular cochains of a space homotopy equivalent to $\mathcal{Z}_{\mathcal{K}}$.

The infinite-dimensional sphere S^∞ is the direct limit (union) of standardly embedded odd-dimensional spheres. Each odd sphere S^{2k+1} can be obtained from S^{2k-1} by attaching two cells of dimensions $2k$ and $2k+1$:

$$S^{2k+1} \cong (S^{2k-1} \cup_f D^{2k}) \cup_g D^{2k+1}.$$

Here the map $f\colon \partial D^{2k} \to S^{2k-1}$ is the identity (and has degree 1), and the map $g\colon \partial D^{2k+1} = S^{2k} \to D^{2k}$ is the projection of the standard sphere onto its equatorial plane (and has degree 0). This implies that S^∞ is contractible and has a cell decomposition with one cell in each dimension; the boundary of an even cell is the closure of an odd cell, and the boundary of an odd cell is zero. The 2-dimensional skeleton of this cell decomposition is the disc D^2 decomposed into three cells as described in Section 4.3. The cellular cochain complex of S^∞ can be identified with the Koszul algebra

$$\Lambda[u] \otimes \mathbb{Z}[v], \quad \deg u = 1, \deg v = 2, \quad du = v,\ dv = 0.$$

The functoriality of the polyhedral product (Proposition 4.1.11 (a)) implies that there is the following deformation retraction onto a cellular subcomplex:

$$\mathcal{Z}_{\mathcal{K}} = (D^2, S^1)^{\mathcal{K}} \hookrightarrow (S^\infty, S^1)^{\mathcal{K}} \longrightarrow (D^2, S^1)^{\mathcal{K}}.$$

Furthermore, the cellular cochains of the polyhedral product $(S^\infty, S^1)^{\mathcal{K}}$ are identified with the Koszul algebra $\Lambda[u_1, \ldots, u_m] \otimes \mathbb{Z}[\mathcal{K}]$, in the same way as $\mathcal{C}^*(\mathcal{Z}_{\mathcal{K}})$ is identified with $R^*(\mathcal{K})$. Since $\mathcal{Z}_{\mathcal{K}} \subset (S^\infty, S^1)^{\mathcal{K}}$ is a deformation retract, the corresponding cellular cochain map

$$\Lambda[u_1, \ldots, u_m] \otimes \mathbb{Z}[\mathcal{K}] = \mathcal{C}^*\big((S^\infty, S^1)^{\mathcal{K}}\big) \longrightarrow \mathcal{C}^*(\mathcal{Z}_{\mathcal{K}}) = R^*(\mathcal{K})$$

induces an isomorphism in cohomology. The above map is a homomorphism of algebras, so it is a quasi-isomorphism. In fact, the cochain homotopy map constructed in the proof of Lemma 3.2.6 is nothing but the cellular cochain map induced by the homotopy above.

Cellular diagonal approximation. Here we establish the cohomology ring isomorphism in Lemma 4.4.1. The difficulty of working with cellular cochains is that they do not admit a functorial associative multiplication. The diagonal map used in the definition of the cohomology product is not cellular, and a cellular approximation cannot be made functorial with respect to arbitrary cellular maps. Here we construct a canonical cellular diagonal approximation $\widetilde{\varDelta}\colon \mathcal{Z}_{\mathcal{K}} \to \mathcal{Z}_{\mathcal{K}} \times \mathcal{Z}_{\mathcal{K}}$ which is functorial with respect to maps of $\mathcal{Z}_{\mathcal{K}}$ induced by simplicial maps of $\mathcal{K}$, and show that the resulting product in the cellular cochains of $\mathcal{Z}_{\mathcal{K}}$ coincides with the product in $R^*(\mathcal{K})$.

The product in the cohomology of a cell complex X is defined as follows (see [**172**], [**115**]). Consider the composite map of cellular cochain complexes

$$\mathcal{C}^*(X) \otimes \mathcal{C}^*(X) \xrightarrow{\ \times\ } \mathcal{C}^*(X \times X) \xrightarrow{\ \widetilde{\varDelta}^*\ } \mathcal{C}^*(X). \tag{4.11}$$

Here the map $\times$ sends a cellular cochain $c_1 \otimes c_2 \in \mathcal{C}^{q_1}(X) \otimes \mathcal{C}^{q_2}(X)$ to the cochain $c_1 \times c_2 \in \mathcal{C}^{q_1+q_2}(X \times X)$, whose value on a cell $e_1 \times e_2 \in X \times X$ is $(-1)^{q_1 q_2} c_1(e_1) c_2(e_2)$. The map $\widetilde{\varDelta}^*$ is induced by a cellular map $\widetilde{\varDelta}$ (a cellular *diagonal approximation*) homotopic to the diagonal $\varDelta\colon X \to X \times X$. In cohomology, map (4.11) induces a multiplication $H^*(X) \otimes H^*(X) \to H^*(X)$ which does not depend on a choice of cellular approximation and is functorial. However, map (4.11) itself is not functorial because the choice of a cellular approximation is not canonical.

Nevertheless, in the case $X = \mathcal{Z}_{\mathcal{K}}$ we can use the following construction.

CONSTRUCTION 4.4.2 (cellular approximation for $\varDelta\colon \mathcal{Z}_{\mathcal{K}} \to \mathcal{Z}_{\mathcal{K}} \times \mathcal{Z}_{\mathcal{K}}$). Consider the map $\widetilde{\varDelta}\colon \mathbb{D} \to \mathbb{D} \times \mathbb{D}$ given in the polar coordinates $z = \rho e^{i\varphi} \in \mathbb{D}$, $0 \leqslant \rho \leqslant 1$, $0 \leqslant \varphi < 2\pi$, by the formula

$$\rho e^{i\varphi} \mapsto \begin{cases} (1 - \rho + \rho e^{2i\varphi}, 1) & \text{for } 0 \leqslant \varphi \leqslant \pi, \\ (1, 1 - \rho + \rho e^{2i\varphi}) & \text{for } \pi \leqslant \varphi < 2\pi. \end{cases} \tag{4.12}$$

It is easy to see that this is a cellular map homotopic to the diagonal $\varDelta\colon \mathbb{D} \to \mathbb{D} \times \mathbb{D}$, and its restriction to the boundary circle $\mathbb{S}$ is a diagonal approximation for $\mathbb{S}$, as described by the following diagram:

$$\begin{tikzcd} \mathbb{S} \arrow[r] \arrow[d, "\widetilde{\varDelta}"'] & \mathbb{D} \arrow[d, "\widetilde{\varDelta}"] \\ \mathbb{S} \times \mathbb{S} \arrow[r] & \mathbb{D} \times \mathbb{D} \end{tikzcd} \tag{4.13}$$

(explicit formulae for the homotopies involved can be found in Exercise 4.4.11). Taking the m-fold product we obtain a cellular approximation $\widetilde{\varDelta}\colon \mathbb{D}^m \to \mathbb{D}^m \times \mathbb{D}^m$. Applying Proposition 4.1.11 (a) to the map of pairs $\widetilde{\varDelta}\colon (\mathbb{D},\mathbb{S}) \to (\mathbb{D}\times\mathbb{D}, \mathbb{S}\times\mathbb{S})$ and observing that $(\mathbb{D}\times\mathbb{D},\mathbb{S}\times\mathbb{S})^{\mathcal{K}} \cong \mathcal{Z}_{\mathcal{K}} \times \mathcal{Z}_{\mathcal{K}}$, we obtain that $\widetilde{\varDelta}\colon \mathbb{D}^m \to \mathbb{D}^m \times \mathbb{D}^m$ restricts to a cellular approximation of the diagonal map of $\mathcal{Z}_{\mathcal{K}}$, as described in the following diagram:

$$\begin{tikzcd}
\mathcal{Z}_{\mathcal{K}} \arrow[r] \arrow[d, "\widetilde{\varDelta}"'] & \mathbb{D}^m \arrow[d, "\widetilde{\varDelta}"] \\
\mathcal{Z}_{\mathcal{K}} \times \mathcal{Z}_{\mathcal{K}} \arrow[r] & \mathbb{D}^m \times \mathbb{D}^m.
\end{tikzcd}$$

Finally, applying Proposition 4.1.11 (c) to diagram (4.13) we obtain that the approximation $\widetilde{\varDelta}$ is functorial with respect to the maps of moment-angle-complexes $\mathcal{Z}_{\mathcal{K}} \to \mathcal{Z}_{\mathcal{L}}$ induced by simplicial maps $\mathcal{K} \to \mathcal{L}$.

LEMMA 4.4.3. *The cellular cochain algebra* $\mathcal{C}^*(\mathcal{Z}_{\mathcal{K}})$ *with the product defined via the diagonal approximation* $\widetilde{\varDelta}\colon \mathcal{Z}_{\mathcal{K}} \to \mathcal{Z}_{\mathcal{K}} \times \mathcal{Z}_{\mathcal{K}}$ *and map* (4.11) *is isomorphic to the algebra* $R^*(\mathcal{K})$. *We therefore have an isomorphism of cohomology rings*

$$H[R^*(\mathcal{K})] \cong H^*(\mathcal{Z}_{\mathcal{K}}).$$

PROOF. We first consider the case $\mathcal{K} = \Delta^0$, i.e. $\mathcal{Z}_{\mathcal{K}} = \mathbb{D}$. The cellular cochain complex has basis of cochains $1 \in \mathcal{C}^0(\mathbb{D})$, $T^* \in \mathcal{C}^1(\mathbb{D})$ and $D^* \in \mathcal{C}^2(\mathbb{D})$ dual to the cells introduced in Section 4.3. The multiplication defined by (4.11) in $\mathcal{C}^*(\mathbb{D})$ is trivial, so we have a ring isomorphism

$$R^*(\Delta^0) = \Lambda[u] \otimes \mathbb{Z}[v]/(v^2 = uv = 0) \longrightarrow \mathcal{C}^*(\mathbb{D}).$$

Taking an m-fold tensor product we obtain a ring isomorphism for $\mathcal{K} = \Delta^{m-1}$:

$$f\colon R^*(\Delta^{m-1}) = \Lambda[u_1,\ldots,u_m] \otimes \mathbb{Z}[v_1,\ldots,v_m]/(v_i^2 = u_iv_i = 0) \longrightarrow \mathcal{C}^*(\mathbb{D}^m).$$

Now for arbitrary $\mathcal{K}$ we have an inclusion $\mathcal{Z}_{\mathcal{K}} \subset \mathbb{D}^m = \mathcal{Z}_{\Delta^{m-1}}$ of a cellular subcomplex and the corresponding ring homomorphism $q\colon \mathcal{C}^*(\mathbb{D}^m) \to \mathcal{C}^*(\mathcal{Z}_{\mathcal{K}})$. Consider the commutative diagram

$$\begin{tikzcd}
R^*(\Delta^{m-1}) \arrow[r, "f"] \arrow[d, "p"'] & \mathcal{C}^*(\mathbb{D}^m) \arrow[d, "q"] \\
R^*(\mathcal{K}) \arrow[r, "g"] & \mathcal{C}^*(\mathcal{Z}_{\mathcal{K}}).
\end{tikzcd}$$

Here the maps p, f and q are ring homomorphisms, and g is an isomorphism of groups by Lemma 4.4.1. We claim that g is also a ring isomorphism. Indeed, take $\alpha, \beta \in R^*(\mathcal{K})$. Since p is onto, we have $\alpha = p(\alpha')$ and $\beta = p(\beta')$. Then

$$g(\alpha\beta) = gp(\alpha'\beta') = qf(\alpha'\beta') = qf(\alpha')qf(\beta') = gp(\alpha')gp(\beta') = g(\alpha)g(\beta),$$

as claimed. Thus, g is a ring isomorphism. □

Main result. By combining the results of Lemmata A.2.10, 3.2.6 and 4.4.3 we obtain the main result of this section:

THEOREM 4.4.4. *There are isomorphisms, functorial in* $\mathcal{K}$, *of bigraded algebras*

$$\begin{aligned} H^{*,*}(\mathcal{Z}_{\mathcal{K}}) &\cong \operatorname{Tor}_{\mathbb{Z}[v_1,\ldots,v_m]}\bigl(\mathbb{Z}[\mathcal{K}],\mathbb{Z}\bigr) \\ &\cong H\bigl[\Lambda[u_1,\ldots,u_m] \otimes \mathbb{Z}[\mathcal{K}], d\bigr], \end{aligned}$$

where the bigrading and the differential on the right hand side are defined by

$$\operatorname{bideg} u_i = (-1, 2), \quad \operatorname{bideg} v_i = (0, 2), \qquad du_i = v_i, \quad dv_i = 0.$$

The algebraic Betti numbers (3.2) of the face ring $\mathbb{Z}[\mathcal{K}]$ therefore acquire a topological interpretation as the bigraded Betti numbers (4.9) of the moment-angle complex $\mathcal{Z}_{\mathcal{K}}$.

Now we combine results of Propositions 3.1.5, 3.2.2 and 4.3.1, Corollary 4.2.3 and Theorem 4.4.4 in the following statement describing the functorial properties of the correspondence $\mathcal{K} \mapsto \mathcal{Z}_{\mathcal{K}}$.

PROPOSITION 4.4.5. *Consider the following functors:*

(a) $\mathcal{Z}$*, the covariant functor* $\mathcal{K} \mapsto \mathcal{Z}_{\mathcal{K}}$ *from the category of finite simplicial complexes and simplicial maps to the category of spaces with torus actions and equivariant maps (the moment-angle complex functor);*

(b) $\mathbf{k}[\cdot]$*, the contravariant functor* $\mathcal{K} \mapsto \mathbf{k}[\mathcal{K}]$ *from simplicial complexes to graded* $\mathbf{k}$*-algebras (the face ring functor);*

(c) Tor-alg*, the contravariant functor*

$$\mathcal{K} \mapsto \operatorname{Tor}_{\mathbf{k}[v_1,\ldots,v_m]}\big(\mathbf{k}[\mathcal{K}], \mathbf{k}\big)$$

from simplicial complexes to bigraded $\mathbf{k}$*-algebras (the* Tor*-algebra functor; it is the composition of* $\mathbf{k}[\cdot]$ *and* $\operatorname{Tor}_{\mathbf{k}[v_1,\ldots,v_m]}(\cdot\,, \mathbf{k})$*);*

(d) H_T^**, the contravariant functor* $X \mapsto H_T^*(X; \mathbf{k})$ *from spaces with torus actions to* $\mathbf{k}$*-algebras (the equivariant cohomology functor);*

(e) H^**, the contravariant functor* $X \mapsto H^*(X; \mathbf{k})$ *from spaces to* $\mathbf{k}$*-algebras (the ordinary cohomology functor).*

Then we have the following identities:

$$H_T^* \circ \mathcal{Z} = \mathbf{k}[\cdot], \qquad H^* \circ \mathcal{Z} = \text{Tor-alg}.$$

The second identity implies that for any simplicial map $\varphi\colon \mathcal{K} \to \mathcal{L}$ *the corresponding cohomology map*

$$\varphi_{\mathcal{Z}}^*\colon H^*(\mathcal{Z}_{\mathcal{L}}) \to H^*(\mathcal{Z}_{\mathcal{K}})$$

coincides with the induced homomorphism of Tor*-algebras* φ_{Tor}^* *from Proposition 3.2.2. In particular, the map* φ *gives rise to a map*

$$H^{-q,2p}(\mathcal{Z}_{\mathcal{L}}) \to H^{-q,2p}(\mathcal{Z}_{\mathcal{K}})$$

of bigraded cohomology.

In the case of Cohen–Macaulay complexes $\mathcal{K}$ (see Section 3.3) we have the following version of Theorem 4.4.4.

PROPOSITION 4.4.6. *Let* $\mathcal{K}$ *be an* $(n-1)$*-dimensional Cohen–Macaulay complex, and let* $\boldsymbol{t}$ *be an hsop in* $\mathbf{k}[\mathcal{K}]$*. Then we have the following isomorphism of algebras:*

$$H^*(\mathcal{Z}_{\mathcal{K}}; \mathbf{k}) \cong \operatorname{Tor}_{\mathbf{k}[v_1,\ldots,v_m]/\boldsymbol{t}}\big(\mathbf{k}[\mathcal{K}]/\boldsymbol{t}, \mathbf{k}\big).$$

PROOF. This follows from Theorem 4.4.4 and Lemma A.3.5. □

Note that the algebra $\mathbf{k}[\mathcal{K}]/\boldsymbol{t}$ is finite-dimensional as a $\mathbf{k}$-vector space, unlike $\mathbf{k}[\mathcal{K}]$. In some circumstances this observation allows us to calculate the cohomology of $\mathcal{Z}_{\mathcal{K}}$ more effectively.

Description of the product in terms of full subcomplexes. The Hochster formula (Theorem 3.2.4) for the components of the Tor-algebra can be used to obtain an alternative description of the product structure in $H^*(\mathcal{Z}_\mathcal{K})$.

We recall from Section 3.2 that the bigraded structure in the Tor-algebra is refined to a multigrading, and the multigraded components of Tor can be calculated in terms of the full subcomplexes of $\mathcal{K}$:

$$\mathrm{Tor}^{-i,2J}_{\mathbb{Z}[v_1,\ldots,v_m]}\bigl(\mathbb{Z}[\mathcal{K}],\mathbb{Z}\bigr)\cong\widetilde{H}^{|J|-i-1}(\mathcal{K}_J),$$

where $J\subset[m]$, see Theorem 3.2.9. Furthermore, the product in the Tor-algebra defines a product in the direct sum $\bigoplus_{p\geqslant 0,\ J\subset[m]}\widetilde{H}^{p-1}(\mathcal{K}_J)$ given by (3.11).

The bigraded structure in the cellular cochain complex of $\mathcal{Z}_\mathcal{K}$ defined in Section 4.3 can be also refined to a multigrading (a $\mathbb{Z}\oplus\mathbb{Z}^m$-grading):

$$\mathcal{C}^*(\mathcal{Z}_\mathcal{K})=\bigoplus_{J\subset[m]}\mathcal{C}^{*,\,2J}(\mathcal{Z}_\mathcal{K}),$$

where $\mathcal{C}^{*,\,2J}(\mathcal{Z}_\mathcal{K})$ is the subcomplex spanned by the cochains $\varkappa(J\backslash I,I)^*$ with $I\subset J$ and $I\in\mathcal{K}$. The bigraded cohomology groups are decomposed as follows:

$$H^{-i,\,2j}(\mathcal{Z}_\mathcal{K})=\bigoplus_{J\subset[m]\colon\,|J|=j}H^{-i,\,2J}(\mathcal{Z}_\mathcal{K}),$$

where $H^{-i,\,2J}(\mathcal{Z}_\mathcal{K})=H^{-i}[\mathcal{C}^{*,\,2J}(\mathcal{Z}_\mathcal{K})]$.

THEOREM 4.4.7 (Baskakov [**18**]). *There are isomorphisms*

$$\widetilde{H}^{p-1}(\mathcal{K}_J)\xrightarrow{\cong}H^{p-|J|,2J}(\mathcal{Z}_\mathcal{K}),$$

which are functorial with respect to simplicial maps and induce a ring isomorphism

$$h\colon\bigoplus_{\substack{p\geqslant 0\\ J\subset[m]}}\widetilde{H}^{p-1}(\mathcal{K}_J)\xrightarrow{\cong}H^*(\mathcal{Z}_\mathcal{K}).$$

PROOF. The statement about the additive isomorphisms follows from Theorems 3.2.9 and 4.4.4. In explicit terms, the cohomology isomorphisms are induced by the cochain isomorphisms given by

$$\begin{aligned}C^{p-1}(\mathcal{K}_J)&\longrightarrow\mathcal{C}^{p-|J|,2J}(\mathcal{Z}_\mathcal{K}),\\ \alpha_L&\longmapsto\varepsilon(L,J)\varkappa(J\setminus L,L)^*\end{aligned}$$

similar to (3.9), where $\alpha_L\in C^{p-1}(\mathcal{K}_J)$ is the cochain dual to a simplex $L\in\mathcal{K}_J$.

The ring isomorphism follows from Proposition 3.2.10 and Theorem 4.4.4. □

For the bigraded and ordinary cohomology of $\mathcal{Z}_\mathcal{K}$ we obtain the following formulae in terms of full subcomplexes of $\mathcal{K}$:

THEOREM 4.4.8. *There are isomorphisms*

$$\begin{aligned}H^{-i,2j}(\mathcal{Z}_\mathcal{K})&\cong\bigoplus_{J\subset[m]\colon\,|J|=j}\widetilde{H}^{j-i-1}(\mathcal{K}_J),\\ H^k(\mathcal{Z}_\mathcal{K})&\cong\bigoplus_{J\subset[m]}\widetilde{H}^{k-|J|-1}(\mathcal{K}_J).\end{aligned}$$

It follows that the cohomology of $\mathcal{Z}_\mathcal{K}$ may have arbitrary torsion:

Corollary 4.4.9. *Any finite abelian group can appear as a summand in a cohomology group of* $H^*(\mathcal{Z}_\mathcal{K})$ *for some* $\mathcal{K}$.

Proof. It follows from the previous theorem that $\widetilde{H}^*(\mathcal{K})$ is a direct summand in $H^*(\mathcal{Z}_\mathcal{K})$ (with appropriate shifts in dimension). Therefore, we can take $\mathcal{K}$ whose simplicial cohomology contains the appropriate torsion. □

Exercises.

4.4.10. Let $\mathbb{S}$ be the standard unit circle decomposed into two cells, where the 0-cell is the unit. The map

$$\widetilde{\varDelta}\colon \mathbb{S}\to\mathbb{S}\times\mathbb{S},\qquad e^{i\varphi}\mapsto\begin{cases}(e^{2i\varphi},1) & \text{for } 0\leqslant\varphi\leqslant\pi,\\ (1,e^{2i\varphi}) & \text{for } \pi\leqslant\varphi<2\pi\end{cases}$$

is a cellular diagonal approximation. It is obtained by restricting map (4.12) to the boundary circle ($\rho=1$). A homotopy F_t between the diagonal $\varDelta\colon\mathbb{S}\to\mathbb{S}\times\mathbb{S}$ ($t=0$) and its cellular approximation $\widetilde{\varDelta}$ ($t=1$) is given by

$$F_t\colon \mathbb{S}\to\mathbb{S}\times\mathbb{S},\qquad e^{i\varphi}\mapsto\begin{cases}\bigl(e^{i(1+t)\varphi},e^{i(1-t)\varphi}\bigr) & \text{for } 0\leqslant\varphi\leqslant\pi,\\ \bigl(e^{i(1-t)\varphi+2\pi it},e^{i(1+t)\varphi-2\pi it}\bigr) & \text{for } \pi\leqslant\varphi<2\pi.\end{cases}$$

4.4.11. Show that the formula

$$\rho e^{i\varphi}\mapsto\begin{cases}\bigl((1-\rho)t+\rho e^{i(1+t)\varphi},(1-\rho)t+\rho e^{i(1-t)\varphi}\bigr) & \text{for } 0\leqslant\varphi\leqslant\pi,\\ \bigl((1-\rho)t+\rho e^{i(1-t)\varphi+2\pi it},(1-\rho)t+\rho e^{i(1+t)\varphi-2\pi it}\bigr) & \text{for } \pi\leqslant\varphi<2\pi.\end{cases}$$

defines a homotopy $G_t\colon\mathbb{D}\to\mathbb{D}\times\mathbb{D}$ between the diagonal $\varDelta\colon\mathbb{D}\to\mathbb{D}\times\mathbb{D}$ ($t=0$) and its approximation $\widetilde{\varDelta}$ (4.13) ($t=1$). For $\rho=1$ the homotopy G_t restricts to the homotopy F_t of the previous exercise.

4.5. Bigraded Betti numbers

Here we describe the main properties of the bigraded Betti numbers of moment-angle complexes and give some examples of explicit calculations.

Lemma 4.5.1. *Let* $\mathcal{K}$ *be a simplicial complex of dimension* $n-1$ *with* $f_0=m$ *vertices and* f_1 *edges, so that* $\dim\mathcal{Z}_\mathcal{K}=m+n$. *We have*

(a) $b^{0,0}(\mathcal{Z}_\mathcal{K})=b^0(\mathcal{Z}_\mathcal{K})=1$ *and* $b^{0,2q}(\mathcal{Z}_\mathcal{K})=0$ *for* $q\neq0$*;*
(b) $b^{-p,2q}=0$ *for* $q>m$ *or* $p>q$*;*
(c) $b^1(\mathcal{Z}_\mathcal{K})=b^2(\mathcal{Z}_\mathcal{K})=0$*;*
(d) $b^3(\mathcal{Z}_\mathcal{K})=b^{-1,4}(\mathcal{Z}_\mathcal{K})=\binom{m}{2}-f_1$*;*
(e) $b^{-p,2q}(\mathcal{Z}_\mathcal{K})=0$ *for* $p\geqslant q>0$ *or* $q-p>n$*;*
(f) $b^{m+n}(\mathcal{Z}_\mathcal{K})=b^{-(m-n),2m}(\mathcal{Z}_\mathcal{K})=\operatorname{rank}\widetilde{H}^{n-1}(\mathcal{K})$.

Proof. We consider the algebra $R^*(\mathcal{K})$ whose cohomology is $H^*(\mathcal{Z}_\mathcal{K})$. Recall that $R^*(\mathcal{K})$ has additive basis of monomials u_Jv_I with $I\in\mathcal{K}$ and $I\cap J=\varnothing$. Since $\operatorname{bideg}v_i=(0,2)$, $\operatorname{bideg}u_j=(-1,2)$, the bigraded component $R^{-p,2q}(\mathcal{K})$ has basis of monomials u_Jv_I with $|I|=q-p$ and $|J|=p$. In particular, $R^{-p,2q}(\mathcal{K})=0$ for $q>m$ or $p>q$, which implies (b). To prove (a) we observe that $R^{0,0}(\mathcal{K})=\mathbf{k}$ and each $v_I\in R^{0,2q}(\mathcal{K})$ with $q>0$ is a coboundary, hence, $H^{0,2q}(\mathcal{Z}_\mathcal{K})=0$ for $q>0$.

Now we prove (e). Let $u_Jv_I\in R^{-p,2q}(\mathcal{K})$; then $|I|=q-p$ and $I\in\mathcal{K}$. Since a simplex of $\mathcal{K}$ has at most n vertices, $R^{-p,2q}(\mathcal{K})=0$ for $q-p>n$. We have $b^{-p,2q}(\mathcal{Z}_\mathcal{K})=0$ for $p>q$ by (b) so we need only to check that $b^{-q,2q}(\mathcal{Z}_\mathcal{K})=0$ for

$q > 0$. The group $R^{-q,2q}(\mathcal{K})$ has basis of monomials u_J with $|J| = q$. Since $d(u_i) = v_i$, there are no nonzero cocycles in $R^{-q,2q}(\mathcal{K})$ for $q > 0$, hence, $H^{-q,2q}(\mathcal{Z}_\mathcal{K}) = 0$.

Statement (c) follows from (e) and (4.10).

We also have $H^3(\mathcal{Z}_\mathcal{K}) = H^{-1,4}(\mathcal{Z}_\mathcal{K})$, by (e). There is a basis in $R^{-1,4}(\mathcal{K})$ consisting of monomials u_jv_i with $i \neq j$. We have $d(u_jv_i) = v_iv_j$ and $d(u_iu_j) = u_jv_i - u_iv_j$. Hence, u_jv_i is a cocycle if and only if $\{i,j\} \notin \mathcal{K}$; in this case the two cocycles u_jv_i and u_iv_j represent the same cohomology class. This proves (d).

It remains to prove (f). The total degree of a monomial $u_Jv_I \in R^*(\mathcal{K})$ is $2|I| + |J|$, and there are constraints $|I| + |J| \leqslant m$ and $|I| \leqslant n$. Therefore, the maximum of the total degree is achieved for $|I| = n$ and $|J| = m - n$. This proves the first identity of (f), and the second follows from Theorem 4.4.8. □

Lemma 4.5.1 shows that nonzero bigraded Betti numbers $b^{r,2q}(\mathcal{Z}_\mathcal{K})$ with $r \neq 0$ appear only in the strip bounded by the lines $r = -1$, $q = m$, $r + q = 1$ and $r + q = n$ in the second quadrant, see Fig. 4.1 (a).



(a) arbitrary $\mathcal{K}$ (b) $|\mathcal{K}| \cong S^{n-1}$

FIGURE 4.1. Possible locations on nonzero $b^{-p,2q}(\mathcal{Z}_\mathcal{K})$ (marked by $*$).

The next result allows us to express the numbers of faces of $\mathcal{K}$ (i.e. its f- and h-vectors) via the bigraded Betti numbers. We consider the Euler characteristics of the complexes $\mathcal{C}^{*,2q}(\mathcal{Z}_\mathcal{K})$ (see (4.8)),

$$\chi_q(\mathcal{Z}_\mathcal{K}) = \sum_{p=0}^{m}(-1)^p \operatorname{rank} \mathcal{C}^{-p,2q}(\mathcal{Z}_\mathcal{K}) = \sum_{p=0}^{m}(-1)^p b^{-p,2q}(\mathcal{Z}_\mathcal{K}) \tag{4.14}$$

and define the generating series

$$\chi(\mathcal{Z}_\mathcal{K}; t) = \sum_{q=0}^{m} \chi_q(\mathcal{Z}_\mathcal{K}) t^{2q}.$$

THEOREM 4.5.2. *The following identity holds for an $(n-1)$-dimensional simplicial complex $\mathcal{K}$ with m vertices:*

$$\chi(\mathcal{Z}_\mathcal{K}; t) = (1 - t^2)^{m-n}(h_0 + h_1t^2 + \cdots + h_nt^{2n}).$$

Here $(h_0, h_1, \ldots, h_n)$ is the h-vector of $\mathcal{K}$.

PROOF. The bigraded component $\mathcal{C}^{-p,2q}(\mathcal{Z}_\mathcal{K})$ has basis of cellular cochains $\varkappa(J,I)^*$ with $I \in \mathcal{K}$, $|I| = q-p$ and $|J| = p$. Therefore, $\operatorname{rank} \mathcal{C}^{-p,2q}(\mathcal{Z}_\mathcal{K}) = f_{q-p-1}\binom{m-q+p}{p}$, where $(f_0, f_1, \ldots, f_{n-1})$ is the f-vector of $\mathcal{K}$ and $f_{-1} = 1$. By substituting this into (4.14) we obtain

$$\chi_q(\mathcal{Z}_\mathcal{K}) = \sum_{j=0}^{m} (-1)^{q-j} f_{j-1}\tbinom{m-j}{q-j}.$$

Then

$$\begin{aligned}\chi(\mathcal{Z}_\mathcal{K};t) &= \sum_{q=0}^{m}\sum_{j=0}^{m} t^{2j}t^{2(q-j)}(-1)^{q-j} f_{j-1}\tbinom{m-j}{q-j} \\ &= \sum_{j=0}^{m} f_{j-1}t^{2j}(1-t^2)^{m-j} = (1-t^2)^m \sum_{j=0}^{n} f_{j-1}(t^{-2}-1)^{-j}.\end{aligned} \tag{4.15}$$

Set $h(t) = h_0 + h_1 t + \cdots + h_n t^n$. Then it follows from (2.3) that

$$t^n h(t^{-1}) = (t-1)^n \sum_{j=0}^{n} f_{j-1}(t-1)^{-j}.$$

By substituting t^{-2} for t in the identity above we finally rewrite (4.15) as

$$\frac{\chi(\mathcal{Z}_\mathcal{K};t)}{(1-t^2)^m} = \frac{t^{-2n}h(t^2)}{(t^{-2}-1)^n} = \frac{h(t^2)}{(1-t^2)^n},$$

which is equivalent to the required identity. □

COROLLARY 4.5.3. *If $\mathcal{K} \neq \Delta^{m-1}$, then the Euler characteristic of $\mathcal{Z}_\mathcal{K}$ is zero.*

PROOF. We have

$$\chi(\mathcal{Z}_\mathcal{K}) = \sum_{p,q=0}^{m} (-1)^{-p+2q} b^{-p,2q}(\mathcal{Z}_\mathcal{K}) = \sum_{q=0}^{m} \chi_q(\mathcal{Z}_\mathcal{K}) = \chi(\mathcal{Z}_\mathcal{K};1) = 0$$

by Theorem 4.5.2 (note that $\mathcal{K} \neq \Delta^{m-1}$ implies that $m > n$). □

We proceed by describing the properties of bigraded Betti numbers for particular classes of simplicial complexes.

DEFINITION 4.5.4. A d-dimensional *pseudomanifold* is a finite simplicial complex $\mathcal{K}$ such that

(a) all maximal simplices of $\mathcal{K}$ have dimension d (i.e. $\mathcal{K}$ is pure d-dimensional);
(b) each $(d-1)$-simplex of $\mathcal{K}$ is the face of exactly two d-simplices of $\mathcal{K}$.
(c) if I and I' are d-simplices of $\mathcal{K}$, then there is a sequence $I = I_1, I_2, \ldots, I_k = I'$ of d-simplices of $\mathcal{K}$ such that I_j and I_{j+1} have a common $(d-1)$-face for $1 \leqslant i \leqslant k-1$.

If $\mathcal{K}$ is a d-dimensional pseudomanifold, then either $H_d(\mathcal{K}) \cong \mathbb{Z}$ or 0 (an exercise). In the former case the pseudomanifold $\mathcal{K}$ is called *orientable*.

LEMMA 4.5.5. *Let $\mathcal{K}$ be an orientable pseudomanifold of dimension $n-1$ with m vertices. Then*

$$H^{m+n}(\mathcal{Z}_\mathcal{K}) = \widetilde{H}^{n-1}(\mathcal{K}) \cong \mathbb{Z}.$$

Under the isomorphism $H^*(\mathcal{Z}_\mathcal{K}) \cong H[R^*(\mathcal{K})]$*, the group above is generated by the class of any monomial* $u_J v_I \in R^*(\mathcal{K})$ *of bidegree* $(-(m-n), 2m)$ *such that* $I \in \mathcal{K}$ *and* $J = [m] \setminus I$.

Proof. The isomorphism of groups follows from Theorem 4.4.8 and the fact that $\mathcal{K}$ is orientable. We have $H^{m+n}(\mathcal{Z}_\mathcal{K}) = H^{-(m-n),2m}(\mathcal{Z}_\mathcal{K})$. The group $R^{-(m-n),2m}(\mathcal{K})$ has basis of monomials $u_J v_I$ with $I \in \mathcal{K},\ |I| = n$ and $J = [m] \setminus I$. Each of these monomials is a cocycle. Let I, I' be two $(n-1)$-simplices of $\mathcal{K}$ having a common $(n-2)$-face. Consider the corresponding cocycles $u_J v_I$ and $u_{J'} v_{I'}$ (where $J = [m] \setminus I,\ J' = [m] \setminus I'$):

$$u_J v_I = u_{j_1} u_{j_2} \cdots u_{j_{m-n}} v_{i_1} \cdots v_{i_{n-1}} v_{i_n},$$
$$u_{J'} v_{I'} = u_{i_n} u_{j_2} \cdots u_{j_{m-n}} v_{i_1} \cdots v_{i_{n-1}} v_{j_1}.$$

Since $\mathcal{K}$ is a pseudomanifold, the $(n-2)$-face $\{i_1, \ldots, i_{n-1}\}$ is contained in exactly two $(n-1)$-faces, namely $I = \{i_1, \ldots, i_{n-1}, i_n\}$ and $I' = \{i_1, \ldots, i_{n-1}, j_1\}$. Therefore we have the following identity in $R^*(\mathcal{K})$

$$\begin{aligned} &d(u_{i_n} u_{j_1} u_{j_2} \cdots u_{j_{m-n}} v_{i_1} \cdots v_{i_{n-1}}) = \\ &\qquad = u_{j_1} u_{j_2} \cdots u_{j_{m-n}} v_{i_1} \cdots v_{i_{n-1}} v_{i_n} - u_{i_n} u_{j_2} \cdots u_{j_{m-n}} v_{i_1} \cdots v_{i_{n-1}} v_{j_1} \end{aligned}$$

Hence, $[u_J v_I] = [u_{J'} v_{I'}]$ (as cohomology classes). Property (c) from the definition of a pseudomanifold implies that all monomials $u_J v_I \in R^{-(m-n),2m}(\mathcal{K})$ represent the same cohomology class up to sign. The isomorphism (3.9) takes $u_J v_I$ to $\pm\alpha_I \in C^{n-1}(\mathcal{K})$, which represents a generator of $\widetilde{H}^{n-1}(\mathcal{K}) \cong \mathbb{Z}$ (see Exercise 4.5.17). □

Remark. If $\mathcal{K}$ is a non-orientable pseudomanifold, then the same argument shows that any monomial $u_J v_I \in R^*(\mathcal{K})$ as above represents the generator of $H^{m+n}(\mathcal{Z}_\mathcal{K}) = H^{n-1}(\mathcal{K}) \cong \mathbb{Z}_2$.

Proposition 4.5.6. *Let* $\mathcal{K}$ *be a triangulated sphere of dimension* $n-1$*. Then Poincaré duality for the moment-angle manifold* $\mathcal{Z}_\mathcal{K}$ *respects the bigrading in cohomology. In particular,*

$$b^{-p,\,2q}(\mathcal{Z}_\mathcal{K}) = b^{-(m-n)+p,\,2(m-q)}(\mathcal{Z}_\mathcal{K}) \quad \textit{for } 0 \leqslant p \leqslant m-n,\ 0 \leqslant q \leqslant m.$$

Proof. The Poincaré duality maps (see Definition 3.4.3) are defined via the cohomology multiplication in $H^*(\mathcal{Z}_\mathcal{K})$, which respects the bigrading. We have $\dim \mathcal{Z}_\mathcal{K} = m + n$, and

$$H^{m+n}(\mathcal{Z}_\mathcal{K}) = \operatorname{Tor}^{-(m-n),\,2m}_{\mathbb{Z}[v_1,\ldots,v_m]}\bigl(\mathbb{Z}[\mathcal{K}], \mathbb{Z}\bigr) \cong \mathbb{Z},$$

by Lemma 4.5.5. This implies the required identity for the Betti numbers. □

Corollary 4.5.7. *Let* $\mathcal{K}$ *be a triangulated* $(n-1)$*-sphere and* $\mathcal{Z}_\mathcal{K}$ *the corresponding moment-angle manifold,* $\dim \mathcal{Z}_\mathcal{K} = m + n$*. Then*

(a) $b^{-p,2q}(\mathcal{Z}_\mathcal{K}) = 0$ *for* $p \geqslant m - n$*, with the only exception* $b^{-(m-n),2m} = 1$*;*
(b) $b^{-p,2q}(\mathcal{Z}_\mathcal{K}) = 0$ *for* $q - p \geqslant n$*, with the only exception* $b^{-(m-n),2m} = 1$.

It follows that if $|\mathcal{K}| \cong S^{n-1}$, then nonzero bigraded Betti numbers $b^{r,2q}(\mathcal{Z}_\mathcal{K})$, except $b^{0,0}(\mathcal{Z}_\mathcal{K})$ and $b^{-(m-n),2m}(\mathcal{Z}_\mathcal{K})$, appear only in the strip bounded by the lines $r = -(m-n-1),\ r = -1,\ r + q = 1$ and $r + q = n - 1$ in the second quadrant, see Fig. 4.1 (b).

A space X is called a *Poincaré duality space* (over $\mathbf{k}$) if $H^*(X;\mathbf{k})$ is a Poincaré algebra (see Definition 3.4.3). We have the following characterisation of moment-angle complexes with Poincaré duality, extending the result of Corollary 4.5.6.

THEOREM 4.5.8. *$\mathcal{Z}_\mathcal{K}$ is a Poincaré duality space over a field $\mathbf{k}$ if and only $\mathcal{K}$ is a Gorenstein complex over $\mathbf{k}$.*

PROOF. Assume that $\mathcal{K}$ is a Gorenstein complex. Consider the algebra T defined in Theorem 3.4.4, i.e. $T=\bigoplus_{i=0}^d T^i$, where $T^i=\mathrm{Tor}^{-i}_{\mathbf{k}[m]}(\mathbf{k}[\mathcal{K}],\mathbf{k})$ and $d=\max\{j\colon \mathrm{Tor}^{-j}_{\mathbf{k}[m]}(\mathbf{k}[\mathcal{K}],\mathbf{k})\neq 0\}$. Since T is Poincaré algebra, $\mathbf{k}\cong T^0\cong \mathrm{Hom}_\mathbf{k}(T^d,T^d)$, which implies that $T^d\cong\mathbf{k}$. Since T has a bigrading, we obtain $T^d=T^{d,2q}$ for some $q\geqslant 0$. Since the multiplication in T respects the bigrading, the isomorphisms $T^i\overset{\cong}{\to}\mathrm{Hom}_\mathbf{k}(T^{d-i},T^d)$ from the definition of a Poincaré algebra split into isomorphisms

$$T^{i,\,2j}\xrightarrow{\cong}\mathrm{Hom}_\mathbf{k}(T^{d-i,\,2(q-j)},T^{d,\,2q}).$$

Let $H^k=H^k(\mathcal{Z}_\mathcal{K};\mathbf{k})$ and $H=\bigoplus_{k=0}^r H^k$; then $H^k=\bigoplus_{i+2j=k}T^{i,\,2j}$ and $r=d+2q$. Therefore, we have the isomorphisms

$$H^k=\bigoplus_{i+2j=k}T^{i,2j}\xrightarrow{\cong}\bigoplus_{i+2j=k}\mathrm{Hom}_\mathbf{k}(T^{d-i,2(q-j)},T^{d,2q})=\mathrm{Hom}_\mathbf{k}(H^{r-k},H^r),$$

which imply that H is a Poincaré algebra.

Now assume that $H=\bigoplus_{k=0}^r H^k$ is a Poincaré algebra. Then

$$\mathbf{k}\cong H^r=\mathrm{Tor}^{-d,\,2q}_{\mathbf{k}[v_1,\dots,v_m]}\bigl(\mathbf{k}[\mathcal{K}],\mathbf{k}\bigr)=T^{d,\,2q}$$

for some $d,q\geqslant 0$. Since the multiplication in the cohomology of $\mathcal{Z}_\mathcal{K}$ respects the bigrading, the isomorphisms $H^k\overset{\cong}{\to}\mathrm{Hom}_\mathbf{k}(H^{r-k},H^r)$ split into isomorphisms

$$H^{-i,\,2j}=T^{i,\,2j}\xrightarrow{\cong}\mathrm{Hom}_\mathbf{k}(T^{d-i,\,2(q-j)},T^{d,\,2q}),$$

which in their turn define the isomorphisms

$$T^i=\bigoplus_j T^{i,\,2j}\xrightarrow{\cong}\bigoplus_j\mathrm{Hom}_\mathbf{k}(T^{d-i,\,2(q-j)},T^d)=\mathrm{Hom}_\mathbf{k}(T^{d-i},T^d).$$

Thus, T is a Poincaré algebra. □

REMARK. We do not assume that $r=\max\{k\colon H^k(\mathcal{Z}_\mathcal{K};\mathbf{k})\neq 0\}$ is equal to $\dim\mathcal{Z}_\mathcal{K}=m+n$ in Theorem 4.5.8. It follows from the proof above that $\mathcal{Z}_\mathcal{K}$ is a Poincaré duality space with $r=\dim\mathcal{Z}_\mathcal{K}$ if and only if $\mathcal{K}$ is a Gorenstein* complex.

Here are some explicit examples of calculations of $H^*(\mathcal{Z}_\mathcal{K})$ using Theorem 4.4.4.

EXAMPLE 4.5.9. Let $\mathcal{K}=\partial\Delta^{m-1}$. Then

$$\mathbb{Z}[\mathcal{K}]=\mathbb{Z}[v_1,\dots,v_m]/(v_1\cdots v_m).$$

The cocycle $u_1v_2v_3\cdots v_m\in\Lambda[u_1,\dots,u_m]\otimes\mathbb{Z}[\mathcal{K}]$ of bidegree $(-1,2m)$ represents a generator of the top degree cohomology group of $\mathcal{Z}_\mathcal{K}\cong S^{2m-1}$.

EXAMPLE 4.5.10. Let $\mathcal{K}$ be the boundary of a 5-gon. We have $\dim\mathcal{Z}_\mathcal{K}=7$. We enumerate the vertices of $\mathcal{K}$ clockwise. The face ring of $\mathcal{K}$ is given in Example 3.1.4.3. The group $H^3(\mathcal{Z}_\mathcal{K})$ has 5 generators corresponding to the diagonals of the 5-gon; these generators are represented by the cocycles $u_iv_{i+2}\in\mathbb{Z}[\mathcal{K}]\otimes\Lambda[u_1,\dots,u_5]$, $1\leqslant i\leqslant 5$ (the summation of indices is modulo 5). A direct calculation shows that $H^4(\mathcal{Z}_\mathcal{K})$ also has 5 generators, represented by the cocycles $u_j u_{j+1} v_{j+3}$, $1 \leqslant j \leqslant 5$. The Betti vector of $\mathcal{Z}_\mathcal{K}$ is therefore given by

$$\big(b^0(\mathcal{Z}_\mathcal{K}), b^1(\mathcal{Z}_\mathcal{K}), \ldots, b^7(\mathcal{Z}_\mathcal{K})\big) = (1, 0, 0, 5, 5, 0, 0, 1).$$

By Lemma 4.5.5, the product of cocycles $u_i v_{i+2}$ and $u_j u_{j+1} v_{j+3}$ represents a generator of $H^7(\mathcal{Z}_\mathcal{K})$ if and only if all the indices $i, i+2, j, j+1, j+3$ are different. Hence, for each cohomology class $[u_i v_{i+2}] \in H^3(\mathcal{Z}_\mathcal{K})$ there is a unique class $[u_j u_{j+1} v_{j+3}] \in H^4(\mathcal{Z}_\mathcal{K})$ such that the product $[u_i v_{i+2}] \cdot [u_j u_{j+1} v_{j+3}]$ is nonzero. These calculations are summarised by the cohomology ring isomorphism

$$H^*(\mathcal{Z}_\mathcal{K}) \cong H^*\big((S^3 \times S^4)^{\#5}\big),$$

in accordance with Exercise 4.1.18.

Example 4.5.11. Now we calculate the Betti numbers and the cohomology product for $\mathcal{Z}_\mathcal{K}$ in the case when $\mathcal{K}$ is a boundary of an m-gon with $m \geqslant 4$.

It follows from Corollary 4.5.7 that the only nonzero bigraded Betti numbers of $\mathcal{Z}_\mathcal{K}$ are $b^{-(p-1),2p}$ for $2 \leqslant p \leqslant m-2$ and $b^{0,0}(\mathcal{Z}_\mathcal{K}) = b^{-(m-2),2m}(\mathcal{Z}_\mathcal{K}) = 1$. The ordinary Betti numbers are therefore given by

$$b^0(\mathcal{Z}_\mathcal{K}) = b^{m+2}(\mathcal{Z}_\mathcal{K}) = 1, \quad b^k(\mathcal{Z}_\mathcal{K}) = b^{-(k-2),2(k-1)}(\mathcal{Z}_\mathcal{K}) \quad \text{for } 3 \leqslant k \leqslant m-1.$$

To calculate $b^{-(k-2),2(k-1)}(\mathcal{Z}_\mathcal{K})$ for $3 \leqslant k \leqslant m-1$ we use the algebra $R^*(\mathcal{K})$ of Construction 3.2.5. We have

$$\begin{aligned}(4.16)\quad b^{-(k-2),2(k-1)}(\mathcal{Z}_\mathcal{K}) &= \dim H^{-(k-2),2(k-1)}[R^*, d] = \\ \dim\ker\big[d\colon R^{-(k-2),2(k-1)} &\to R^{-(k-3),2(k-1)}\big] - \dim dR^{-(k-1),2(k-1)}.\end{aligned}$$

Since $H^{-(k-1),2(k-1)}(\mathcal{Z}_\mathcal{K}) = 0$ for $k > 1$, the differential d from $R^{-(k-1),2(k-1)}$ is monomorphic, and

$$\dim dR^{-(k-1),2(k-1)} = \dim R^{-(k-1),2(k-1)}.$$

Similarly, since $H^{-(k-3),2(k-1)}(\mathcal{Z}_\mathcal{K}) = 0$ for $k \leqslant m$, the differential from $R^{-(k-2),2(k-1)}$ is epimorphic, and

$$\begin{aligned}\dim\ker\big[d\colon R^{-(k-2),2(k-1)} &\to R^{-(k-3),2(k-1)}\big] \\ &= \dim R^{-(k-2),2(k-1)} - \dim R^{-(k-3),2(k-1)}.\end{aligned}$$

Substituting the above two expressions into (4.16) and using formula (3.3) for the dimensions of $R^{-p,2q}$, we calculate

$$\begin{aligned}b^k(\mathcal{Z}_\mathcal{K}) &= b^{-(k-2),2(k-1)}(\mathcal{Z}_\mathcal{K}) \\ &= \dim R^{-(k-2),2(k-1)} - \dim R^{-(k-3),2(k-1)} - \dim R^{-(k-1),2(k-1)} \\ &= m\tbinom{m-1}{k-2} - m\tbinom{m-2}{k-3} - \tbinom{m}{k-1} \\ &= (k-2)\tbinom{m-2}{k-1} + (m-k)\tbinom{m-2}{m-k+1} \quad \text{for } 3 \leqslant k \leqslant m-1.\end{aligned}$$

This additive calculation is valid for any field $\mathbf{k}$, and therefore for $\mathbf{k} = \mathbb{Z}$ as well, because the cohomology of $\mathcal{Z}_\mathcal{K}$ does not have torsion in this example (this follows from Theorem 4.4.8).

The product of any three classes of positive degree in $H^*(\mathcal{Z}_\mathcal{K})$ is zero (i.e. the *cohomology product length* of $\mathcal{Z}_\mathcal{K}$ is 2). Indeed, if $\alpha_i \in H^{-(p_i-1),2p_i}(\mathcal{Z}_\mathcal{K})$ for $i = 1, 2, 3$, then

$$\alpha_1\alpha_2\alpha_3 \in H^{-(p_1+p_2+p_3-3),2(p_1+p_2+p_3)}(\mathcal{Z}_\mathcal{K}),$$

which is zero by Lemma 4.5.1 (e) (note that $n = 2$ in this example). Hence all nontrivial products in $H^*(\mathcal{Z}_\mathcal{K})$ arise from Poincaré duality.

The above calculations of the Betti numbers together with the observations about the cohomology product can be summarised by saying that $\mathcal{Z}_\mathcal{K}$ cohomologically looks like a connected sum of sphere products, namely,

$$H^*(\mathcal{Z}_\mathcal{K}) \cong H^*\Bigl(\mathop{\#}_{k=3}^{m-1}\bigl(S^k \times S^{m+2-k}\bigr)^{\#(k-2)\binom{m-2}{k-1}}\Bigr), \tag{4.17}$$

as rings.

The cohomology isomorphism of (4.17) is induced by a homeomorphism of manifolds. This is true even in a more general situation, when $\mathcal{K}$ is the boundary of a stacked polytope (see Definition 1.4.8):

THEOREM 4.5.12 (McGavran, see [**27**, Theorem 6.3]). *Let $\mathcal{K}$ be the boundary of a stacked polytope of dimension n with m vertices. Then the corresponding moment-angle manifold is homeomorphic to a connected sum of sphere products,*

$$\mathcal{Z}_\mathcal{K} \cong \mathop{\#}_{k=3}^{m-n+1}\bigl(S^k \times S^{m+n-k}\bigr)^{\#(k-2)\binom{m-n}{k-1}}.$$

For $n = 2$ we obtain a homeomorphism of manifolds underlying the isomorphism (4.17) (note that any 2-polytope is stacked).

The bigraded Betti numbers for stacked polytopes can also be calculated:

THEOREM 4.5.13 ([**215**], [**59**], [**141**]). *Let $\mathcal{K}$ be as in Theorem 4.5.12 and $n \geqslant 3$. Then the nonzero bigraded Betti numbers of $\mathcal{Z}_\mathcal{K}$ are given by*

$$\begin{aligned} b^{0,0}(\mathcal{Z}_\mathcal{K}) &= b^{-(m-n),2m}(\mathcal{Z}_\mathcal{K}) = 1,\\ b^{-i,2(i+1)}(\mathcal{Z}_\mathcal{K}) &= i\tbinom{m-n}{i+1} \quad \text{for } 1 \leqslant i \leqslant m-n-1,\\ b^{-i,2(i+n-1)}(\mathcal{Z}_\mathcal{K}) &= (m-n-i)\tbinom{m-n}{m-n+1-i} \quad \text{for } 1 \leqslant i \leqslant m-n-1. \end{aligned}$$

Exercises.

4.5.14. A triangulated manifold $\mathcal{K}$ is a pseudomanifold.

4.5.15. If $\mathcal{K}$ is a d-dimensional pseudomanifold, then either $H_d(\mathcal{K}) \cong \mathbb{Z}$ or $H_d(\mathcal{K}) \cong 0$. What happens for homology and cohomology with coefficients in a field $\mathbf{k}$?

4.5.16. If $\mathcal{K}$ is an orientable d-dimensional pseudomanifold (i.e. $H_d(\mathcal{K}) \cong \mathbb{Z}$) then $H^d(\mathcal{K}) \cong \mathbb{Z}$, and if $\mathcal{K}$ is non-orientable then $H^d(\mathcal{K}) \cong \mathbb{Z}_2$. What happens for cohomology with coefficients in a field $\mathbf{k}$?

4.5.17. Let $\mathcal{K}$ be an orientable d-dimensional pseudomanifold. The homology group $H_d(\mathcal{K}) \cong \mathbb{Z}$ is generated by the class of simplicial chain $\langle\mathcal{K}\rangle = \sum_{I\in\mathcal{K},\,\dim I=d} I$ where the d-simplices $I \in \mathcal{K}$ are oriented properly (the *fundamental homology class* of $\mathcal{K}$). The cohomology group $H^d(\mathcal{K}) \cong \mathbb{Z}$ is generated by the class of any cochain α_I taking value 1 on an oriented d-simplex $I \in \mathcal{K}$ and vanishing on all other simplices.

4.5.18. Calculate $H^*(\mathcal{Z}_\mathcal{K})$ where $\mathcal{K}$ is the boundary of a pentagon using Theorem 4.4.8 and the description of the cohomology product given in Proposition 3.2.10 (or Exercise 3.2.14).

4.6. Coordinate subspace arrangements

Here we establish a homotopy equivalence between the moment-angle complex $\mathcal{Z}_\mathcal{K}$ and the complement of the arrangement of coordinate subspaces in $\mathbb{C}^m$ corresponding to a simplicial complex $\mathcal{K}$. As a corollary we obtain an explicit description of the cohomology ring of a coordinate subspace arrangement complement. In some cases, knowing the cohomology ring allows us to identify the homotopy type of arrangement complements.

Coordinate subspace arrangements already appeared in Section 3.1 as affine algebraic varieties corresponding to face rings (see Proposition 3.1.12). Here we consider these arrangements from the general point of view.

A *coordinate subspace* in $\mathbb{C}^m$ can be given as

$$L_I = \{(z_1,\ldots,z_m) \in \mathbb{C}^m \colon z_{i_1} = \cdots = z_{i_k} = 0\}, \tag{4.18}$$

where $I = \{i_1,\ldots,i_k\}$ is a subset of $[m]$.

Construction 4.6.1. We assign to a simplicial complex $\mathcal{K}$ the arrangement of complex coordinate subspaces (or *coordinate subspace arrangement*) given by

$$\mathcal{A}(\mathcal{K}) = \{L_I \colon I \notin \mathcal{K}\}.$$

We denote by $U(\mathcal{K})$ the complement to $\mathcal{A}(\mathcal{K})$ in $\mathbb{C}^m$, that is,

$$U(\mathcal{K}) = \mathbb{C}^m \setminus \bigcup_{I \notin \mathcal{K}} L_I. \tag{4.19}$$

Observe that if $\mathcal{K}' \subset \mathcal{K}$ is a subcomplex, then $U(\mathcal{K}') \subset U(\mathcal{K})$.

Proposition 4.6.2. *The assignment $\mathcal{K} \mapsto U(\mathcal{K})$ defines a bijective inclusion-preserving correspondence between simplicial complexes on the set $[m]$ and complements of coordinate subspace arrangements in $\mathbb{C}^m$.*

Proof. We need to reconstruct a simplicial complex from the complement and check that it indeed defines the inverse correspondence. Let $\mathcal{A}$ be a coordinate subspace arrangement in $\mathbb{C}^m$, and let U be its complement. Set

$$\mathcal{K}(U) = \{I \subset [m] \colon L_I \cap U \neq \varnothing\}.$$

It is easy to see that $\mathcal{K}(U)$ is a simplicial complex satisfying $U(\mathcal{K}(U)) = U$ and $\mathcal{K}(U(\mathcal{K})) = \mathcal{K}$. □

If $\{i\}$ is a ghost vertex of $\mathcal{K}$, then the coordinate subspace arrangement $\mathcal{A}(\mathcal{K})$ contains the hyperplane $\{z_i = 0\}$. The arrangement $\mathcal{A}(\mathcal{K})$ does not contain hyperplanes if and only the vertex set of $\mathcal{K}$ is the whole $[m]$.

The complement $U(\mathcal{K})$ is an example of a polyhedral product space (see Construction 4.1.9), as is shown by the next proposition.

Proposition 4.6.3. $U(\mathcal{K}) = (\mathbb{C}, \mathbb{C}^\times)^\mathcal{K}$.

Proof. Given a point $\boldsymbol{z} = (z_1,\ldots,z_m) \in \mathbb{C}^m$, consider its zero set $\omega(\boldsymbol{z}) = \{i \in [m] \colon z_i = 0\} \subset [m]$. We have

$$\begin{aligned} U(\mathcal{K}) &= \mathbb{C}^m \setminus \bigcup_{I \notin \mathcal{K}} L_I = \mathbb{C}^m \setminus \bigcup_{I \notin \mathcal{K}} \{\boldsymbol{z} \colon \omega(\boldsymbol{z}) \supset I\} = \mathbb{C}^m \setminus \bigcup_{I \notin \mathcal{K}} \{\boldsymbol{z} \colon \omega(\boldsymbol{z}) = I\} \\ &= \bigcup_{I \in \mathcal{K}} \{\boldsymbol{z} \colon \omega(\boldsymbol{z}) = I\} = \bigcup_{I \in \mathcal{K}} \{\boldsymbol{z} \colon \omega(\boldsymbol{z}) \subset I\} = \bigcup_{I \in \mathcal{K}} (\mathbb{C}, \mathbb{C}^\times)^I = (\mathbb{C}, \mathbb{C}^\times)^\mathcal{K}. \quad \square \end{aligned}$$

Example 4.6.4.
1. If $\mathcal{K} = \Delta^{m-1}$, then $U(\mathcal{K}) = \mathbb{C}^m$.
2. If $\mathcal{K} = \partial\Delta^{m-1}$, then $U(\mathcal{K}) = \mathbb{C}^m \setminus \{0\}$.
3. Let $\mathcal{K}$ be the discrete complex with m vertices. Then

$$U(\mathcal{K}) = \mathbb{C}^m \setminus \bigcup_{1\leqslant i<j\leqslant m} \{z_i = z_j = 0\}$$

is the complement to all coordinate subspaces of codimension two.

4. More generally, if $\mathcal{K}$ is the i-dimensional skeleton of Δ^{m-1}, then $U(\mathcal{K})$ is the complement to all coordinate subspaces of codimension $i + 2$.

Since each coordinate subspace is invariant under the standard action of $\mathbb{T}^m$ on $\mathbb{C}^m$, the complement $U(\mathcal{K})$ is also a $\mathbb{T}^m$-invariant subset in $\mathbb{C}^m$.

A *deformation retraction* of a space X onto a subspace A is a homotopy $F_t\colon X \to X$, $t \in \mathbb{I}$, such that $F_0 = \mathrm{id}$ (the identity map), $F_1(X) = A$ and $F_t|_A = \mathrm{id}$ for all t. The term 'deformation retraction' is often used only for the last map $f = F_1\colon X \to A$; this map is a homotopy equivalence.

Theorem 4.6.5. *The moment-angle complex $\mathcal{Z}_\mathcal{K}$ is a $\mathbb{T}^m$-invariant subspace of $U(\mathcal{K})$, and there is a $\mathbb{T}^m$-equivariant deformation retraction*

$$\mathcal{Z}_\mathcal{K} \hookrightarrow U(\mathcal{K}) \xrightarrow{\simeq} \mathcal{Z}_\mathcal{K}.$$

Proof. We have $\mathcal{Z}_\mathcal{K} = (\mathbb{D},\mathbb{S})^\mathcal{K} \subset (\mathbb{C},\mathbb{C}^\times)^\mathcal{K} = U(\mathcal{K})$ by the functoriality of the polyhedral product, so the moment-angle complex $\mathcal{Z}_\mathcal{K}$ is indeed contained in the complement $U(\mathcal{K})$ as a $\mathbb{T}^m$-invariant subset.

The deformation retraction $U(\mathcal{K}) \to \mathcal{Z}_\mathcal{K}$ will be constructed by induction. We remove simplices from Δ^{m-1} until we obtain $\mathcal{K}$, in such a way that we get a simplicial complex at each intermediate step.

The base of induction is clear: if $\mathcal{K} = \Delta^{m-1}$, then $U(\mathcal{K}) = \mathbb{C}^m$, $\mathcal{Z}_\mathcal{K} = \mathbb{D}^m$, and the retraction $\mathbb{C}^m \to \mathbb{D}^m$ is evident.

The orbit space $\mathcal{Z}_\mathcal{K}/\mathbb{T}^m$ is the cubical complex $\mathrm{cc}(\mathcal{K}) = (\mathbb{I}, 1)^\mathcal{K}$ (see Construction 2.9.11). The orbit space $U(\mathcal{K})/\mathbb{T}^m$ can be identified with

$$U(\mathcal{K})_\geqslant = U(\mathcal{K}) \cap \mathbb{R}^m_\geqslant = (\mathbb{R}_\geqslant, \mathbb{R}_>)^\mathcal{K}$$

where $\mathbb{R}^m_\geqslant$ is viewed as a subset in $\mathbb{C}^m$.

We shall first construct a deformation retraction $r\colon U(\mathcal{K})_\geqslant \to \mathrm{cc}(\mathcal{K})$ of orbit spaces, and then cover it by a deformation retraction $\widetilde{r}\colon U(\mathcal{K}) \to \mathcal{Z}_\mathcal{K}$.

Now assume that $\mathcal{K}$ is obtained from a simplicial complex $\mathcal{K}'$ by removing one maximal simplex $J = \{j_1, \ldots, j_k\}$, i.e. $\mathcal{K} \cup J = \mathcal{K}'$. Then the cubical complex $\mathrm{cc}(\mathcal{K}')$ is obtained from $\mathrm{cc}(\mathcal{K})$ by adding a single k-dimensional face $C_J = (\mathbb{I}, 1)^J$. We also have $U(\mathcal{K}) = U(\mathcal{K}') \setminus L_J$, so that

$$U(\mathcal{K})_\geqslant = U(\mathcal{K}')_\geqslant \setminus \{\boldsymbol{y}\colon y_{j_1} = \cdots = y_{j_k} = 0\}.$$

We may assume by induction that there is a deformation retraction $r'\colon U(\mathcal{K}')_\geqslant \to \mathrm{cc}(\mathcal{K}')$ such that $\omega(r'(\boldsymbol{y})) = \omega(\boldsymbol{y})$, where $\omega(\boldsymbol{y})$ is the set of zero coordinates of $\boldsymbol{y}$. In particular, r' restricts to a deformation retraction

$$r'\colon U(\mathcal{K}')_\geqslant \setminus \{\boldsymbol{y}\colon y_{j_1} = \cdots = y_{j_k} = 0\} \longrightarrow \mathrm{cc}(\mathcal{K}') \setminus \boldsymbol{y}_J$$

where $\boldsymbol{y}_J$ is the point with coordinates $y_{j_1} = \cdots = y_{j_k} = 0$ and $y_j = 1$ for $j \notin J$.

Since $J \notin \mathcal{K}$, we have $\boldsymbol{y}_J \notin \mathrm{cc}(\mathcal{K})$. On the other hand, $\boldsymbol{y}_J$ belongs to the extra face $C_J = (\mathbb{I}, 1)^J$ of $\mathrm{cc}(\mathcal{K}')$. We therefore may apply the deformation retraction r_J

shown in Fig. 4.2 on the face C_J, with centre at $\boldsymbol{y}_J$. In coordinates, a homotopy F_t



FIGURE 4.2. Retraction $r_J\colon\ \mathrm{cc}(\mathcal{K}')\setminus \boldsymbol{y}_J \to \mathrm{cc}(\mathcal{K})$.

between the identity map $\mathrm{cc}(\mathcal{K}')\setminus \boldsymbol{y}_J \to \mathrm{cc}(\mathcal{K}')\setminus \boldsymbol{y}_J$ (for $t=0$) and the retraction $r_J\colon\ \mathrm{cc}(\mathcal{K}')\setminus \boldsymbol{y}_J \to \mathrm{cc}(\mathcal{K})$ (for $t=1$) is given by

$$\begin{aligned} F_t\colon\ \mathrm{cc}(\mathcal{K}')\setminus \boldsymbol{y}_J &\longrightarrow \mathrm{cc}(\mathcal{K}')\setminus \boldsymbol{y}_J,\\ (y_1,\dots,y_m,t) &\longmapsto (y_1+t\alpha_1y_1,\dots,y_m+t\alpha_my_m). \end{aligned}$$

where

$$\alpha_i=\begin{cases}\frac{1-\max_{j\in J}y_j}{\max_{j\in J}y_j}, & \text{if } i\in J,\\ 0, & \text{if } i\notin J,\end{cases}\qquad \text{for } 1\leqslant i\leqslant m.$$

We observe that $\omega(F_t(\boldsymbol{y}))=\omega(\boldsymbol{y})$ for any t and $\boldsymbol{y}\in\mathrm{cc}(\mathcal{K}')$. Now, the composition

$$r\colon U(\mathcal{K})_{\geqslant}=U(\mathcal{K}')_{\geqslant}\setminus\{\boldsymbol{y}\colon y_{j_1}=\dots=y_{j_k}=0\}\xrightarrow{r'}\mathrm{cc}(\mathcal{K}')\setminus \boldsymbol{y}_J\xrightarrow{r_J}\mathrm{cc}(\mathcal{K}) \tag{4.20}$$

is a deformation retraction, and it satisfies $\omega(r(\boldsymbol{y}))=\omega(\boldsymbol{y})$ as this is true for r_J and r. The inductive step is now complete. The required retraction $\widetilde{r}\colon U(\mathcal{K})\to\mathcal{Z}_{\mathcal{K}}$ covers r as shown in the following commutative diagram:

$$\begin{tikzcd}
\mathcal{Z}_{\mathcal{K}} \arrow[r, hook] \arrow[d, "\mu"] & U(\mathcal{K}) \arrow[r, "\widetilde{r}"] \arrow[d, "\mu"] & \mathcal{Z}_{\mathcal{K}} \arrow[d, "\mu"] \\
\mathrm{cc}(\mathcal{K}) \arrow[r, hook] & U_{\geqslant}(\mathcal{K}) \arrow[r, "r"] & \mathrm{cc}(\mathcal{K})
\end{tikzcd}$$

Explicitly, $\widetilde{r}$ is decomposed inductively in a way similar to (4.20),

$$\widetilde{r}\colon U(\mathcal{K})=U(\mathcal{K}')\setminus L_J\xrightarrow{\widetilde{r}'}\mathcal{Z}_{\mathcal{K}'}\setminus\mu^{-1}(\boldsymbol{y}_J)\xrightarrow{\widetilde{r}_J}\mathcal{Z}_{\mathcal{K}},$$

where $\mu^{-1}(\boldsymbol{y}_J)=\prod_{j\in J}\{0\}\times\prod_{j\notin J}\mathbb{S}$, and $\widetilde{r}_J$ is given in coordinates $(z_1,\dots,z_m)=\bigl(\sqrt{y_1}e^{i\varphi_1},\dots,\sqrt{y_m}e^{i\varphi_m}\bigr)$ by

$$\bigl(\sqrt{y_1}e^{i\varphi_1},\dots,\sqrt{y_m}e^{i\varphi_m}\bigr)\mapsto\bigl(\sqrt{y_1+\alpha_1y_1}e^{i\varphi_1},\dots,\sqrt{y_m+\alpha_my_m}e^{i\varphi_m}\bigr)$$

with α_i as above. □

Since $U(\mathcal{K})$ and $\mathcal{Z}_{\mathcal{K}}$ are homotopy equivalent, we can use the results on the cohomology of $\mathcal{Z}_{\mathcal{K}}$ (such as Theorems 4.4.4 and 4.4.8) to describe the cohomology rings of coordinate subspace arrangement complements.

Example 4.6.6. Let $\mathcal{K}$ be the set of m disjoint points. Then $\mathcal{Z}_{\mathcal{K}}$ is homotopy equivalent to the complement $U(\mathcal{K})$ of Example 4.6.4.3, and

$$\mathbb{Z}[\mathcal{K}] = \mathbb{Z}[v_1, \dots, v_m]/(v_i v_j,\ i \neq j).$$

The subspace of cocycles in $R^*(\mathcal{K})$ has basis of monomials

$$u_{i_1} u_{i_2} \cdots u_{i_{k-1}} v_{i_k} \quad \text{with } i_p \neq i_q \text{ for } p \neq q.$$

Since the total degree of $u_{i_1} u_{i_2} \cdots u_{i_{k-1}} v_{i_k}$ is $k+1$, the space of cocycles of degree $k+1$ has dimension $m\binom{m-1}{k-1}$. The subspace of coboundaries of degree $k+1$ is spanned by the elements of the form

$$d(u_{i_1} \cdots u_{i_k})$$

and has dimension $\binom{m}{k}$. Therefore,

$$\begin{aligned}
&\dim H^0(U(\mathcal{K})) = 1,\\
&\dim H^1(U(\mathcal{K})) = H^2(U(\mathcal{K})) = 0,\\
&\dim H^{k+1}(U(\mathcal{K})) = m\tbinom{m-1}{k-1} - \tbinom{m}{k} = (k-1)\tbinom{m}{k}, \quad \text{for } 2 \leqslant k \leqslant m,
\end{aligned}$$

and the multiplication in the cohomology of $U(\mathcal{K})$ is trivial.

The calculation of the previous example shows that if $\mathcal{K}$ is the set of m points, then there is a cohomology ring isomorphism

$$H^*\bigl(U(\mathcal{K})\bigr) \cong H^*\Bigl(\bigvee_{k=2}^{m} \bigl(S^{k+1}\bigr)^{\vee(k-1)\binom{m}{k}}\Bigr), \tag{4.21}$$

where $X^{\vee k}$ denotes the k-fold wedge of a space X. This cohomology isomorphism is induced by a homotopy equivalence, as is shown by the following result.

Theorem 4.6.7 ([**103**], [**104**, Corollary 9.5]). *Let $\mathcal{K}$ be the i-dimensional skeleton of the simplex Δ^{m-1}, so that $U(\mathcal{K})$ is the complement to all coordinate planes of codimension $i+2$ in $\mathbb{C}^m$. Then $U(\mathcal{K})$ is homotopy equivalent to a wedge of spheres:*

$$U(\mathcal{K}) \simeq \bigvee_{k=i+2}^{m} \bigl(S^{i+k+1}\bigr)^{\vee\binom{m}{k}\binom{k-1}{i+1}}.$$

The proof uses the interpretation of $\mathcal{Z}_{\mathcal{K}}$ as the homotopy fibre of the inclusion $i\colon (\mathbb{C}P^\infty)^{\mathcal{K}} \hookrightarrow (\mathbb{C}P^\infty)^m$ (Theorem 4.2.2), and will be given in Section 8.3. For $i = 0$ we obtain the homotopy equivalence behind cohomology isomorphism (4.21).

Real coordinate subspace arrangements

$$U_{\mathbb{R}}(\mathcal{K}) = (\mathbb{R}, \mathbb{R}^\times)^{\mathcal{K}}$$

in $\mathbb{R}^m$ are defined and treated similarly; there are real analogues of all results and constructions of this section. (The real version of Theorem 4.6.7 is much simpler: see Exercise 4.1.16).

A coordinate subspace can be given either by setting some coordinates to zero, as in (4.18), or as the linear span of a subset of the standard basis $\boldsymbol{e}_1, \dots, \boldsymbol{e}_m$. The latter approach leads to an alternative way of parametrising complements of coordinate subspace arrangements by simplicial complexes, which is related to the former one by Alexander duality.

Given a subset $I \subset [m]$ we set $S_I = \mathbb{C}\langle \boldsymbol{e}_i\colon i \in I\rangle$ (the $\mathbb{C}$-span of the basis vectors corresponding to I), and use the notation $\widehat{I} = [m] \setminus I$ and $\widehat{\mathcal{K}} = \{\widehat{I} \subset [m]\colon I \notin \mathcal{K}\}$

from Construction 2.4.1. Then the coordinate subspace arrangement corresponding to a simplicial complex $\mathcal{K}$ can be written in the following two ways:

$$\mathcal{A}(\mathcal{K}) = \{L_I \colon I \notin \mathcal{K}\} = \{S_{\widehat{I}} \colon \widehat{I} \in \widehat{\mathcal{K}}\}.$$

Using Alexander duality we can reformulate the description of the cohomology of $U(\mathcal{K})$ in terms of full subcomplexes of $\mathcal{K}$ (Theorem 4.4.8) as follows.

Proposition 4.6.8. *There are isomorphisms*

$$\widetilde{H}^q\bigl(U(\mathcal{K})\bigr) \cong \bigoplus_{\widehat{I}\in\widehat{\mathcal{K}}} \widetilde{H}_{2m-2|\widehat{I}|-q-2}(\operatorname{lk}_{\widehat{\mathcal{K}}} \widehat{I}).$$

Proof. By Theorems 4.6.5 and 4.4.8,

$$H^q\bigl(U(\mathcal{K})\bigr) \cong \bigoplus_{I\subset[m]} \widetilde{H}^{q-|I|-1}(\mathcal{K}_I).$$

Nonempty simplices $I \in \mathcal{K}$ do not contribute to the sum above, since the corresponding subcomplexes $\mathcal{K}_I$ are contractible. Since $\widetilde{H}^{-1}(\varnothing) = \mathbb{Z}$, the empty simplex contributes $\mathbb{Z}$ to $H^0(U(\mathcal{K}))$. Therefore, we can rewrite the isomorphism above as

$$\widetilde{H}^q\bigl(U(\mathcal{K})\bigr)\cong \bigoplus_{I\notin\mathcal{K}} \widetilde{H}^{q-|I|-1}(\mathcal{K}_I).$$

Using Alexander duality (Proposition 2.4.6) we calculate

$$\widetilde{H}^{q-|I|-1}(\mathcal{K}_I) \cong \widetilde{H}_{|I|-3-q+|I|+1}(\operatorname{lk}_{\widehat{\mathcal{K}}} \widehat{I}) = \widetilde{H}_{2m-2|\widehat{I}|-q-2}(\operatorname{lk}_{\widehat{\mathcal{K}}} \widehat{I}),$$

where $\widehat{I} = [m]\backslash I$ is a simplex of $\widehat{\mathcal{K}}$. □

Proposition 4.6.8 is a particular case of the well-known *Goresky–MacPherson formula* [**102**, Chapter III], which calculates the (co)homology groups of the complement of an arrangement of affine subspaces in terms of its *intersection poset*. In the case of coordinate subspace arrangements $\mathcal{A}(\mathcal{K})$ the intersection poset is the face poset of the dual complex $\widehat{\mathcal{K}}$. For more on the relationships between general affine subspace arrangements and moment-angle complexes see [**45**, Chapter 8].

Exercises.

4.6.9. The affine algebraic variety $X(\mathcal{K})$ corresponding to the face ring $\mathbb{C}[\mathcal{K}]$ (see Proposition 3.1.12) and the coordinate subspace arrangement $\mathcal{A}(\mathcal{K})$ of Construction 4.6.1 are related by the identity $X(\widehat{\mathcal{K}}) = \mathcal{A}(\mathcal{K})$, where $\widehat{\mathcal{K}} = \{I \subset [m] \colon [m] \setminus I \notin \mathcal{K}\}$ is the Alexander dual complex.

4.6.10. Show directly that the complement to the 3 coordinate lines in $\mathbb{C}^3$ is homotopy equivalent to the wedge of spheres $S^3\vee S^3\vee S^3\vee S^4\vee S^4$; this corresponds to $m = 3$ in (4.21).

4.6.11. Show directly (without referring to Theorem 4.6.5 and Proposition 4.2.4) that the complement $U(\mathcal{K})$ is 2-connected if $\mathcal{K}$ does not have ghost vertices, and that $U(\mathcal{K})$ is $2q$-connected if $\mathcal{K}$ is q-neighbourly.

Moment-angle complexes: additional topics

4.7. Free and almost free torus actions on $\mathcal{Z}_\mathcal{K}$

Here consider free and almost free actions of toric subgroups $T^k \subset \mathbb{T}^m$ on $\mathcal{Z}_\mathcal{K}$. As usual, $\mathcal{K}$ is an $(n-1)$-dimensional simplicial complex on $[m]$, and $\mathcal{Z}_\mathcal{K}$ is the corresponding moment-angle complex.

We start with a simple characterisation of the isotropy subgroups of the standard $\mathbb{T}^m$-action on $\mathcal{Z}_\mathcal{K}$. For each $I \subset [m]$ we consider the coordinate subtorus

$$\mathbb{T}^I = \{(t_1, \dots, t_m) \in \mathbb{T}^m : t_j = 1 \text{ for } j \notin I\} = \prod_{i \in I} \mathbb{T} \;\subset\; \mathbb{T}^m$$

(note that $\mathbb{T}^I = (\mathbb{T}, 1)^I$ in the notation of Construction 4.1.9).

Proposition 4.7.1. *Let $\boldsymbol{z} \in \mathcal{Z}_\mathcal{K}$, and set $\omega(\boldsymbol{z}) = \{i \in [m] : z_i = 0\} \in \mathcal{K}$. Then the isotropy subgroup of $\boldsymbol{z}$ with respect to the $\mathbb{T}^m$-action is $\mathbb{T}^{\omega(\boldsymbol{z})}$. Furthermore, each coordinate subtorus $\mathbb{T}^I$ for $I \in \mathcal{K}$ is the isotropy subgroup for a point $\boldsymbol{z} \in \mathcal{Z}_\mathcal{K}$.*

Proof. An element $\boldsymbol{t} = (t_1, \dots, t_m) \in \mathbb{T}^m$ fixes $\boldsymbol{z}$ if and only if $t_i = 1$ whenever $z_i \neq 0$, which is equivalent to that $\boldsymbol{t} \in \mathbb{T}^{\omega(\boldsymbol{z})}$. The last statement is also clear: $\mathbb{T}^I$ is the isotropy subgroup for any $\boldsymbol{z} \in (\mathbb{D}, \mathbb{S})^I \subset \mathcal{Z}_\mathcal{K}$ with $\omega(\boldsymbol{z}) = I$. □

Recall that an action of a group on a topological space is *almost free* if all isotropy subgroups are finite.

Definition 4.7.2. We define the *free toral rank* of $\mathcal{Z}_\mathcal{K}$, denoted $\operatorname{ftr} \mathcal{Z}_\mathcal{K}$, as the maximal dimension of toric subgroups $T^k \subset \mathbb{T}^m$ acting on $\mathcal{Z}_\mathcal{K}$ freely. Similarly, the *almost free toral rank* of $\mathcal{Z}_\mathcal{K}$, denoted $\operatorname{atr} \mathcal{Z}_\mathcal{K}$, is the maximal dimension of toric subgroups $T^k \subset \mathbb{T}^m$ acting on $\mathcal{Z}_\mathcal{K}$ almost freely.

Proposition 4.7.3. *Let $\mathcal{K}$ be a simplicial complex of dimension $n-1$ on m vertices and $\mathcal{K} \neq \Delta^{m-1}$. The toral ranks of $\mathcal{Z}_\mathcal{K}$ satisfy the following inequalities:*

$$1 \leqslant \operatorname{ftr} \mathcal{Z}_\mathcal{K} \leqslant \operatorname{atr} \mathcal{Z}_\mathcal{K} \leqslant m - n.$$

Proof. By Proposition 4.7.1, isotropy subgroups for the $\mathbb{T}^m$-action on $\mathcal{Z}_\mathcal{K}$ are coordinate subgroups of the form $\mathbb{T}^I$. The diagonal circle in $\mathbb{T}^m$ intersects each of these coordinate subgroups trivially (since $I \neq [m]$), and therefore acts freely on $\mathcal{Z}_\mathcal{K}$. This proves the first inequality. The second is obvious. To prove the third one, assume that $T^k \subset \mathbb{T}^m$ acts almost freely on $\mathcal{Z}_\mathcal{K}$. Then the intersection of T^k with every $\mathbb{T}^m$-isotropy subgroup $\mathbb{T}^I$ is a finite group. Choose a maximal simplex $I \in \mathcal{K}$, $|I| = n$. Then $\mathbb{T}^I \cap T^k$ can be finite only if $k \leqslant m - n$. □

The map $\mathbb{R}^m \to \mathbb{T}^m$, $(\varphi_1, \dots, \varphi_m) \mapsto (e^{2\pi i\varphi_1}, \dots, e^{2\pi i\varphi_m})$, identifies $\mathbb{T}^m$ with the quotient $\mathbb{R}^m/\mathbb{Z}^m$. Subtori $T^k \subset \mathbb{T}^m$ of dimension k bijectively correspond to unimodular sublattices $L \subset \mathbb{Z}^m$ of rank k (a sublattice is *unimodular* if it is a direct summand in $\mathbb{Z}^m$). The inclusion $T^k \subset \mathbb{T}^m$ can be viewed as $L_{\mathbb{R}}/L \subset \mathbb{R}^m/\mathbb{Z}^m$, where $L_{\mathbb{R}}$ is the k-dimensional subspace in $\mathbb{R}^m$ spanned by L.

Choosing a basis in L we obtain an integer $m \times k$-matrix $S = (s_{ij})$, $1 \leqslant i \leqslant m$, $1 \leqslant j \leqslant k$, so that L is identified with the image of $S\colon \mathbb{Z}^k \to \mathbb{Z}^m$. The k-torus T^k is the image of the corresponding monomorphism of tori $\mathbb{T}^k \to \mathbb{T}^m$, namely,

$$T^k = \bigl\{\bigl(e^{2\pi i(s_{11}\psi_1+\cdots+s_{1k}\psi_k)}, \ldots, e^{2\pi i(s_{m1}\psi_1+\cdots+s_{mk}\psi_k)}\bigr)\bigr\} \subset \mathbb{T}^m, \tag{4.22}$$

where $(\psi_1, \ldots, \psi_k) \in \mathbb{R}^k$. Since L is unimodular, the columns of S form a part of basis of the lattice $\mathbb{Z}^m$.

LEMMA 4.7.4. *Let T^k be a k-dimensional subtorus in $\mathbb{T}^m$ and let L be the corresponding unimodular sublattice of rank k in $\mathbb{Z}^m$. Let $S = (s_{ij})$, $1 \leqslant i \leqslant m$, $1 \leqslant j \leqslant k$, be a matrix defining L, so that T^k is given by* (4.22).

(a) *The torus T^k acts on $\mathcal{Z}_{\mathcal{K}}$ almost freely if and only if for each $I \in \mathcal{K}$ the intersection of subspaces $L_{\mathbb{R}}$ and $\mathbb{R}^I$ in $\mathbb{R}^m$ is zero. Equivalently, the $(m-|I|) \times k$-matrix $S_{\widehat{I}}$ obtained by deleting from S the rows with numbers $i \in I$ has rank k.*

(b) *The torus T^k acts on $\mathcal{Z}_{\mathcal{K}}$ freely if and only if for each $I \in \mathcal{K}$ the sublattice spanned by L and $\mathbb{Z}^I$ in $\mathbb{Z}^m$ is unimodular of rank $k + |I|$. Equivalently, the columns of the $(m-|I|) \times k$-matrix $S_{\widehat{I}}$ form a part of basis of $\mathbb{Z}^{m-|I|}$.*

PROOF. We prove (a) first. By Proposition 4.7.1, the T^k-action on $\mathcal{Z}_{\mathcal{K}}$ is almost free if and only if the intersection $T^k \cap \mathbb{T}^I \subset \mathbb{T}^m$ is finite for each $I \in \mathcal{K}$. This intersection can be identified with the kernel of the map $f\colon T^k \times \mathbb{T}^I \to \mathbb{T}^m$ (the product of the inclusion maps $T^k \to \mathbb{T}^m$ and $\mathbb{T}^I \to \mathbb{T}^m$). This kernel is finite if and only if the corresponding map of real spaces $L_{\mathbb{R}} \times \mathbb{R}^I \to \mathbb{R}^m$ is injective, which is equivalent to that $L_{\mathbb{R}} \cap \mathbb{R}^I = \{0\}$. Let $I = \{i_1, \ldots, i_p\}$, then the matrix of f has the form $(S\,|\,\boldsymbol{e}_{i_1}|\cdots|\boldsymbol{e}_{i_p})$, where $\boldsymbol{e}_i$ is the ith standard basis column vector. Clearly, this matrix has rank $k + |I|$ if and only if the matrix $S_{\widehat{I}}$ has rank k.

Now we prove (b). The T^k-action on $\mathcal{Z}_{\mathcal{K}}$ is free if and only if the the kernel of $f\colon T^k \times \mathbb{T}^I \to \mathbb{T}^m$ is trivial for each $I \in \mathcal{K}$, i.e. $T^k \times \mathbb{T}^I$ embeds as a subtorus. This is equivalent to the conditions stated in (b). □

The following algebraic lemma will be required for the algebraic characterisation of maximal-dimensional subtori acting on $\mathcal{Z}_{\mathcal{K}}$ freely or almost freely:

LEMMA 4.7.5. *Let $\mathbf{k}$ be a field or $\mathbb{Z}$, and assume given a diagram*

$$\begin{tikzcd}
 & & 0 \arrow[d] & & \\
 & & \mathbf{k}^r \arrow[d, "i_1"] & & \\
0 \arrow[r] & \mathbf{k}^s \arrow[r, "i_2"] & \mathbf{k}^m \arrow[r, "p_2"] \arrow[d, "p_1"] & \mathbf{k}^{m-s} \arrow[r] & 0 \\
 & & \mathbf{k}^{m-r} \arrow[d] & & \\
 & & 0 & &
\end{tikzcd}$$

in which both vertical and horizontal lines are short exact sequences of $\mathbf{k}$*-vector spaces or free abelian groups. Then* p_1i_2 *is surjective (respectively, injective or split injective) if and only if* p_2i_1 *is surjective (respectively, injective or split injective).*

Proof. This is a simple diagram chase. Assume first that p_1i_2 is surjective. Take $\alpha \in \mathbf{k}^{m-s}$; we need to cover it by an element in $\mathbf{k}^r$. There is $\beta \in \mathbf{k}^m$ such that $p_2(\beta) = \alpha$. If $\beta \in i_1(\mathbf{k}^r)$, then we are done. Otherwise set $\gamma = p_1(\beta) \neq 0$. Since p_1i_2 is surjective, we can choose $\delta \in \mathbf{k}^s$ such that $p_1i_2(\delta) = \gamma$. Set $\eta = i_2(\delta) \neq 0$. Hence, $p_1(\eta) = p_1(\beta)(= \gamma)$ and there is $\xi \in \mathbf{k}^r$ such that $i_1(\xi) = \beta - \eta$. Then $p_2i_1(\xi) = p_2(\beta - \eta) = \alpha - p_2i_2(\delta) = \alpha$. Thus, p_2i_1 is surjective.

Now assume that p_1i_2 is injective. Suppose $p_2i_1(\alpha) = 0$ for a nonzero $\alpha \in \mathbf{k}^r$. Set $\beta = i_1(\alpha) \neq 0$. Since $p_2(\beta) = 0$, there is a nonzero $\gamma \in \mathbf{k}^s$ such that $i_2(\gamma) = \beta$. Then $p_1i_2(\gamma) = p_1(\beta) = p_1i_1(\alpha) = 0$. This contradicts the assumption that p_1i_2 is injective. Thus, p_2i_1 is injective.

Finally, if p_1i_2 is split injective, then its dual map $i_2^*p_1^*\colon \mathbf{k}^{m-r} \to \mathbf{k}^s$ is surjective. Then $i_1^*p_2^*\colon \mathbf{k}^{m-s} \to \mathbf{k}^r$ is also surjective. Thus, p_2i_1 is split injective. □

Let $\boldsymbol{t} = (t_1, \ldots, t_n) \in \mathbb{Z}[\mathcal{K}]$ be a linear sequence given by

$$t_i = \lambda_{i1}v_1 + \cdots + \lambda_{im}v_m, \quad \text{for } 1 \leqslant i \leqslant n. \tag{4.23}$$

We consider the integer $n \times m$-matrix $\Lambda = (\lambda_{ij})$, $1 \leqslant i \leqslant n$, $1 \leqslant j \leqslant m$. It defines a homomorphism of lattices $\Lambda\colon \mathbb{Z}^m \to \mathbb{Z}^n$ and a homomorphism of tori $\Lambda\colon \mathbb{T}^m \to \mathbb{T}^n$.

Theorem 4.7.6. *The following conditions are equivalent:*

(a) *the sequence* $(t_1, \ldots, t_n)$ *given by* (4.23) *is an lsop in the rational face ring* $\mathbb{Q}[\mathcal{K}]$*;*

(b) *the kernel* $T_\Lambda = \operatorname{Ker}(\Lambda\colon \mathbb{T}^m \to \mathbb{T}^n)$ *is a product of an* $(m-n)$*-torus and a finite group, and* T_Λ *acts almost freely on* $\mathcal{Z}_{\mathcal{K}}$*.*

Proof. We first observe that under any of conditions (a) or (b) the rational map $\Lambda\colon \mathbb{Q}^m \to \mathbb{Q}^n$ is surjective. For each simplex $I \in \mathcal{K}$ we consider the diagram

$$\begin{tikzcd}
 & & 0 \arrow[d] & & \\
 & & \mathbb{Q}^{m-n} \arrow[d, "i_1"] & & \\
0 \arrow[r] & \mathbb{Q}^{|I|} \arrow[r, "i_2"] & \mathbb{Q}^m \arrow[r, "p_2"] \arrow[d, "\Lambda"] & \mathbb{Q}^{m-|I|} \arrow[r] & 0 \\
 & & \mathbb{Q}^n \arrow[d] & & \\
 & & 0 & &
\end{tikzcd}$$

where i_2 is the inclusion of the coordinate subspace $\mathbb{Q}^I \to \mathbb{Q}^m$. The map Λi_2 is given by the $n \times |I|$ matrix $\Lambda_I = (\lambda_{ij})$, $1 \leqslant i \leqslant n$, $j \in I$. By Lemma 3.3.2, the sequence $(t_1, \ldots, t_n)$ is a rational lsop if and only if the rank of Λ_I is $|I|$ for each $I \in \mathcal{K}$. Hence condition (a) of the theorem is equivalent to the injectivity of the map Λi_2 for each $I \in \mathcal{K}$.

On the other hand, by Lemma 4.7.4 (a), the action of the $(m-n)$-torus T_Λ on $\mathcal{Z}_{\mathcal{K}}$ is almost free if and only if $i_1(\mathbb{Q}^{m-n}) \cap \mathbb{Q}^I = \{0\}$ for each $I \in \mathcal{K}$. The latter condition is equivalent to that $i_1(\mathbb{Q}^{m-n}) \cap \operatorname{Ker} p_2 = \{0\}$, i.e. that $p_2 i_1$ is injective. Hence condition (b) of the theorem is equivalent to the injectivity of the map $p_2 i_1$ for each $I \in \mathcal{K}$. Now the theorem follows from Lemma 4.7.5. □

The almost free toral rank of $\mathcal{Z}_\mathcal{K}$ can now be easily determined:

COROLLARY 4.7.7. *We have* $\operatorname{atr} \mathcal{Z}_\mathcal{K} = m - n$, *i.e. for each simplicial complex* $\mathcal{K}$ *of dimension* $(n-1)$ *on* $[m]$ *there is an* $(m-n)$*-dimensional subtorus in* $\mathbb{T}^m$ *acting on* $\mathcal{Z}_\mathcal{K}$ *almost freely.*

PROOF. Choose a rational lsop in $\mathbb{Q}[\mathcal{K}]$ by Theorem A.3.10, and multiply it by a common denominator to get an integral sequence (4.23). It is still an lsop in $\mathbb{Q}[\mathcal{K}]$ (but it may fail to be an lsop in $\mathbb{Z}[\mathcal{K}]$), and therefore the $(m-n)$-torus T_Λ acts on $\mathcal{Z}_\mathcal{K}$ almost freely by Theorem 4.7.6. □

There is an analogue of Theorem 4.7.6 for free torus actions:

THEOREM 4.7.8. *The following conditions are equivalent:*

(a) *the sequence* $(t_1, \ldots, t_n)$ *given by* (4.23) *is an lsop in* $\mathbb{Z}[\mathcal{K}]$*;*
(b) $T_\Lambda = \operatorname{Ker}(\Lambda\colon \mathbb{T}^m \to \mathbb{T}^n)$ *is an* $(m-n)$*-torus acting freely on* $\mathcal{Z}_\mathcal{K}$*.*

PROOF. The argument is the same as in the proof of Theorem 4.7.6: Lemma 4.7.5 is now applied to the diagram of integral lattices instead of rational vector spaces. □

Nevertheless, there is no analogue of Corollary 4.7.7 for free torus actions, as integral lsop's may fail to exist. Indeed, the free toral rank of $\mathcal{Z}_\mathcal{K}$ where $\mathcal{K}$ is the boundary of a cyclic n-polytope with $m \geqslant 2^n$ vertices is strictly less than $m-n$, as shown by Example 3.3.4. The free toral rank of $\mathcal{Z}_\mathcal{K}$ is a combinatorial characteristic of $\mathcal{K}$, also known as the *Buchstaber invariant*. Its determination is a very subtle problem; see [**85**] and [**95**] for some partial results in this direction.

There is the following important conjecture of equivariant topology and rational homotopy theory concerning almost free torus action.

CONJECTURE 4.7.9 (Toral Rank Conjecture, Halperin [**114**]). *Assume that a torus* T^k *acts almost freely on a finite-dimensional topological space* X*. Then*

$$\dim H^*(X;\mathbb{Q}) \geqslant 2^k,$$

i.e. the total dimension of the cohomology of X *is at least that of the torus* T^k*.*

The Toral Rank Conjecture is valid for $k \leqslant 3$ and is open in general. See [**190**] and [**222**] for the discussion of the current status of this conjecture.

In the case of moment-angle complexes we have the following result:

THEOREM 4.7.10 ([**56**], [**221**]). *Let* $\mathcal{K}$ *be a simplicial complex of dimension* $n-1$ *with* m *vertices, and let* $\mathcal{Z}_\mathcal{K}$ *be the corresponding moment-angle complex. Then*

$$\operatorname{rank} H^*(\mathcal{Z}_\mathcal{K}) = \sum_{k=0}^{m+n} \operatorname{rank} H^k(\mathcal{Z}_\mathcal{K}) \geqslant 2^{m-n}.$$

The proof of this theorem given in [**221**] uses a construction of independent interest and a couple of technical lemmata. We include this proof below.

COROLLARY 4.7.11. *The Toral Rank Conjecture is valid for subtori* $T^k \subset \mathbb{T}^m$ *acting almost freely on* $\mathcal{Z}_\mathcal{K}$*.*

The following is a particular case of the so-called *simplicial wedge construction* [**189**]. It has been brought into toric topology by the work [**12**].

CONSTRUCTION 4.7.12 (Simplicial doubling). Let $\mathcal{K}$ be a simplicial complex on the vertex set $[m]$. The *double* of $\mathcal{K}$ is the simplicial complex $\mathcal{D}(\mathcal{K})$ on the vertex set $[2m] = \{1, 1', 2, 2', \dots, m, m'\}$ whose missing faces (minimal non-faces) are $\{i_1, i_1', \dots, i_k, i_k'\}$ where $\{i_1, \dots, i_k\}$ is a missing face of $\mathcal{K}$. In other words, $\mathcal{D}(\mathcal{K})$ is determined by its face ring given by

$$\mathbf{k}\bigl[\mathcal{D}(\mathcal{K})\bigr] = \mathbf{k}[v_1, v_{1'}, \dots, v_m, v_{m'}]/\bigl(\{v_{i_1}v_{i_1'}\cdots v_{i_k}v_{i_k'}\}\colon \{i_1, \dots, i_k\} \notin \mathcal{K}\bigr).$$

EXAMPLE 4.7.13.
1. If $\mathcal{K} = \Delta^{m-1}$ (the full simplex on m vertices), then $\mathcal{D}(\mathcal{K}) = \Delta^{2m-1}$.
2. If $\mathcal{K} = \partial\Delta^{m-1}$, then $\mathcal{D}(\mathcal{K}) = \partial\Delta^{2m-1}$.

The doubling construction interacts nicely with the polyhedral product:

THEOREM 4.7.14. *Let (X, A) be a pair of spaces, let $\mathcal{K}$ be a simplicial complex on $[m]$ and $\mathcal{D}(\mathcal{K})$ its double. Then*

$$(X, A)^{\mathcal{D}(\mathcal{K})} = (X \times X, X \times A \cup A \times X)^{\mathcal{K}}.$$

PROOF. Set $(Y, B) = (X \times X, X \times A \cup A \times X)$. Given a point $\boldsymbol{y} = (y_1, \dots, y_m) \in Y^m$, we set

$$\omega_Y(\boldsymbol{y}) = \{i \in [m]\colon y_i \notin B\} \subset [m].$$

Similarly, given $\boldsymbol{x} = (x_1, x_{1'}, \dots, x_m, x_{m'}) \in X^{2m}$, we set

$$\omega_X(\boldsymbol{x}) = \bigl\{j \in \{1, 1', \dots, m, m'\}\colon x_j \notin A\bigr\} \subset \{1, 1', \dots, m, m'\}.$$

We identify $\boldsymbol{y}$ with $\boldsymbol{x}$ by the formula $(y_1, \dots, y_m) = ((x_1, x_{1'}), \dots, (x_m, x_{m'})) \in Y^m = X^{2m}$. It follows from the definition of the polyhedral product that $\boldsymbol{y} \notin (Y, B)^{\mathcal{K}}$ if and only if $\omega_Y(\boldsymbol{y}) \notin \mathcal{K}$. The latter is equivalent to the condition $\omega_X(\boldsymbol{x}) \notin \mathcal{D}(\mathcal{K})$, since if $\omega_Y(\boldsymbol{y}) = \{i_1, \dots, i_k\}$ then $\omega_X(\boldsymbol{x}) \supset \{i_1, i_1', \dots, i_k, i_k'\}$. Therefore,

$$\boldsymbol{y} \notin (Y, B)^{\mathcal{K}} \iff \boldsymbol{x} \notin (X, A)^{\mathcal{D}(\mathcal{K})},$$

which implies that $(X, A)^{\mathcal{D}(\mathcal{K})} = (Y, B)^{\mathcal{K}}$. □

REMARK. The simplicial wedge [**189**], [**12**] is a generalisation of the doubling construction, in which the ith vertex of $\mathcal{K}$ is replaced by a j_i-tuple of vertices, for some vector $\boldsymbol{j} = (j_1, \dots, j_m)$ of natural numbers. The double corresponds to $\boldsymbol{j} = (2, \dots, 2)$. There is an analogue of Theorem 4.7.14 in this setting, see [**12**, §7].

As an important consequence of Theorem 4.7.14 we obtain the following relationship between the moment-angle complex $\mathcal{Z}_\mathcal{K}$ and its real analogue $\mathcal{R}_\mathcal{K}$:

COROLLARY 4.7.15. *We have $\mathcal{Z}_\mathcal{K} \cong \mathcal{R}_{\mathcal{D}(\mathcal{K})}$.*

PROOF. Apply Theorem 4.7.14 to the pair $(X, A) = (D^1, S^0)$, observing that $(D^1 \times D^1, D^1 \times S^0 \cup S^0 \times D^1) \cong (D^2, S^1)$. □

LEMMA 4.7.16. *Let (X, A) be a pair of cell complexes such that A has a collar neighbourhood $U(A)$ in X (i.e. there is a homeomorphism of pairs $(U(A), A) \cong (A \times [0, 1), A \times \{0\})$). Let $Y = X_1 \cup_A X_2$ be the space obtained by attaching two copies of X along A. Then*

$$\operatorname{rank} H^*(Y) \geqslant \operatorname{rank} H^*(A).$$

Proof. The assumption on (X, A) implies that we can apply the Mayer–Vietoris sequence to the decomposition $Y = X_1 \cup_A X_2$:

$$\cdots \xrightarrow{\beta_{k-1}} H^{k-1}(A) \xrightarrow{\delta_{k-1}} H^k(Y) \xrightarrow{\alpha_k} H^k(X_1) \oplus H^k(X_2) \xrightarrow{\beta_k} H^k(A) \to \cdots .$$

The map β_k is $i_1^* \oplus (-i_2^*)$, where $i_1\colon A \to X_1$ and $i_2\colon A \to X_2$ are the inclusions. Since $X_1 = X_2 = X$, and the inclusions i_1 and i_2 coincide, we have $\operatorname{rank}\operatorname{Ker}\beta_k \geqslant \operatorname{rank} H^k(X)$ and $\operatorname{rank}\operatorname{Im}\beta_k \leqslant \operatorname{rank} H^k(X)$. Using these inequalities we calculate

$$\begin{aligned}\operatorname{rank} H^k(Y) &= \operatorname{rank}\operatorname{Ker}\alpha_k + \operatorname{rank}\operatorname{Im}\alpha_k = \operatorname{rank}\operatorname{Im}\delta_{k-1} + \operatorname{rank}\operatorname{Ker}\beta_k\\ &\geqslant \operatorname{rank} H^{k-1}(A) - \operatorname{rank}\operatorname{Im}\beta_{k-1} + \operatorname{rank} H^k(X)\\ &\geqslant \operatorname{rank} H^{k-1}(A) - \operatorname{rank} H^{k-1}(X) + \operatorname{rank} H^k(X).\end{aligned}$$

The required inequality is obtained by summing up over k. □

Theorem 4.7.17. *Let $\mathcal{K}$ be a simplicial complex of dimension $n-1$ with m vertices, and let $\mathcal{R}_\mathcal{K}$ be the corresponding real moment-angle complex. Then*

$$\operatorname{rank} H^*(\mathcal{R}_\mathcal{K}) \geqslant 2^{m-n'} \geqslant 2^{m-n},$$

where n' is the minimum of the cardinality of maximal simplices of $\mathcal{K}$ (so that $n' = n = \dim\mathcal{K} + 1$ if and only if $\mathcal{K}$ is pure).

Proofs of Theorems 4.7.10 and 4.7.17. We first prove the inequality for $\mathcal{R}_\mathcal{K}$, by induction on the number of vertices m. For $m = 1$ the statement is clear. We embed $\mathcal{R}_\mathcal{K}$ as a subcomplex in the 'big' cube $[-1, 1]^m$ (see Construction 4.1.5) with coordinates $\boldsymbol{u} = (u_1, \ldots, u_m)$, $-1 \leqslant u_i \leqslant 1$. Assume that the first vertex of $\mathcal{K}$ belongs to an $(n'-1)$-dimensional maximal simplex of $\mathcal{K}$, and consider the following subspaces of $\mathcal{R}_\mathcal{K}$:

$$X_+ = \{\boldsymbol{u} \in \mathcal{R}_\mathcal{K}\colon u_1 \geqslant 0\}, \quad X_- = \{\boldsymbol{u} \in \mathcal{R}_\mathcal{K}\colon u_1 \leqslant 0\},$$
$$A = X_+ \cap X_- = \{\boldsymbol{u} \in \mathcal{R}_\mathcal{K}\colon u_1 = 0\}.$$

Applying Lemma 4.7.16 to the decomposition $\mathcal{R}_\mathcal{K} = X_+ \cup_A X_-$ we obtain

$$\operatorname{rank} H^*(\mathcal{R}_\mathcal{K}) \geqslant \operatorname{rank} H^*(A).$$

On the other hand, A is the disjoint union of 2^{m-m_1-1} copies of $\mathcal{R}_{\mathrm{lk}_\mathcal{K}\{1\}}$, where m_1 is the number of vertices of $\mathrm{lk}_\mathcal{K}\{1\}$. Since $\{1\}$ is a vertex of a maximal simplex of $\mathcal{K}$ of minimal cardinality n', the minimal cardinality of maximal simplices in $\mathrm{lk}_\mathcal{K}\{1\}$ is $n_1' = n' - 1$. Now using the inductive hypothesis we obtain

$$\operatorname{rank} H^*(A) = 2^{m-m_1-1} \operatorname{rank} H^*(\mathcal{R}_{\mathrm{lk}_\mathcal{K}\{1\}}) \geqslant 2^{m-m_1-1} 2^{m_1-n_1'} = 2^{m-n'}.$$

Theorem 4.7.17 is therefore proved.

To prove Theorem 4.7.10 we use the fact that $\mathcal{Z}_\mathcal{K} \cong \mathcal{R}_{\mathcal{D}(\mathcal{K})}$, and observe that the numbers $m - n$ (and $m - n'$) for $\mathcal{K}$ and $\mathcal{D}(\mathcal{K})$ coincide. □

Using Theorems 4.4.4 and 4.4.8 we may reformulate Theorem 4.7.10 in both algebraic and combinatorial terms:

Theorem 4.7.18. *Let $\mathcal{K}$ be a simplicial complex of dimension $n-1$ with m vertices, and let $\mathbf{k}$ be a field. Then*

$$\dim \operatorname{Tor}_{\mathbf{k}[v_1,\ldots,v_m]}\bigl(\mathbf{k}[\mathcal{K}], \mathbf{k}\bigr) = \sum_{J\subset[m],\, k\geqslant 0} \dim \widetilde{H}^{k-|J|-1}(\mathcal{K}_J; \mathbf{k}) \geqslant 2^{m-n}.$$

As a corollary we obtain that the weak Horrocks Conjecture (Conjecture A.2.12) holds for a particular class of rings:

COROLLARY 4.7.19. *Let $\mathcal{K}$ be a Cohen–Macaulay simplicial complex (over a field $\mathbf{k}$) of dimension $n-1$ with m vertices. Let $\boldsymbol{t}=(t_1,\ldots,t_n)$ be an lsop in $\mathbf{k}[\mathcal{K}]$, so that $\mathbf{k}[m]/\boldsymbol{t}\cong\mathbf{k}[w_1,\ldots,w_{m-n}]$ and $\dim\mathbf{k}[\mathcal{K}]/\boldsymbol{t}<\infty$. Then*

$$\dim\operatorname{Tor}_{\mathbf{k}[w_1,\ldots,w_{m-n}]}\bigl(\mathbf{k}[\mathcal{K}]/\boldsymbol{t},\mathbf{k}\bigr)\geqslant 2^{m-n},$$

i.e. the weak Horrocks Conjecture holds for the rings $\mathbf{k}[\mathcal{K}]/\boldsymbol{t}$.

PROOF. This follows from the previous theorem and Proposition 4.4.6. □

Exercises.

4.7.20. Show that $\operatorname{ftr}\mathcal{Z}_\mathcal{K}=1$ if and only if $\mathcal{K}=\partial\Delta^{m-1}$.

4.7.21. The Toral Rank Conjecture fails if $\dim X=\infty$.

4.7.22. Show that the doubling operation respects the join, that is, $\mathcal{D}(\mathcal{K}*\mathcal{L})=\mathcal{D}(\mathcal{K})*\mathcal{D}(\mathcal{L})$.

4.7.23. Assume that $\mathcal{K}$ is the boundary complex of a simplicial n-polytope $Q\subset\mathbb{R}^n$ with m vertices $v_1,\ldots,v_m$. Then $\mathcal{D}(\mathcal{K})$ is the boundary of a simplicial polytope $\mathcal{D}(Q)$ of dimension $m+n$ with $2m$ vertices, which can be obtained in the following way. We embed $\mathbb{R}^n$ as the coordinate subspace in $\mathbb{R}^{m+n}$ on the last n coordinates. For each vertex $v_i\in Q\subset\mathbb{R}^n$ take the line $l_i\subset\mathbb{R}^{m+n}$ through v_i parallel to the ith coordinate line of $\mathbb{R}^{m+n}$, for $1\leqslant i\leqslant m$. Then replace each v_i by a pair of points $v_i',v_i''\in l_i$ such that v_i the centre of the segment with the vertices v_i',v_i''. Then the boundary of

$$\mathcal{D}(Q)=\operatorname{conv}(v_1',v_1'',\ldots,v_m',v_m'')\subset\mathbb{R}^{m+n}$$

is $\mathcal{D}(\mathcal{K})$.

4.7.24. There is the following generalisation of Theorem 4.7.14. Let

$$(\boldsymbol{X},\boldsymbol{A})=\bigl\{(X_1,A_1),(X_{1'},A_{1'}),\ldots,(X_m,A_m),(X_{m'},A_{m'})\bigr\}$$

be a set of $2m$ pairs of spaces. Define a new set $(\boldsymbol{Y},\boldsymbol{B})=\{(Y_1,B_1),\ldots,(Y_m,B_m)\}$ of m pairs, where

$$(Y_i,B_i)=\bigl(X_i\times X_{i'},X_i\times A_{i'}\cup A_i\times X_{i'}\bigr).$$

Then

$$(\boldsymbol{X},\boldsymbol{A})^{\mathcal{D}(\mathcal{K})}\cong(\boldsymbol{Y},\boldsymbol{B})^{\mathcal{K}}.$$

Further generalisations can be found in [**12**, §7].

4.7.25. The inequality $\operatorname{rank}H^*(\mathcal{R}_\mathcal{K})\geqslant 2^{m-n}$ of Theorem 4.7.17 (or the inequality $\operatorname{rank}H^*(\mathcal{Z}_\mathcal{K})\geqslant 2^{m-n}$) turns into identity if and only if

$$\mathcal{K}=\partial\Delta^{k_1-1}*\partial\Delta^{k_2-1}*\cdots*\partial\Delta^{k_p-1}*\Delta^{m-s-1},$$

where $s=k_1+\cdots+k_p$ and the join factor Δ^{m-s-1} is void if $s=m$ (compare Exercise 3.3.19). In this case both $\mathcal{R}_\mathcal{K}$ and $\mathcal{Z}_\mathcal{K}$ are products of spheres and a disc.

4.8. Massey products in the cohomology of moment-angle complexes

Here we address the question of existence of nontrivial triple Massey products in the Koszul complex

$$[\Lambda[u_1, \dots, u_m] \otimes \mathbb{Z}[\mathcal{K}], d]$$

of the face ring, and therefore in the cohomology of $\mathcal{Z}_\mathcal{K}$. The general definition of Massey products in the cohomology of a differential graded algebra is reviewed in Section A.4 of the Appendix. A geometrical approach to constructing nontrivial triple Massey products in the Koszul complex of the face ring was developed by Baskakov in **[19]** as an extension of the cohomology calculation in Theorem 4.4.7. It is well-known that non-trivial higher Massey products obstruct the *formality* of a differential graded algebra, which in our case leads to a family of nonformal moment-angle manifolds $\mathcal{Z}_\mathcal{K}$ (see Section B.2 for the background material).

Construction 4.8.1 (Baskakov **[19]**). Let $\mathcal{K}_i$ be a triangulation of a sphere S^{n_i-1} with $|V_i| = m_i$ vertices, $i = 1, 2, 3$. Set $m = m_1 + m_2 + m_3$, $n = n_1 + n_2 + n_3$,

$$\mathcal{K} = \mathcal{K}_1 * \mathcal{K}_2 * \mathcal{K}_3, \quad \mathcal{Z}_\mathcal{K} = \mathcal{Z}_{\mathcal{K}_1} \times \mathcal{Z}_{\mathcal{K}_2} \times \mathcal{Z}_{\mathcal{K}_3}.$$

Then $\mathcal{K}$ is a triangulation of S^{n-1} and therefore $\mathcal{Z}_\mathcal{K}$ is an $(m+n)$-manifold.

We choose maximal simplices $I_1 \in \mathcal{K}_1$, $I_2', I_2'' \in \mathcal{K}_2$ such that $I_2' \cap I_2'' = \varnothing$, and $I_3 \in \mathcal{K}_3$. Set

$$\widetilde{\mathcal{K}} = \mathrm{ss}_{I_1 \cup I_2'}(\mathrm{ss}_{I_2'' \cup I_3} \mathcal{K}),$$

where ss_I denotes the stellar subdivision at I, see Definition 2.7.1. Then $\widetilde{\mathcal{K}}$ is a triangulation of S^{n-1} with $m + 2$ vertices. Take generators

$$\beta_i \in \widetilde{H}^{n_i-1}(\widetilde{\mathcal{K}}_{V_i}) \cong \widetilde{H}^{n_i-1}(S^{n_i-1}), \quad \text{for } i = 1, 2, 3,$$

where $\widetilde{\mathcal{K}}_{V_i}$ is the restriction of $\widetilde{\mathcal{K}}$ to the vertex set of $\mathcal{K}_i$, and set

$$\alpha_i = h(\beta_i) \in H^{n_i-m_i, 2m_i}(\mathcal{Z}_{\widetilde{\mathcal{K}}}) \subset H^{m_i+n_i}(\mathcal{Z}_{\widetilde{\mathcal{K}}}),$$

where h is the isomorphism of Theorem 4.4.7. Then

$$\beta_1\beta_2 \in \widetilde{H}^{n_1+n_2-1}(\widetilde{\mathcal{K}}_{V_1 \sqcup V_2}) \cong \widetilde{H}^{n_1+n_2-1}(S^{n_1+n_2-1} \setminus pt) = 0,$$

hence $\alpha_1\alpha_2 = h(\beta_1\beta_2) = 0$, and similarly $\alpha_2\alpha_3 = 0$. Therefore, the triple Massey product $\langle \alpha_1, \alpha_2, \alpha_3 \rangle \subset H^{m+n-1}(\mathcal{Z}_{\widetilde{\mathcal{K}}})$ is defined. By definition, it is the set of cohomology classes represented by the cocycles $(-1)^{\deg a_1+1} a_1 f + e a_3$ where a_i is a cocycle representing α_i, and e, f are cochains satisfying $de = a_1 a_2$, $df = a_2 a_3$.

For the simplest example of this series, take $\mathcal{K}_i = S^0$ (two points), so that $\mathcal{K}$ is the boundary of an octahedron, and $\widetilde{\mathcal{K}}$ is obtained by applying stellar subdivisions at two skew edges. We shall consider this example in more detail below.

Recall that a Massey product is *trivial* if it contains zero.

Theorem 4.8.2. *The above defined triple Massey product*

$$\langle \alpha_1, \alpha_2, \alpha_3 \rangle \subset H^{m+n-1}(\mathcal{Z}_{\widetilde{K}})$$

is nontrivial.

Proof. Consider the subcomplex of $\widetilde{\mathcal{K}}$ consisting of the two new vertices added to $\mathcal{K}$ in the process of stellar subdivision. By Proposition 4.1.11 (b), the inclusion of this subcomplex induces an embedding of a 3-dimensional sphere $S^3 \hookrightarrow \mathcal{Z}_{\widetilde{K}}$. Since the two new vertices are not joined by an edge in $\mathcal{Z}_{\widetilde{\mathcal{K}}}$, the embedded 3-sphere defines a non-trivial class $x \in H_3(\mathcal{Z}_{\widetilde{\mathcal{K}}})$. Its Poincaré dual cohomology class in $H^{m+n-1}(\mathcal{Z}_{\widetilde{\mathcal{K}}})$ is contained in the Massey product $\langle\alpha_1, \alpha_2, \alpha_3\rangle$. We need only to check that this element cannot be turned into zero by adding elements from the indeterminacy of the Massey product, i.e. from the subspace

$$\alpha_1 \cdot H^{m_2+m_3+n_2+n_3-1}(\mathcal{Z}_{\widetilde{\mathcal{K}}}) + \alpha_3 \cdot H^{m_1+m_2+n_1+n_2-1}(\mathcal{Z}_{\widetilde{\mathcal{K}}}).$$

To do this we use the multigraded structure in $H^*(\mathcal{Z}_{\widetilde{\mathcal{K}}})$. The multigraded components of the group $H^{m_2+m_3+n_2+n_3-1}(\mathcal{Z}_{\widetilde{\mathcal{K}}})$ different from the component determined by the full subcomplex $\widetilde{\mathcal{K}}_{V_2 \sqcup V_3}$ do not affect the nontriviality of the Massey product, while the multigraded component corresponding to $\widetilde{\mathcal{K}}_{V_2 \sqcup V_3}$ is zero since $\widetilde{\mathcal{K}}_{V_2 \sqcup V_3} \cong S^{n_2+n_3-1} \setminus pt$ is contractible. The group $H^{m_1+m_2+n_1+n_2-1}(\mathcal{Z}_{\widetilde{\mathcal{K}}})$ is considered similarly. It follows that the Massey product contains a unique nonzero element in its multigraded component and therefore it is nontrivial. □

Corollary 4.8.3. *Let $\widetilde{\mathcal{K}}$ be a triangulated sphere obtained from another sphere $\mathcal{K}$ by applying two stellar subdivisions as described above. Then the corresponding 2-connected moment-angle manifold $\mathcal{Z}_{\widetilde{\mathcal{K}}}$ is nonformal.*

In the proof of Theorem 4.8.2 the nontriviality of the Massey product is established geometrically. A parallel argument may be carried out algebraically using the Koszul complex or its quotient algebra $R^*(\mathcal{K})$, as illustrated in the next example. To be precise, the nonformality of a manifold $\mathcal{Z}_{\mathcal{K}}$ is equivalent to the nonformality of its *singular* cochain algebra $C^*(\mathcal{Z}_{\mathcal{K}}; \mathbb{Q})$ (or Sullivan's algebra $A_{PL}(\mathcal{Z}_{\mathcal{K}})$), while the rational Koszul complex or the algebra $R^*(\mathcal{K}) \otimes \mathbb{Q}$ are quasi-isomorphic to the *cellular* cochain algebra $\mathcal{C}^*(\mathcal{Z}_{\mathcal{K}}; \mathbb{Q})$. However, this difference is irrelevant: it can be easily seen that the existence of a nontrivial triple Massey product for the cellular cochain algebra $\mathcal{C}^*(\mathcal{Z}_{\mathcal{K}}; \mathbb{Q})$ implies that $\mathcal{Z}_{\mathcal{K}}$ is nonformal (an exercise, or see [**75**, Proposition 5.6.1]).

Example 4.8.4. We consider the simplest case of Construction 4.8.1, when $\widetilde{\mathcal{K}}$ is obtained from the boundary of an octahedron by applying stellar subdivisions at two skew edges. Then $\widetilde{\mathcal{K}} = \mathcal{K}_P$ is the nerve complex of a simple 3-polytope P obtained by truncating a cube at two edges as shown in Figure 4.3. The face ring is given by

$$\mathbb{Z}[\mathcal{K}_P] = \mathbb{Z}[v_1, \ldots, v_6, w_1, w_2]/\mathcal{I}_{\mathcal{K}_P},$$

where v_i, $i = 1, \ldots, 6$, are the generators coming from the facets of the cube and w_1, w_2 are the generators corresponding to the two new facets, and

$$\mathcal{I}_{\mathcal{K}_P} = (v_1v_2, v_3v_4, v_5v_6, w_1w_2, v_1v_3, v_4v_5, w_1v_3, w_1v_6, w_2v_2, w_2v_4).$$

We denote the corresponding exterior generators of $R^*(\mathcal{K}_P)$ by $u_1, \ldots, u_6, t_1, t_2$; they satisfy $du_i = v_i$ and $dt_i = w_i$. Consider the cocycles

$$a = v_1u_2, \quad b = v_3u_4, \quad c = v_5u_6$$

and the corresponding cohomology classes $\alpha, \beta, \gamma \in H^{-1,4}[R^*(\mathcal{K})]$. The equations

$$ab = de, \quad bc = df$$

have a solution $e = 0$, $f = v_5u_3u_4u_6$, so the triple Massey product $\langle\alpha, \beta, \gamma\rangle \in H^{-4,12}[R^*(\mathcal{K})]$ is defined. This Massey product is represented by the cocycle

$$af + ec = v_1v_5u_2u_3u_4u_6$$



FIGURE 4.3.

and is nontrivial. The differential graded algebra $R^*(\mathcal{K}_P)$ and the 11-dimensional manifold $\mathcal{Z}_{\mathcal{K}_P}$ are not formal.

REMARK. We may truncate another edge of the polytope P from the previous example to obtain a 3-dimensional associahedron As^3, shown in Figure 1.4 (left). By considering similar nontrivial Massey products (now there will be three of them, corresponding to each pair of truncated edges) we deduce that the 12-dimensional moment-angle manifold corresponding to As^3 is also nonformal.

In view of Theorem 4.8.2, the question arises of describing the class of simplicial complexes $\mathcal{K}$ for which the algebra $R^*(\mathcal{K})$ (equivalently, the Koszul algebra $[\Lambda[u_1,\ldots,u_m]\otimes\mathbb{Z}[\mathcal{K}],d]$) is formal. For example, this is the case if $\mathcal{K}$ is the boundary of a polygon or, more generally, if $\mathcal{K}$ is of the form described in Theorem 4.5.12.

Triple Massey products in the cohomology of $\mathcal{Z}_{\mathcal{K}}$ were further studied in the work of Denham and Suciu [**75**]. According to [**75**, Theorem 6.1.1], there exists a nontrivial triple Massey product of 3-dimensional cohomology classes $\alpha,\beta,\gamma\in H^3(\mathcal{Z}_{\mathcal{K}})$ if and only if the 1-skeleton of $\mathcal{K}$ contains an induced subgraph isomorphic to one of the five explicitly described 'obstruction' graphs. In [**75**, Example 8.5.1] there is also constructed an example of $\mathcal{K}$ for which the corresponding $\mathcal{Z}_{\mathcal{K}}$ has an *indecomposable* triple Massey product in the cohomology (a triple Massey product is indecomposable if it does not contain a cohomology class that can be written as a product of two cohomology classes of positive dimension).

To conclude this section, we mention that the algebraic study of Massey products in the cohomology of Koszul complexes has a long history. It goes back to the work of Golod [**100**], who studied the Poincaré series of $\mathrm{Tor}_R(\mathbf{k},\mathbf{k})$ for a Nötherian local ring R. The main result of Golod is a calculation of the Poincaré series for the class of rings whose Koszul complexes have all Massey products vanishing (including the cohomology multiplication). Such rings were called *Golod* in the monograph [**113**] of Gulliksen and Levin, where the reader can find a detailed exposition of Golod's theorem together with several further applications.

DEFINITION 4.8.5. We refer to a simplicial complex $\mathcal{K}$ as *Golod* if its face ring $\mathbf{k}[\mathcal{K}]$ has the Golod property, i.e. if the multiplication and all higher Massey products in the Koszul complex $[\Lambda[u_1,\ldots,u_m]\otimes\mathbf{k}[\mathcal{K}],d]$ are trivial.

Golod complexes were studied in [**120**], where several combinatorial criteria for Golodness were given. The appearance of moment-angle complexes added a topological dimension to the whole study. A wide class of Golod complexes $\mathcal{K}$ was described in [**104**, Theorem 11.2] using the theory of moment-angle complexes. This class includes skeleta of simplices considered in Theorem 4.6.7, and, more generally, all *shifted* complexes. For all such $\mathcal{K}$ the corresponding moment-angle complex $\mathcal{Z}_\mathcal{K}$ is homotopy equivalent to a wedge of spheres.

Exercises.

4.8.6. If the cellular cochain algebra $\mathcal{C}^*(\mathcal{Z}_\mathcal{K};\mathbb{Q})$ carries a nontrivial triple Massey product, then $\mathcal{Z}_\mathcal{K}$ is nonformal.

4.9. Moment-angle complexes from simplicial posets

Simplicial posets $\mathcal{S}$ generalise naturally abstract simplicial complexes (see Section 2.8). Algebraic properties of their face rings $\mathbf{k}[\mathcal{S}]$ were discussed in Section 3.5. Following the categorical description of the moment-angle complex $\mathcal{Z}_\mathcal{K}$ outlined in the end of Construction 4.1.1, it is easy to extend the definition of $\mathcal{Z}_\mathcal{K}$ to simplicial posets. The resulting space $\mathcal{Z}_\mathcal{S}$ carries a torus action, and its equivariant and ordinary cohomology is expressed in terms of the face ring $\mathbb{Z}[\mathcal{S}]$ in the same way as for the standard moment-angle complexes $\mathcal{Z}_\mathcal{K}$. Simplicial posets and associated moment-angle complexes therefore provide a broader context for studying the link between torus actions and combinatorial commutative algebra. These developments are originally due to Lü and Panov [**142**].

Let $\mathcal{S}$ be a finite simplicial poset with the vertex set $V(\mathcal{S})=[m]$.

Construction 4.9.1 (moment-angle complex). We consider the face category $\textsc{cat}(\mathcal{S})$ whose objects are elements $\sigma\in\mathcal{S}$ and there is a morphism from σ to τ whenever $\sigma\leqslant\tau$. For each element $\sigma\in\mathcal{S}$ we define the following subset in the standard unit polydisc $\mathbb{D}^m\subset\mathbb{C}^m$:

$$(D^2,S^1)^\sigma=\bigl\{(z_1,\ldots,z_m)\in\mathbb{D}^m\colon |z_j|^2=1 \text{ for } j\not\leqslant\sigma\bigr\}.$$

Then $(D^2,S^1)^\sigma$ is homeomorphic to a product of $|\sigma|$ discs and $m-|\sigma|$ circles. We have an inclusion $(D^2,S^1)^\tau\subset(D^2,S^1)^\sigma$ whenever $\tau\leqslant\sigma$. Now define a diagram

$$\begin{aligned}\mathcal{D}_\mathcal{S}(D^2,S^1)\colon \textsc{cat}(\mathcal{S})&\longrightarrow \textsc{top},\\ \sigma&\longmapsto(D^2,S^1)^\sigma,\end{aligned}$$

which maps a morphism $\sigma\leqslant\tau$ of $\textsc{cat}(\mathcal{S})$ to the inclusion $(D^2,S^1)^\sigma\subset(D^2,S^1)^\tau$.

We define the *moment-angle complex* corresponding to $\mathcal{S}$ by

$$\mathcal{Z}_\mathcal{S}=\operatorname{colim}\mathcal{D}_\mathcal{S}(D^2,S^1)=\operatorname*{colim}_{\sigma\in\mathcal{S}}(D^2,S^1)^\sigma.$$

The space $\mathcal{Z}_\mathcal{S}$ is therefore glued from the blocks $(D^2,S^1)^\sigma$ according to the poset relation in $\mathcal{S}$. When $\mathcal{S}$ is (the face poset of) a simplicial complex $\mathcal{K}$ it becomes the standard moment-angle complex $\mathcal{Z}_\mathcal{K}$.

Since every subset $(D^2,S^1)^\sigma\subset\mathbb{D}^m$ is invariant with respect to the coordinatewise action of the m-torus $\mathbb{T}^m$, the moment-angle complex $\mathcal{Z}_\mathcal{S}$ acquires a $\mathbb{T}^m$-action.

This definition extends to a set $(\boldsymbol{X},\boldsymbol{A})$ of m pairs of spaces (see Construction 4.1.9), so we define the *polyhedral product* of $(\boldsymbol{X},\boldsymbol{A})$ corresponding to $\mathcal{S}$ by

$$(\boldsymbol{X},\boldsymbol{A})^\mathcal{S}=\operatorname{colim}\mathcal{D}_\mathcal{S}(\boldsymbol{X},\boldsymbol{A})=\operatorname*{colim}_{\sigma\in\mathcal{S}}(\boldsymbol{X},\boldsymbol{A})^\sigma.$$

The construction of the polyhedral product $(\boldsymbol{X},\boldsymbol{A})^{\mathcal{S}}$ is functorial in all arguments: there are straightforward analogues of Propositions 4.1.11 and 4.1.12, which are proved in a similar way.

EXAMPLE 4.9.2. Let $\mathcal{S}$ be the simplicial poset of Fig. 3.3 (a). Then $\mathcal{Z}_{\mathcal{S}}$ is obtained by gluing two copies of $D^2\times D^2$ along their boundary $S^3=D^2\times S^1\cup S^1\times D^2$. Therefore, $\mathcal{Z}_{\mathcal{S}}\cong S^4$. Here, $\mathcal{K}_{\mathcal{S}}=\Delta^1$ (a segment), and the moment-angle complex map induced by the map $\mathcal{S}\to\mathcal{K}_{\mathcal{S}}$ (2.9) folds S^4 onto D^4. Similarly, if $\mathcal{S}$ is of Fig. 3.3 (b), then $\mathcal{Z}_{\mathcal{S}}\cong S^6$. Note that even-dimensional spheres do not appear as moment-angle complexes $\mathcal{Z}_{\mathcal{K}}$ for simplicial complexes $\mathcal{K}$.

The *join* of simplicial posets $\mathcal{S}_1$ and $\mathcal{S}_2$ is the simplicial poset $\mathcal{S}_1*\mathcal{S}_2$ whose elements are pairs (σ_1,σ_2), with $(\sigma_1,\sigma_2)\leqslant(\tau_1,\tau_2)$ whenever $\sigma_1\leqslant\tau_1$ in $\mathcal{S}_1$ and $\sigma_2\leqslant\tau_2$ in $\mathcal{S}_2$. The following properties of $\mathcal{Z}_{\mathcal{S}}$ are similar to those of $\mathcal{Z}_{\mathcal{K}}$.

THEOREM 4.9.3.

(a) $\mathcal{Z}_{\mathcal{S}_1*\mathcal{S}_2}\cong\mathcal{Z}_{\mathcal{S}_1}\times\mathcal{Z}_{\mathcal{S}_2}$;
(b) *the quotient* $\mathcal{Z}_{\mathcal{S}}/\mathbb{T}^m$ *is homeomorphic to the cone over* $|\mathcal{S}|$;
(c) *if* $|\mathcal{S}|\cong S^{n-1}$, *then* $\mathcal{Z}_{\mathcal{S}}$ *is a manifold of dimension* $m+n$.

PROOF. Statements (a) and (b) are proved in the same way as the corresponding statements for simplicial complexes, see Section 4.1. To prove (c) we use the 'dual' decomposition of the boundary of the n-ball cone $|\mathcal{S}|$ into faces, in the same way as in the proof of Theorem 4.1.4. □

CONSTRUCTION 4.9.4 (cell decomposition). We proceed by analogy with the construction of Section 4.3. The disc $\mathbb{D}$ is decomposed into 3 cells: the point $1\in\mathbb{D}$ is the 0-cell; the complement to 1 in the boundary circle is the 1-cell, which we denote T; and the interior of $\mathbb{D}$ is the 2-cell, which we denote D. The polydisc $\mathbb{D}^m$ then acquires the product cell decomposition, with each $(D^2,S^1)^\sigma\subset(D^2,S^1)^\tau$ being an inclusion of cellular subcomplexes for $\sigma\leqslant\tau$. We therefore obtain a cell decomposition of $\mathcal{Z}_{\mathcal{S}}$ (although this time it is not a subcomplex in $\mathbb{D}^m$ in general!). Each cell in $\mathcal{Z}_{\mathcal{S}}$ is determined by an element $\sigma\in\mathcal{S}$ and a subset $\omega\in V(\mathcal{S})$ with $V(\sigma)\cap\omega=\varnothing$. Such a cell is a product of $|\sigma|$ cells of D-type, $|\omega|$ cells of T-type and the rest of 1-type. We denote this cell by $\kappa(\omega,\sigma)$.

The resulting cellular cochain complex $\mathcal{C}^*(\mathcal{Z}_{\mathcal{S}})$ has an additive basis consisting of cochains $\kappa(\omega,\sigma)^*$ dual to the corresponding cells. We introduce a $\mathbb{Z}\oplus\mathbb{Z}^m$-grading on the cochains by setting

$$\operatorname{mdeg}\kappa(\omega,\sigma)^*=(-|\omega|,2V(\sigma)+2\omega),$$

where we think of both $V(\sigma)$ and ω as vectors in $\{0,1\}^m\subset\mathbb{Z}^m$. The cellular differential preserves the $\mathbb{Z}^m$-part of the multigrading, so we obtain a decomposition

$$\mathcal{C}^*(\mathcal{Z}_{\mathcal{S}})=\bigoplus_{\boldsymbol{a}\in\mathbb{Z}^m}\mathcal{C}^{*,2\boldsymbol{a}}(\mathcal{Z}_{\mathcal{K}})$$

into a sum of subcomplexes. The only nontrivial subcomplexes are those for which $\boldsymbol{a}$ is in $\{0,1\}^m$. The cellular cohomology of $\mathcal{Z}_{\mathcal{S}}$ thereby acquires a multigrading, and we define the *multigraded Betti numbers* $b^{-i,2\boldsymbol{a}}(\mathcal{Z}_{\mathcal{S}})$ by

$$b^{-i,2\boldsymbol{a}}(\mathcal{Z}_{\mathcal{S}})=\operatorname{rank}H^{-i,2\boldsymbol{a}}(\mathcal{Z}_{\mathcal{S}}),\quad\text{for }1\leqslant i\leqslant m,\ \boldsymbol{a}\in\mathbb{Z}^m.$$

For the ordinary Betti numbers we have $b^k(\mathcal{Z}_{\mathcal{S}})=\sum_{-i+2|\boldsymbol{a}|=k}b^{-i,2\boldsymbol{a}}(\mathcal{Z}_{\mathcal{S}})$.

The map of moment-angle complexes $\mathcal{Z}_{\mathcal{S}_1} \to \mathcal{Z}_{\mathcal{S}_2}$ induced by a simplicial poset map $\mathcal{S}_1 \to \mathcal{S}_2$ is clearly a cellular map, and therefore the cellular cohomology is functorial with respect to such maps.

We now recall from Section 3.5 that the face ring $\mathbb{Z}[\mathcal{S}]$ is a $\mathbb{Z}^m$-graded $\mathbb{Z}[v_1, \ldots, v_m]$-module via the map sending each v_i identically, and we have the $\mathbb{Z} \oplus \mathbb{Z}^m$-graded Tor-algebra of $\mathbb{Z}[\mathcal{S}]$:

$$\operatorname{Tor}_{\mathbb{Z}[v_1,\ldots,v_m]}\bigl(\mathbb{Z}[\mathcal{S}], \mathbb{Z}\bigr) = \bigoplus_{i \geqslant 0,\, \boldsymbol{a} \in \mathbb{N}^m} \operatorname{Tor}^{-i,\,2\boldsymbol{a}}_{\mathbb{Z}[v_1,\ldots,v_m]}\bigl(\mathbb{Z}[\mathcal{S}], \mathbb{Z}\bigr).$$

Theorem 4.9.5. *There is a graded ring isomorphism*

$$H^*(\mathcal{Z}_{\mathcal{S}}) \cong \operatorname{Tor}_{\mathbb{Z}[v_1,\ldots,v_m]}(\mathbb{Z}[\mathcal{S}], \mathbb{Z})$$

whose graded components are given by the group isomorphisms

$$H^p(\mathcal{Z}_{\mathcal{S}}) \cong \bigoplus_{-i+2|\boldsymbol{a}|=p} \operatorname{Tor}^{-i,\,2\boldsymbol{a}}_{\mathbb{Z}[v_1,\ldots,v_m]}(\mathbb{Z}[\mathcal{S}], \mathbb{Z}) \tag{4.24}$$

in each degree p. Here $|\boldsymbol{a}| = j_1 + \cdots + j_m$ for $\boldsymbol{a} = (j_1, \ldots, j_m)$.

Using the Koszul complex we restate the above theorem as follows:

Theorem 4.9.6. *There is a graded ring isomorphism*

$$H^*(\mathcal{Z}_{\mathcal{S}}) \cong H\bigl[\Lambda[u_1, \ldots, u_m] \otimes \mathbb{Z}[\mathcal{S}], d\bigr],$$

where the $\mathbb{Z} \oplus \mathbb{Z}^m$-grading and the differential on the right hand side are defined by

$$\operatorname{mdeg} u_i = (-1, 2\boldsymbol{e}_i), \quad \operatorname{mdeg} v_\sigma = (0, 2V(\sigma)), \qquad du_i = v_i, \quad dv_\sigma = 0,$$

and $\boldsymbol{e}_i \in \mathbb{Z}^m$ is the ith basis vector, for $i = 1, \ldots, m$.

Proof. The proof follows the lines of the proof of Theorem 4.4.4, but the analogues of Lemmata 3.2.6 and 4.4.3 are proved in a different way.

We first set up the quotient differential graded ring

$$R^*(\mathcal{S}) = \Lambda[u_1, \ldots, u_m] \otimes \mathbb{Z}[\mathcal{S}]/\mathcal{I}_R$$

where $\mathcal{I}_R$ is the ideal generated by the elements

$$u_i v_\sigma \quad \text{with } i \in V(\sigma), \quad \text{and} \quad v_\sigma v_\tau \quad \text{with } \sigma \wedge \tau \neq \hat{0}.$$

Note that the latter condition is equivalent to $V(\sigma) \cap V(\tau) \neq \varnothing$. The ring $R^*(\mathcal{S})$ will serve as an algebraic model for the cellular cochains of $\mathcal{Z}_{\mathcal{S}}$.

We need to prove an analogue of Lemma 3.2.6, i.e. show that the quotient map

$$\varrho\colon \Lambda[u_1, \ldots, u_m] \otimes \mathbb{Z}[\mathcal{S}] \to R^*(\mathcal{S})$$

induces an isomorphism in cohomology. Instead of constructing a chain homotopy directly, we shall identify both $R^*(\mathcal{S})$ and $\Lambda[u_1, \ldots, u_m] \otimes \mathbb{Z}[\mathcal{S}]$ with the cellular cochains of homotopy equivalent spaces.

Theorem 3.5.7 implies that $R^*(\mathcal{S})$ has basis of monomials $u_\omega v_\sigma$ where $\omega \subset V(\mathcal{S})$, $\sigma \in \mathcal{S}$, $\omega \cap V(\sigma) = \varnothing$, and $u_\omega = u_{i_1} \ldots u_{i_k}$ for $\omega = \{i_1, \ldots, i_k\}$. In particular, $R^*(\mathcal{S})$ is a free abelian group of finite rank. The map

$$g\colon R^*(\mathcal{S}) \to \mathcal{C}^*(\mathcal{Z}_{\mathcal{S}}), \quad u_\omega v_\sigma \mapsto \kappa(\omega, \sigma)^* \tag{4.25}$$

is an isomorphism of cochain complexes. Indeed, the additive bases of the two groups are in one-to-one correspondence, and the differential in $R^*(\mathcal{S})$ acts (in the case $|\omega| = 1$ and $i \notin V(\sigma)$) as

$$d(u_i v_\sigma) = v_i v_\sigma = \sum_{\eta \in i \vee \sigma} v_\eta.$$

This is exactly how the cellular differential in $\mathcal{C}^*(\mathcal{Z}_\mathcal{S})$ acts on $\kappa(i,\sigma)^*$. The case of arbitrary ω is treated similarly. It follows that we have an isomorphism of cohomology groups $H^j[R^*(\mathcal{S})] \cong H^j(\mathcal{Z}_\mathcal{S})$ for all j.

The differential ring $[\Lambda[u_1,\dots,u_m] \otimes \mathbb{Z}[\mathcal{S}], d]$ also may be identified with the cellular cochains of a certain space. Namely, consider the polyhedral product $(S^\infty, S^1)^\mathcal{S}$. Then we show in the same way as in Subsection 4.4 that there is an isomorphism of cochain complexes

$$g' \colon \Lambda[u_1,\dots,u_m] \otimes \mathbb{Z}[\mathcal{S}] \to \mathcal{C}^*\bigl((S^\infty, S^1)^\mathcal{S}\bigr).$$

Furthermore, the standard functoriality arguments give a deformation retraction

$$\mathcal{Z}_\mathcal{S} = (D^2, S^1)^\mathcal{S} \hookrightarrow (S^\infty, S^1)^\mathcal{S} \longrightarrow (D^2, S^1)^\mathcal{S}$$

onto a cellular subcomplex. Hence the cochain map $\mathcal{C}^*((S^\infty, S^1)^\mathcal{S}) \to \mathcal{C}^*(\mathcal{Z}_\mathcal{S})$ corresponding to the inclusion $\mathcal{Z}_\mathcal{S} \hookrightarrow (S^\infty, S^1)^\mathcal{S}$ induces an isomorphism in cohomology.

Summarising the above observations we obtain the commutative square

$$\begin{tikzcd}
\Lambda[u_1,\dots,u_m] \otimes \mathbb{Z}[\mathcal{S}] \arrow[r, "g'"] \arrow[d, "\varrho"'] & \mathcal{C}^*((S^\infty, S^1)^\mathcal{S}) \arrow[d] \\
R^*(\mathcal{S}) \arrow[r, "g"] & \mathcal{C}^*(\mathcal{Z}_\mathcal{S})
\end{tikzcd} \tag{4.26}$$

in which the horizontal arrows are isomorphisms of cochain complexes, and the right vertical arrow induces an isomorphism in cohomology. It follows that the left arrow also induces an isomorphism in cohomology, as claimed.

The additive isomorphism of (4.24) now follows from (4.26). To establish the ring isomorphism we need to analyse the multiplication of cellular cochains.

We consider the diagonal approximation map $\widetilde{\Delta} \colon \mathbb{D}^m \to \mathbb{D}^m \times \mathbb{D}^m$ given on each coordinate by (4.12). It restricts to a map $(D^2, S^1)^\sigma \to (D^2, S^1)^\sigma \times (D^2, S^1)^\sigma$ for every $\sigma \in \mathcal{S}$ and gives rise to a map of diagrams

$$\mathcal{D}_\mathcal{S}(D^2, S^1) \to \mathcal{D}_\mathcal{S}(D^2, S^1) \times \mathcal{D}_\mathcal{S}(D^2, S^1).$$

By definition, the colimit of the latter is $\mathcal{Z}_{\mathcal{S} * \mathcal{S}}$, which is identified with $\mathcal{Z}_\mathcal{S} \times \mathcal{Z}_\mathcal{S}$. We therefore obtain a cellular approximation $\widetilde{\Delta} \colon \mathcal{Z}_\mathcal{S} \to \mathcal{Z}_\mathcal{S} \times \mathcal{Z}_\mathcal{S}$ for the diagonal map of $\mathcal{Z}_\mathcal{S}$. It induces a ring structure on the cellular cochains via the composition

$$\mathcal{C}^*(\mathcal{Z}_\mathcal{S}) \otimes \mathcal{C}^*(\mathcal{Z}_\mathcal{S}) \xrightarrow{\;\times\;} \mathcal{C}^*(\mathcal{Z}_\mathcal{S} \times \mathcal{Z}_\mathcal{S}) \xrightarrow{\;\widetilde{\Delta}^*\;} \mathcal{C}^*(\mathcal{Z}_\mathcal{S}).$$

We claim that, with this multiplication in $\mathcal{C}^*(\mathcal{Z}_\mathcal{S})$, the map (4.25) becomes a ring isomorphism. To see this we first observe that since (4.25) is a linear map, it is enough to consider the product of two generators $u_\omega v_\sigma$ and $u_\psi v_\tau$. If any two of the subsets ω, $V(\sigma)$, ψ and $V(\tau)$ have nonempty intersection, then $u_\omega v_\sigma \cdot u_\psi v_\tau = 0$ in $R^*(\mathcal{S})$. Otherwise (if all four subsets are disjoint) we have

$$g(u_\omega v_\sigma \cdot u_\psi v_\tau) = g\bigl(u_{\omega \sqcup \psi} \cdot \sum_{\eta \in \sigma \vee \tau} v_\eta\bigr) = \sum_{\eta \in \sigma \vee \tau} \kappa(\omega \sqcup \psi, \eta)^*. \tag{4.27}$$



FIGURE 4.4. 'Ball with corners' Q dual to the simplicial poset $\mathcal{S}$.

We also observe that for any cell $\kappa(\chi,\eta)$ of $\mathcal{Z}_{\mathcal{S}}$ (with $\chi\cap V(\eta)=\varnothing$) we have

$$\widetilde{\Delta}\kappa(\chi,\eta)=\sum_{\substack{\omega\sqcup\psi=\chi\\ \sigma\vee\tau\,\ni\,\eta}}\kappa(\omega,\sigma)\times\kappa(\psi,\tau).$$

Therefore,

$$\begin{aligned}g(u_\omega v_\sigma)\cdot g(u_\psi v_\tau)&=\kappa(\omega,\sigma)^*\cdot\kappa(\psi,\tau)^*\\&=\widetilde{\Delta}^*\bigl(\kappa(\omega,\sigma)\times\kappa(\psi,\tau)\bigr)^*=\sum_{\eta\,\in\,\sigma\vee\tau}\kappa(\omega\sqcup\psi,\eta)^*.\end{aligned}$$

Comparing this with (4.27) we deduce that (4.25) is a ring map. □

REMARK. Using the monoid structure on $\mathbb{D}$ as in Proposition 4.1.12 one easily sees that the construction of $\mathcal{Z}_{\mathcal{S}}$ is functorial with respect to maps of simplicial posets. This together with Proposition 3.5.12 makes the isomorphism of Theorem 4.9.5 functorial.

Using Hochster's formula for simplicial posets (Theorem 3.5.14) we can calculate the cohomology of $\mathcal{Z}_{\mathcal{S}}$ via the cohomology of full subposets $\mathcal{S}_J\subset\mathcal{S}$. Here is an example of calculation using this method.

EXAMPLE 4.9.7. Let $\mathcal{S}$ be the simplicial poset shown on Fig. 4.4 (right). It has $m=5$ vertices, 9 edges and 6 triangular faces. The dual 3-dimensional 'ball with corners' Q (see the proof of Theorem 4.9.3) is shown in Figure 4.4 (left). We denote its facets $F_1,\dots,F_5$, edges e,f,g and the vertex $\sigma=F_3\cap f$ as shown. The corresponding moment-angle complex $\mathcal{Z}_{\mathcal{S}}$ is an 8-dimensional manifold.

The face ring $\mathbb{Z}[\mathcal{S}]$ is the quotient of the polynomial ring

$$\mathbb{Z}[\mathcal{S}]=\mathbb{Z}[v_1,\dots,v_5,v_e,v_f,v_g],\quad \deg v_i=2,\quad \deg v_e=\deg v_f=\deg v_e=4$$

by the relations

$$\begin{aligned}&v_1v_2=v_e+v_f+v_g,\\&v_3v_4=v_3v_5=v_4v_5=v_3v_e=v_4v_f=v_5v_g=v_ev_f=v_ev_g=v_ev_f=0.\end{aligned}$$

The other generators and relations in the original presentation of $\mathbb{Z}[\mathcal{S}]$ can be derived from the above; e.g., $v_\sigma=v_3v_f$.

Given $J \subset [m]$ we define the following subset in the boundary of Q:

$$Q_J = \bigcup_{j \in J} F_j \subset Q.$$

By analogy with Proposition 3.2.11 we prove that

$$H^{-i,\,2J}(\mathcal{Z}_{\mathcal{S}}) \cong \widetilde{H}^{|J|-i-1}(Q_J). \tag{4.28}$$

Using this formula we calculate the nontrivial cohomology groups of $\mathcal{Z}_{\mathcal{S}}$ as follows:

$$\begin{array}{rll}
H^{0,(0,0,0,0,0)}(\mathcal{Z}_{\mathcal{S}}) = \widetilde{H}^{-1}(\varnothing) = \mathbb{Z} & & 1\\
H^{-1,(0,0,2,2,0)}(\mathcal{Z}_{\mathcal{S}}) = \widetilde{H}^{0}(F_3 \cup F_4) = \mathbb{Z} & & u_3v_4\\
H^{-1,(0,0,2,0,2)}(\mathcal{Z}_{\mathcal{S}}) = \widetilde{H}^{0}(F_3 \cup F_5) = \mathbb{Z} & & u_5v_3\\
H^{-1,(0,0,0,2,2)}(\mathcal{Z}_{\mathcal{S}}) = \widetilde{H}^{0}(F_4 \cup F_5) = \mathbb{Z} & & u_4v_5\\
H^{-2,(0,0,2,2,2)}(\mathcal{Z}_{\mathcal{S}}) = \widetilde{H}^{0}(F_3 \cup F_4 \cup F_5) = \mathbb{Z} \oplus \mathbb{Z} & & u_5u_3v_4,\ u_5u_4v_3\\
H^{0,(2,2,0,0,0)}(\mathcal{Z}_{\mathcal{S}}) = \widetilde{H}^{1}(F_1 \cup F_2) = \mathbb{Z} \oplus \mathbb{Z} & & v_e,\ v_f\\
H^{-1,(2,2,2,0,0)}(\mathcal{Z}_{\mathcal{S}}) = \widetilde{H}^{1}(F_1 \cup F_2 \cup F_3) = \mathbb{Z} & & u_3v_e\\
H^{-1,(2,2,0,2,0)}(\mathcal{Z}_{\mathcal{S}}) = \widetilde{H}^{1}(F_1 \cup F_2 \cup F_4) = \mathbb{Z} & & u_4v_f\\
H^{-1,(2,2,0,0,2)}(\mathcal{Z}_{\mathcal{S}}) = \widetilde{H}^{1}(F_1 \cup F_2 \cup F_5) = \mathbb{Z} & & u_5v_g\\
H^{-2,(2,2,2,2,2)}(\mathcal{Z}_{\mathcal{S}}) = \widetilde{H}^{2}(F_1 \cup \cdots \cup F_5) = \mathbb{Z} & & u_5u_4v_3v_f = u_5u_4v_\sigma
\end{array}$$

It follows that the ordinary (1-graded) Betti numbers of $\mathcal{Z}_{\mathcal{S}}$ are given by the sequence $(1,0,0,3,4,3,0,0,1)$. In the right column of the table above we include the cocycles in the differential graded ring $\Lambda[u_1,\ldots,u_5] \otimes \mathbb{Z}[\mathcal{S}]$ representing generators of the corresponding cohomology group. This allows us to determine the ring structure in $H^*(\mathcal{Z}_{\mathcal{S}})$. For example,

$$[u_5u_3v_4] \cdot [v_f] = [u_5u_3v_4v_f] = 0 = [u_5u_4v_3] \cdot [v_e].$$

On the other hand,

$$\begin{aligned}
[u_5u_3v_4] \cdot [v_e] = -[u_3u_5v_4v_e] = -[u_3u_4v_5v_e]&\\
= [u_3u_4v_5v_f] = [u_5u_4v_3v_f] = [u_5u_4v_3] \cdot [v_f]&.
\end{aligned}$$

Here we have used the relations $d(u_3u_4u_5v_e) = u_3u_4v_5v_e - u_3u_5v_4v_e$ and $d(u_1u_3u_4v_2v_5) = u_3u_4v_5v_e + u_3u_4v_5v_f$. All nontrivial products come from Poincaré duality. These calculations are summarised by the cohomology ring isomorphism

$$H^*(\mathcal{Z}_{\mathcal{S}}) \cong H^*\bigl((S^3 \times S^5)^{\#3} \,\#\, (S^4 \times S^4)^{\#2}\bigr)$$

where the manifold on the right hand side is the connected sum of three copies of $S^3 \times S^5$ and two copies of $S^4 \times S^4$. We expect that the isomorphism above is induced by a homeomorphism.

Exercises.

4.9.8. Generalise Proposition 4.2.1 to simplicial posets, i.e. establish a ring isomorphism $H^*\bigl((\mathbb{C}P^\infty, pt)^{\mathcal{S}}\bigr) \cong \mathbb{Z}[\mathcal{S}]$.

CHAPTER 5

Toric varieties and manifolds

A toric variety is an algebraic variety on which an *algebraic torus* $(\mathbb{C}^\times)^n$ acts with a dense (Zariski open) orbit. An algebraic torus contains a (compact) torus T^n, so toric varieties are toric spaces in our usual sense. Toric varieties are described by combinatorial-geometric objects, rational fans (see Section 2.1), and the combinatorics of the fan determines the orbit structure of the torus action.

Toric varieties were introduced in 1970 in the pioneering work of Demazure on *Cremona group*. The geometry of toric varieties, or *toric geometry*, very quickly became one of the most fascinating topics in algebraic geometry and found applications in many other mathematical sciences, sometimes distant from each other. We have already mentioned the proof for the necessity part of the g-theorem for simplicial polytopes given by Stanley. Other remarkable applications include counting lattice points and volumes of lattice polytopes; relations with Newton polytopes and the number of solutions of a system of algebraic equations (after Khovanskii and Kushnirenko); discriminants, resultants and hypergeometric functions (after Gelfand, Kapranov and Zelevinsky); reflexive polytopes and mirror symmetry for Calabi–Yau toric hypersurfaces and complete intersections (after Batyrev). Standard references on toric geometry include Danilov's survey [**68**] and books by Oda [**173**], Fulton [**96**] and Ewald [**86**]. The most recent exhaustive account by Cox, Little and Schenck [**66**] covers many new applications, including those mentioned above. Without attempting to give another review of toric geometry, in this chapter we collect the basic definitions and constructions, and emphasise topological and combinatorial aspects of toric varieties.

We review the three main approaches to toric varieties in the appropriate sections: the 'classical' construction via fans, the 'algebraic quotient' construction, and the 'symplectic reduction' construction. The intersection of Hermitian quadrics appearing as an intermediate stage in the symplectic construction of toric varieties (the level set for the moment map) links toric geometry to moment-angle complexes. This link will be developed further in the next chapter. In the last section we consider the topological properties which will be taken as the base for different topological generalisations of toric varieties in the later chapters.

A basic knowledge of algebraic geometry would much help the reader of this chapter, although it is not absolutely necessary.

5.1. Classical construction from rational fans

An *algebraic torus* T^{a} is a commutative complex algebraic group isomorphic to a product $(\mathbb{C}^\times)^n$ of copies of the multiplicative group $\mathbb{C}^\times = \mathbb{C} \setminus \{0\}$. It contains a compact torus T^n as a Lie (but not algebraic) subgroup.

We shall often identify an algebraic torus with the standard model $(\mathbb{C}^\times)^n$.

Definition 5.1.1. A *toric variety* is a normal complex algebraic variety V containing an algebraic torus $(\mathbb{C}^\times)^n$ as a Zariski open subset in such a way that the natural action of $(\mathbb{C}^\times)^n$ on itself extends to an action on V.

It follows that $(\mathbb{C}^\times)^n$ acts on V with a dense orbit. Toric varieties originally appeared as equivariant compactifications of an algebraic torus, although non-compact (e.g., affine) examples are now of equal importance.

Example 5.1.2. The algebraic torus $(\mathbb{C}^\times)^n$ and the affine space $\mathbb{C}^n$ are the simplest examples of toric varieties. A compact example is given by the projective space $\mathbb{C}P^n$ on which the torus acts in homogeneous coordinates as follows:

$$(t_1,\ldots,t_n)\cdot(z_0:z_1:\ldots:z_n)=(z_0:t_1z_1:\ldots:t_nz_n).$$

Algebraic geometry of toric varieties is translated completely into the language of combinatorial and convex geometry. Namely, there is a bijective correspondence between rational fans in an n-dimensional space (see Section 2.1) and complex n-dimensional toric varieties. Under this correspondence,

$$\begin{aligned}\text{cones} &\longleftrightarrow \text{affine varieties}\\ \text{complete fans} &\longleftrightarrow \text{compact varieties}\\ \text{normal fans of polytopes} &\longleftrightarrow \text{projective varieties}\\ \text{regular fans} &\longleftrightarrow \text{nonsingular varieties}\\ \text{simplicial fans} &\longleftrightarrow \text{orbifolds}\end{aligned}$$

We review this construction below; the details can be found in the sources mentioned above. Following the algebraic tradition, we use the coordinate-free notation here. We consider a lattice N of rank n (isomorphic to the standard lattice $\mathbb{Z}^n$), and denote by $N_\mathbb{R}$ its ambient n-dimensional real vector space $N\otimes_\mathbb{Z}\mathbb{R}\cong\mathbb{R}^n$. We also define the algebraic torus $T_N^{\mathrm{a}}=N\otimes_\mathbb{Z}\mathbb{C}^\times\cong(\mathbb{C}^\times)^n$. All cones and fans in this chapter are rational.

Construction 5.1.3. We first describe how to assign an affine toric variety to a cone $\sigma\subset N_\mathbb{R}$. Consider the dual cone $\check{\sigma}\subset N^*_\mathbb{R}$ (see (2.1)) and denote by

$$S_\sigma=\check{\sigma}\cap N^*$$

the set of its lattice points. Then S_σ is a finitely generated semigroup (with respect to addition). Let $A_\sigma=\mathbb{C}[S_\sigma]$ be the semigroup ring of S_σ. It is a commutative finitely generated $\mathbb{C}$-algebra, with a $\mathbb{C}$-vector space basis $\{\chi^{\boldsymbol{u}}\colon \boldsymbol{u}\in S_\sigma\}$. The multiplication in A_σ is defined via the addition in S_σ:

$$\chi^{\boldsymbol{u}}\cdot\chi^{\boldsymbol{u}'}=\chi^{\boldsymbol{u}+\boldsymbol{u}'},$$

so χ^0 is the unit. The *affine toric variety* V_σ corresponding to σ is the affine algebraic variety corresponding to A_σ:

$$V_\sigma=\operatorname{Spec}(A_\sigma).$$

By choosing a multiplicative generator set in A_σ we represent it as a quotient

$$A_\sigma=\mathbb{C}[x_1,\ldots,x_r]/\mathcal{I};$$

then the variety V_σ is the common zero set of polynomials from the ideal $\mathcal{I}$. Each point of V_σ corresponds to a semigroup homomorphism $\operatorname{Hom}_{\mathrm{sg}}(S_\sigma,\mathbb{C}_\mathrm{m})$, where $\mathbb{C}_\mathrm{m}=\mathbb{C}^\times\cup\{0\}$ is the multiplicative semigroup of complex numbers.

Now if τ is a face of σ, then $\check{\sigma} \subset \check{\tau}$, and the inclusion of semigroup algebras $\mathbb{C}[S_\sigma] \to \mathbb{C}[S_\tau]$ induces a morphism $V_\tau \to V_\sigma$, which is an inclusion of a Zariski open subset. This allows us to glue the affine varieties V_σ corresponding to all cones σ in a fan Σ into an algebraic variety V_Σ, referred to as the *toric variety* corresponding to the fan Σ. More formally, V_Σ may be defined as the *colimit* of algebraic varieties V_σ over the partially ordered set of cones of Σ:

$$V_\Sigma = \operatorname*{colim}_{\sigma \in \Sigma} V_\sigma.$$

Here is the crucial point: the fact that the cones σ patch into a fan Σ guarantees that the variety V_Σ obtained by gluing the pieces V_σ is Hausdorff in the usual topology. In algebraic geometry, the Hausdorffness is replaced by the related notion of separatedness: a variety V is *separated* if the image of the diagonal map $\varDelta \colon V \to V \times V$ is Zariski closed. A separated variety is Hausdorff in the usual topology.

LEMMA 5.1.4. *If a collection of cones $\{\sigma\}$ forms a fan Σ, then the variety $V_\Sigma = \operatorname{colim}_{\sigma \in \Sigma} V_\sigma$ is separated.*

PROOF. Using the separatedness criterion of [**201**, Ch. V, §4.3] (see also [**68**, Proposition 5.4]), it is enough to verify the following: if cones σ and σ' intersect in a common face τ, then the diagonal map $V_\tau \to V_\sigma \times V_{\sigma'}$ is a closed embedding. This is equivalent to the assertion that the natural homomorphism $A_\sigma \otimes A_{\sigma'} \to A_\tau$ is surjective. To prove this, we use Separation Lemma (Lemma 2.1.2). According to it, there is a linear function $\boldsymbol{u}$ which is nonnegative on σ, nonpositive on σ' and the intersection of the hyperplane $\boldsymbol{u}^\perp$ with σ is τ. Now take $\boldsymbol{u}' \in \check{\tau}$, i.e. $\boldsymbol{u}'$ is nonnegative on τ. Then there is an integer $k \geqslant 0$ such that $\boldsymbol{u}' + k\boldsymbol{u}$ is nonnegative on σ, i.e. $\boldsymbol{u}' + k\boldsymbol{u} \in \check{\sigma}$. Then $\boldsymbol{u}' = (\boldsymbol{u}' + k\boldsymbol{u}) + (-k\boldsymbol{u}) \in \check{\sigma} + \check{\sigma}'$. It follows that $S_\sigma \oplus S_{\sigma'} \to S_\tau$ is surjective map of vector spaces (or semigroups), hence $A_\sigma \otimes A_{\sigma'} \to A_\tau$ is a surjective homomorphism. $\square$

We shall consider only separated varieties in what follows.

The variety V_σ carries an algebraic action of the torus $T_N^{\mathrm{a}} = N \otimes_{\mathbb{Z}} \mathbb{C}^\times$

$$T_N^{\mathrm{a}} \times V_\sigma \to V_\sigma, \quad (\boldsymbol{t}, x) \mapsto \boldsymbol{t} \cdot x \tag{5.1}$$

which is defined as follows. A point $\boldsymbol{t} \in T_N^{\mathrm{a}}$ is determined by a group homomorphism $N^* \to \mathbb{C}^\times$. In coordinates, the homomorphism $\mathbb{Z}^n \cong N^* \to \mathbb{C}^\times$ corresponding to $\boldsymbol{t} = (t_1, \ldots, t_n)$ is given by

$$\boldsymbol{u} = (u_1, \ldots, u_n) \mapsto \boldsymbol{t}(\boldsymbol{u}) = (t_1^{u_1} \cdots t_n^{u_n}).$$

A point $x \in V_\sigma$ corresponds to a semigroup homomorphism $S_\sigma \to \mathbb{C}_{\mathrm{m}}$. Then we define $\boldsymbol{t} \cdot x$ as the point in V_σ corresponding to the semigroup homomorphism $S_\sigma \to \mathbb{C}_{\mathrm{m}}$ given by

$$\boldsymbol{u} \mapsto t(\boldsymbol{u}) x(\boldsymbol{u}).$$

The homomorphism of algebras $A_\sigma \to A_\sigma \otimes \mathbb{C}[N^*]$ dual to the action (5.1) maps $\chi^{\boldsymbol{u}}$ to $\chi^{\boldsymbol{u}} \otimes \chi^{\boldsymbol{u}}$ for $\boldsymbol{u} \in S_\sigma$. If $\sigma = \{\boldsymbol{0}\}$, then we obtain the multiplication in the algebraic group T_N^{a}. The actions on the varieties V_σ are compatible with the inclusions of open sets $V_\tau \to V_\sigma$ corresponding to the inclusions of faces $\tau \subset \sigma$. Therefore, for each fan Σ we obtain a T_N^{a}-action on the variety V_Σ, which extends the T_N^{a}-action on itself.

EXAMPLE 5.1.5. Let $N = \mathbb{Z}^n$ and let σ be the cone spanned by the first k basis vectors $\boldsymbol{e}_1, \ldots, \boldsymbol{e}_k$, where $0 \leqslant k \leqslant n$. The semigroup $S_\sigma = \sigma^\vee \cap N^*$ is generated by the dual elements $\boldsymbol{e}_1^*, \ldots, \boldsymbol{e}_k^*$ and $\pm\boldsymbol{e}_{k+1}^*, \ldots, \pm\boldsymbol{e}_n^*$. Therefore,

$$A_\sigma \cong \mathbb{C}[x_1, \ldots, x_k, x_{k+1}, x_{k+1}^{-1}, \ldots, x_n, x_n^{-1}],$$

where we set $x_i = \chi^{\boldsymbol{e}_i^*}$. It follows that the corresponding affine variety is

$$V_\sigma \cong \mathbb{C} \times \cdots \times \mathbb{C} \times \mathbb{C}^\times \times \cdots \times \mathbb{C}^\times = \mathbb{C}^k \times (\mathbb{C}^\times)^{n-k}.$$

In particular, for $k = n$ we obtain an n-dimensional affine space, and for $k = 0$ (i.e. $\sigma = \{0\}$) we obtain the algebraic torus $(\mathbb{C}^\times)^n$.

EXAMPLE 5.1.6. Let $\sigma \subset \mathbb{R}^2$ be the cone generated by the vectors $\boldsymbol{e}_2$ and $2\boldsymbol{e}_1 - \boldsymbol{e}_2$ (note that these two vectors do not span $\mathbb{Z}^2$, so this cone is not regular). The dual cone $\sigma^\vee$ is generated by $\boldsymbol{e}_1^*$ and $\boldsymbol{e}_1^* + 2\boldsymbol{e}_2^*$. The semigroup S_σ is generated by $\boldsymbol{e}_1^*$, $\boldsymbol{e}_1^* + \boldsymbol{e}_2^*$ and $\boldsymbol{e}_1^* + 2\boldsymbol{e}_2^*$, with one relation among them. Therefore,

$$A_\sigma = \mathbb{C}[x, xy, xy^2] \cong \mathbb{C}[u, v, w]/(v^2 - uw)$$

and V_σ is a quadratic cone (a singular variety).



FIGURE 5.1. Complete fans in $\mathbb{R}^2$

EXAMPLE 5.1.7. Let Σ be the complete fan in $\mathbb{R}^2$ with the following three maximal cones: the cone σ_0 generated by $\boldsymbol{e}_1$ and $\boldsymbol{e}_2$, the cone σ_1 generated by $\boldsymbol{e}_2$ and $-\boldsymbol{e}_1 - \boldsymbol{e}_2$, and the cone σ_2 generated by $-\boldsymbol{e}_1 - \boldsymbol{e}_2$ and $\boldsymbol{e}_1$, see Fig. 5.1 (a). Then each affine variety V_{σ_i} is isomorphic to $\mathbb{C}^2$, with coordinates (x, y) for σ_0, $(x^{-1}, x^{-1}y)$ for σ_1, and (y^{-1}, xy^{-1}) for σ_2. These three affine charts glue together into the complex projective plane $V_\Sigma = \mathbb{C}P^2$ in the standard way: if $(z_0 : z_1 : z_2)$ are the homogeneous coordinats in $\mathbb{C}P^2$, then we have $x = z_1/z_0$ and $y = z_2/z_0$.

EXAMPLE 5.1.8. Fix $k \in \mathbb{Z}$ and consider the complete fan in $\mathbb{R}^2$ with the four two-dimensional cones generated by the pairs of vectors $(\boldsymbol{e}_1, \boldsymbol{e}_2)$, $(\boldsymbol{e}_1, -\boldsymbol{e}_2)$, $(-\boldsymbol{e}_1 + k\boldsymbol{e}_2, -\boldsymbol{e}_2)$ and $(-\boldsymbol{e}_1 + k\boldsymbol{e}_2, \boldsymbol{e}_2)$, see Fig 5.1 (b). It can be shown that the corresponding toric variety F_k is the projectivisation $\mathbb{C}P(\underline{\mathbb{C}} \oplus \mathcal{O}(k))$ of the sum of a trivial line bundle $\underline{\mathbb{C}}$ and the kth power $\mathcal{O}(k) = \gamma^{\otimes k}$ of the canonical line bundle γ over $\mathbb{C}P^1$ (an exercise). These 2-dimensional complex varieties F_k are known as *Hirzebruch surfaces.*

The inclusion poset of closures of T_N^{a}-orbits of V_Σ is isomorphic to the reversed inclusion poset of faces of Σ. That is, k-dimensional cones of Σ correspond to codimension-k orbits of the algebraic torus action on V_Σ. In particular,

n-dimensional cones correspond to fixed points, and the apex (the zero cone) corresponds to the dense orbit. Furthermore, if a subcollection of cones of Σ forms a fan Σ', then the toric variety $V_{\Sigma'}$ is embedded into V_Σ as a Zariski open subset.

We recall from Section 2.1 that a fan is called simplicial (respectively, regular) if each of its cones is generated by a part of basis of the space $N_\mathbb{R}$ (respectively, of the lattice N), and a fan is called complete if the union of its cones is the whole $N_\mathbb{R}$.

A toric variety V_Σ is compact (in usual topology) if and only if the fan Σ is complete. If Σ is a simplicial fan, then V_Σ is an *orbifold*, that is, it is locally isomorphic to a quotient of $\mathbb{C}^n$ by a finite group action. A toric variety V_Σ is nonsingular (smooth) if and only if the fan Σ is regular.

Exercises.

5.1.9. Let Σ be the 'multifan' in $\mathbb{R}^1$ consisting of two identical 1-dimensional cones generated by $\boldsymbol{e}_1$ and a 0-dimensional cone $\mathbf{0}$. Describe the algebraic variety V_Σ obtained by gluing the affine varieties corresponding to this 'multifan' and show that V_Σ is not separated (or non-Hausdorff in the usual topology).

5.1.10. Describe the toric variety corresponding to the fan with 3 one-dimensional cones generated by the vectors $\boldsymbol{e}_1$, $\boldsymbol{e}_2$ and $-\boldsymbol{e}_1 - \boldsymbol{e}_2$.

5.1.11. Show that the toric variety of Example 5.1.8 is isomorphic to the Hirzebruch surface $F_k = \mathbb{C}P(\underline{\mathbb{C}} \oplus \mathcal{O}(k))$.

5.1.12. Show that the Hirzebruch surface F_k is homeomorphic to $S^2 \times S^2$ for even k and is homeomorphic to $\mathbb{C}P^2 \# \overline{\mathbb{C}P}^2$ for odd k, where $\#$ denotes the connected sum, and $\overline{\mathbb{C}P}^2$ is $\mathbb{C}P^2$ with the orientation reversed.

5.2. Projective toric varieties and polytopes

Construction 5.2.1 (projective toric varieties). Let P be a convex polytope with vertices in the dual lattice N^* (a *lattice polytope*), and let Σ_P be the normal fan of P (see Construction 2.1.3). Since $P \subset N^*_\mathbb{R}$, the fan Σ_P belongs to the space $N_\mathbb{R}$. It has a maximal cone σ_v for each vertex $v \in P$. The dual cone σ_v^* is the "vertex cone" at v, generated by all vectors pointing from v to other points of P.

Define the toric variety $V_P = V_{\Sigma_P}$. Since the normal fan Σ_P does not depend on the linear size of the polytope, we may assume that for each vertex v the semigroup S_{σ_v} is generated by the lattice points of the polytope (this can always be achieved by replacing P by kP with sufficiently large k). Since N^* is the lattice of characters of the algebraic torus $T_N^{\mathrm{a}} = N \otimes_\mathbb{Z} \mathbb{C}^\times$ (that is, $N = \mathrm{Hom}_\mathbb{Z}(T_N^{\mathrm{a}}, \mathbb{C}^\times)$), the lattice points of the polytope $P \subset N^*$ define an embedding of T_N^{a} in the torus $(\mathbb{C}^\times)^{|N^* \cap P|}$, where $|N^* \cap P|$ is the number of lattice points in P. It can be shown that the toric variety V_P is the projective closure of the torus T_N^{a} under this embedding (see [**66**, §2.2] or [**96**, §3.4]). In particular, the variety V_P is *projective*, that is, it can be given by a set of homogeneous equations in $\mathbb{C}P^{|N^* \cap P|}$. It follows that toric varieties arising from polytopes are projective. The converse is also true: the fan corresponding to a projective toric variety is the normal fan of a lattice polytope.

The polytope P carries more geometric information than the normal fan Σ_P: different lattice polytopes with the same normal fan Σ correspond to different projective embeddings of the toric variety V_Σ.

Nonsingular projective toric varieties correspond to lattice polytopes P which are simple and Delzant (that is, for each vertex v, the normal vectors of facets meeting at v form a basis of the lattice N).

Example 5.2.2. The fan Σ described in Example 2.1.4 is regular, but cannot be obtained as the normal fan of a simple polytope. The corresponding 3-dimensional toric variety V_Σ is compact and nonsingular, but not projective.

We note that although the fan Σ from the previous example cannot be realised *geometrically* as the fan over the faces of a simplicial polytope (or, equivalently, as the normal fan of a simple polytope), its underlying simplicial complex $\mathcal{K}_\Sigma$ is nevertheless *combinatorially* equivalent to the boundary complex of a simplicial polytope (namely, an octahedron with a pyramid over one of its facets, see Fig. 2.1). In other words, using the terminology of Section 2.5, the star-shaped sphere triangulation $\mathcal{K}_\Sigma$ is polytopal (in the combinatorial sense).

There are, of course, nonpolytopal star-shaped spheres, such as the Barnette sphere $\mathcal{K}$ (see Example 2.5.8). It is easy to see that a simplicial fan realising the Barnette sphere can be chosen rational. However, to realise a nonpolytopal sphere by a *regular* fan is a more difficult task. In fact, not such examples are known:

Problem 5.2.3. Does there exist a complete regular fan Σ whose underlying complex $\mathcal{K}_\Sigma$ is a nonpolytopal sphere triangulation?

This question is also important for the study of *quasitoric manifolds* and other topological generalisations of toric varieties discussed in the later chapters.

We observe that each combinatorial simple polytope admits a convex realisation as a lattice polytope. Indeed by a small perturbation of the defining inequalities in (1.1) we can make all of them rational (that is, with rational $\boldsymbol{a}_i$ and b_i). Such a perturbation does not change the combinatorial type, as the half-spaces defined by the inequalities are in general position. As a result, we obtain a simple polytope P' of the same combinatorial type with rational vertex coordinates. To get a lattice polytope (say, with vertices in the standard lattice $\mathbb{Z}^n$) we just take the magnified polytope kP' for appropriate $k \in \mathbb{Z}$. Similarly, by perturbing the vertices instead of the hyperplanes, we can obtain a lattice realisation for an arbitrary simplicial polytope (and we can obtain a rational fan realisation of any starshaped sphere triangulation). However, this argument do not work for convex polytopes which are neither simple nor simplicial. In fact, there are exist *nonrational* combinatorial polytopes, which cannot be realised with rational vertex coordinates, see [**228**, Example 6.21] and the discussion there.

Toric geometry, even in its topological part, does not translate to a purely combinatorial study of fans and polytopes: the underlying convex geometry is what really matters. This is illustrated by the simple observation that different realisations of a combinatorial polytope by lattice polytopes often produce different (even topologically) toric varieties:

Example 5.2.4. The complete regular fan Σ_k corresponding to the Hirzebruch surface F_k (see Fig. 5.1 (b)) is the normal fan of a lattice quadrilateral (trapezoid), e.g. given by

$$P_k = \{(x_1, x_2) \in \mathbb{R}^2 \colon x_1 \geqslant 0,\ x_2 \geqslant 0,\ -x_1 + kx_2 \geqslant -1,\ -x_2 \geqslant -1\}.$$

The polytopes P_k corresponding to different k are combinatorially equivalent (as they are all quadrilaterals), but the topology of the corresponding toric varieties F_k is different for even and odd k (see Exercise 5.1.12).

Furthermore, there exist combinatorial simple polytopes that do not admit any lattice realisation P with smooth V_P:

EXAMPLE 5.2.5 ([**70**, 1.22]). Let P be the dual of a 2-neighbourly simplicial n-polytope (e.g., a cyclic polytope of dimension $n \geqslant 4$, see Example 1.1.18) with $m \geqslant 2^n$ vertices. Then for any lattice realisation of P the corresponding normal fan Σ_P is not regular, and the toric variety V_P is singular. Indeed, assume that the normal fan Σ_P is regular. Since the 1-skeleton of $\mathcal{K}_P$ is a complete graph, each pair of primitive generators $\boldsymbol{a}_i, \boldsymbol{a}_j$ of one-dimensional cones of Σ_P is a part of basis of $\mathbb{Z}^n$, and therefore $\boldsymbol{a}_i$ and $\boldsymbol{a}_j$ must be different modulo 2. This is a contradiction, since the number of different nonzero vectors in $\mathbb{Z}_2^n$ is $2^n - 1$.

Exercises.

5.2.6. Describe explicitly a rational fan realising the Barnette sphere by writing down its primitive integral generator vectors.

5.2.7. Write down a system of homogeneous equations defining each Hirzebruch surface in a projective space (hint: use Construction 5.2.1).

5.3. Cohomology of toric manifolds

A *toric manifold* is a smooth compact toric variety. (Compactness will be always understood in the sense of usual topology; it corresponds to algebraic geometer's notion of *completeness.*) Toric manifolds V_Σ correspond to complete regular fans Σ. *Projective toric manifolds* V_P correspond to lattice polytopes P whose normal fans are regular.

The cohomology of a toric manifold V_Σ can be calculated effectively from the fan Σ. The Betti numbers are determined by the combinatorics of Σ only, while the ring structure of $H^*(V_\Sigma)$ depends on the geometric data. The required combinatorial ingredients are the h-vector $\boldsymbol{h}(\mathcal{K}_\Sigma) = (h_0, h_1, \dots, h_n)$ (see Definition 2.2.5) of the underlying simplicial complex $\mathcal{K}_\Sigma$ and its face ring $\mathbb{Z}[\mathcal{K}_\Sigma]$ (Definition 3.1.1). The geometric data consists of the primitive generators $\boldsymbol{a}_1, \dots, \boldsymbol{a}_m$ of one-dimensional cones (edges) of Σ.

THEOREM 5.3.1 (Danilov–Jurkiewicz). *Let Σ be a complete regular fan in $N_\mathbb{R} \cong \mathbb{R}^n$, and V_Σ the corresponding toric manifold. Then*

(a) *the homology groups of V_Σ vanish in odd dimensions, and are free abelian in even dimensions, with ranks given by*

$$b_{2i}(V_\Sigma) = h_i(\mathcal{K}_\Sigma),$$

where $h_i(\mathcal{K}_\Sigma)$, $i = 0, 1, \dots, n$, are the components of the h-vector of $\mathcal{K}_\Sigma$;

(b) *the cohomology ring of V_Σ is given by*

$$H^*(V_\Sigma; \mathbb{Z}) = \mathbb{Z}[v_1, \dots, v_m]/(\mathcal{I}_{\mathcal{K}_\Sigma} + \mathcal{J}_\Sigma) = \mathbb{Z}[\mathcal{K}_\Sigma]/\mathcal{J}_\Sigma,$$

where the v_is are the 2-dimensional cohomology classes dual to the invariant divisors D_i corresponding to the edges of Σ, and $\mathcal{J}_\Sigma$ is the ideal generated by all linear forms $\sum_{j=1}^m \langle \boldsymbol{u}, \boldsymbol{a}_j \rangle v_j$, $\boldsymbol{u} \in N^*$.

This theorem was proved by Jurkiewicz for projective toric manifolds and by Danilov [**68**, Theorem 10.8] in the general case. We shall give a topological proof of a more general result in Section 7.2.

To obtain an explicit presentation of the ring $H^*(V_\Sigma)$ we choose a basis of N and write the vectors $\boldsymbol{a}_j$ in coordinates: $\boldsymbol{a}_j = (a_{j1}, \ldots, a_{jn})^t$, $1 \leqslant j \leqslant m$. Then the ideal J_Σ is generated by the n linear forms

$$t_i = a_{1i}v_1 + \cdots + a_{mi}v_m \in \mathbb{Z}[v_1, \ldots, v_m], \quad 1 \leqslant i \leqslant n.$$

By Lemma 3.3.2, the sequence $t_1, \ldots, t_n$ is an lsop in the Cohen–Macaulay ring $\mathbb{Z}[\mathcal{K}_\Sigma]$, so it is a regular sequence. Hence, $\mathbb{Z}[\mathcal{K}_\Sigma]$ is a free $\mathbb{Z}[t_1, \ldots, t_n]$-module, and statement (a) of Theorem 5.3.1 follows from (b) and Theorem 3.1.10.

Remark. Theorem 5.3.1 remains valid for complete simplicial fans and corresponding toric orbifolds if the integer coefficients are replaced by the rationals [**68**]. The integral cohomology of toric orbifolds often has torsion, and the ring structure is subtle even in the simplest case of weighted projective spaces [**129**], [**13**].

It follows from Theorem 5.3.1 that the cohomology ring of V_Σ is generated by two-dimensional classes. This is the first property to check if one wishes to determine whether a given algebraic variety or smooth manifold has a structure of a toric manifold. For instance, this rules out flag varieties and Grassmanians different from projective spaces. Another important property of toric manifolds is that the *Chow ring* of V_Σ coincides with its integer cohomology ring [**96**, § 5.1].

Assume now that $\Sigma = \Sigma_P$ is the normal fan of a lattice polytope $P = P(A, \boldsymbol{b})$ given by (1.1). Let V_P be the corresponding projective toric variety, see Construction 5.2.1. In the notation of Theorem 5.3.1, the linear combination $D_P = b_1D_1 + \cdots + b_mD_m$ is an *ample divisor* on V_P (see, e.g., [**66**, Proposition 6.1.10]). This means that, when k is sufficiently large, kD_P is a hyperplane section divisor for a projective embedding $V_P \subset \mathbb{C}P^r$. In fact, the space of sections $H^0(V_P, kD_P)$ of (the line bundle corresponding to) kD_P has basis corresponding to the lattice points in kP. One may take k so that kP has 'enough lattice points' to get an embedding of V_P into the projectivisation of $H^0(V_P, kD_P)$; this is exactly the embedding described in Construction 5.2.1. Let $\omega = b_1v_1 + \cdots + b_mv_m \in H^2(V_P; \mathbb{C})$ be the complex cohomology class of D_P.

Theorem 5.3.2 (Hard Lefschetz theorem for toric orbifolds). *Let P be a lattice simple polytope* (1.1)*, let V_P be the corresponding projective toric variety, and let $\omega = b_1v_1 + \cdots + b_mv_m \in H^2(V_P; \mathbb{C})$ be the class defined above. Then the maps*

$$H^{n-i}(V_P; \mathbb{C}) \xrightarrow{\omega^i} H^{n+i}(V_P; \mathbb{C})$$

are isomorphisms for all $i = 1, \ldots, n$.

If V_P is smooth, then it is Kähler, and ω is the class of the Kähler 2-form.

The proof of the Hard Lefschetz Theorem is well beyond the scope of this book. In fact, it is a corollary of a more general version of Hard Lefschetz Theorem for the (middle perversity) *intersection cohomology*, which is valid for all projective varieties (not necessary orbifolds). See the discussion in [**96**, §5.2] or [**66**, §12.6].

Now we are ready to give Stanley's argument for the 'only if' part of the g-theorem for simple polytopes:

Proof of the necessity part of Theorem 1.4.14. We need to establish conditions (a)–(c) for a combinatorial simple polytope. Realise it by a lattice polytope $P \subset \mathbb{R}^n$ as described in the previous section. Let V_P be the corresponding toric variety. Part (a) is already proved (Theorem 1.3.4). It follows from Theorem 5.3.2 that the multiplication by $\omega \in H^2(V_P;\mathbb{Q})$ is a monomorphism $H^{2i-2}(V_P;\mathbb{Q}) \to H^{2i}(V_P;\mathbb{Q})$ for $i \leqslant \left[\frac{n}{2}\right]$. This together with part (a) of Theorem 5.3.1 gives that $h_{i-1} \leqslant h_i$ for $0 \leqslant i \leqslant \left[\frac{n}{2}\right]$, thus proving (b). To prove (c), define the graded commutative $\mathbb{Q}$-algebra $A = H^*(V_P;\mathbb{Q})/(\omega)$, where (ω) is the ideal generated by ω. Then $A^0 = \mathbb{Q}$, $A^{2i} = H^{2i}(V_P;\mathbb{Q})/(\omega \cdot H^{2i-2}(V_P;\mathbb{Q}))$ for $1 \leqslant i \leqslant \left[\frac{n}{2}\right]$, and A is generated by degree-two elements (since so is $H^*(V_P;\mathbb{Q})$). It follows from Theorem 1.4.12 that the numbers $\dim A^{2i} = h_i - h_{i-1},\ 0 \leqslant i \leqslant \left[\frac{n}{2}\right]$, are the components of an M-vector, thus proving (c) and the whole theorem. □

Remark. The Dehn–Sommerville equations now can be interpreted as Poincaré duality for V_P. (Even though V_P may be not smooth, the rational cohomology algebra of a toric orbifold satisfies Poincaré duality.)

The Hard Lefschetz Theorem holds for projective varieties only. Therefore, Stanley's argument cannot be generalised to nonpolytopal spheres. However, the cohomology of toric varieties can be used to prove statements generalising the g-theorem in a different direction, namely, to the case of general (not necessarily simple or simplicial) convex polytopes. So suppose P is a convex lattice n-polytope, and V_P is the corresponding projective toric variety. If P is not simple then V_P has worse than orbifold-type singularities and its ordinary cohomology behaves badly. The Betti numbers of V_P are not determined by the combinatorial type of P and do not satisfy Poincaré duality. On the other hand, the dimensions $\widehat{h}_i = \dim IH_{2i}(V_P)$ of the intersection homology groups of V_P are combinatorial invariants of P (see the description in [**206**] or [**66**, §12.5]). The vector

$$\widehat{\boldsymbol{h}}(P) = (\widehat{h}_0, \widehat{h}_1, \ldots, \widehat{h}_n)$$

is called the *intersection h-vector*, or the *toric h-vector* of P. If P is simple, then the toric h-vector coincides with the standard h-vector, but in general $\widehat{\boldsymbol{h}}(P)$ is not determined by the face numbers of P. The toric h-vector satisfies the 'Dehn–Sommerville equations' $\widehat{h}_i = \widehat{h}_{n-i}$, and the Hard Lefschetz Theorem for intersection cohomology shows that it also satisfies the GLBC inequalities:

$$\widehat{h}_0 \leqslant \widehat{h}_1 \leqslant \cdots \leqslant \widehat{h}_{\left[\frac{n}{2}\right]}.$$

In the case when P cannot be realised by a lattice polytope (i.e., when P is nonrational), the toric h-vector can still be defined combinatorially, but the GLBC inequalities require a separate proof. Partial results in this direction were obtained by several people, before the Hard Lefschetz Theorem for nonrational polytopes was eventually proved in the work of Karu [**131**]. This result also gives a purely combinatorial proof of the Hard Lefschetz Theorem for projective toric varieties.

5.4. Algebraic quotient construction

Along with the classical construction of toric varieties from fans, described in Section 5.1, there is an alternative way to define a toric variety: as the quotient of a Zariski open subset in $\mathbb{C}^m$ (more precisely, the complement of a coordinate subspace arrangement) by an action of an abelian algebraic group (a product of an algebraic torus and a finite group). Different versions of this construction, which we refer to as simply the 'quotient construction', have appeared in the work of several authors since the early 1990s. In our exposition we mainly follow the work of Cox [**65**] (and also its modernised exposition in [**66**, Chapter 5]); more historical remarks can be also found in these sources.

Quotients in algebraic geometry. Taking quotients of algebraic varieties by algebraic group actions is tricky for both topological and algebraic reasons. First, as algebraic groups (such as algebraic tori) are not compact, their orbits may be not closed, and the quotients may be non-Hausdorff. Second, even if the quotient is Hausdorff as a topological space, it may fail to be an algebraic variety. This may be remedied to some extent by the notion of the categorical quotient.

Let X be an algebraic variety with an action of an affine algebraic group G. An algebraic variety Y is called a *categorical quotient* of X by the action of G if there exists a morphism $\pi\colon X \to Y$ which is constant on G-orbits of X and has the following universal property: for any morphism $\varphi\colon X \to Z$ which is constant on G-orbits, there is a unique morphism $\widehat{\varphi}\colon Y \to Z$ such that $\widehat{\varphi} \circ \pi = \varphi$. This is described by the diagram

$$\begin{tikzcd}
X \arrow[rr, "\varphi"] \arrow[dr, "\pi"'] & & Z \\
& Y \arrow[ur, dashed, "\widehat{\varphi}"'] &
\end{tikzcd}$$

A categorical quotient Y is unique up to isomorphism, and we denote it by $X/\!\!/G$.

Assume first that $X = \operatorname{Spec} A$ is an affine variety, where $A = \mathbb{C}[X]$ is the algebra of regular functions on X. Let $\mathbb{C}[X]^G$ be the subalgebra of G-invariant functions (i.e. such functions f that $f(gx) = f(x)$ for any $g \in G$ and $x \in X$). If G is an algebraic torus (or any *reductive* affine algebraic group), then $\mathbb{C}[X]^G$ is finitely generated. The corresponding affine variety $\operatorname{Spec} \mathbb{C}[X]^G$ is the categorical quotient $X/\!\!/G$. The quotient morphism $\pi\colon X \to X/\!\!/G$ is dual to the inclusion of algebras $\mathbb{C}[X]^G \to \mathbb{C}[X]$. The morphism π is surjective and induces a one-to-one correspondence between $X/\!\!/G$ and *closed* G-orbits of X (i.e. $\pi^{-1}(x)$ contains a unique closed G-orbit for any $x \in X/\!\!/G$, see [**66**, Proposition 5.0.7]).

Therefore, if all G-orbits of an affine variety X are closed, then the categorical quotient $X/\!\!/G$ is identified as a topological space with the ordinary 'topological' quotient X/G. In algebraic geometry quotients of this type are called *geometric* and also denoted by X/G.

Example 5.4.1. Let $\mathbb{C}^\times$ act on $\mathbb{C} = \operatorname{Spec}(\mathbb{C}[z])$ by scalar multiplication. There are two orbits: the closed orbit 0 and the open orbit $\mathbb{C}^\times$. The topological quotient $\mathbb{C}/\mathbb{C}^\times$ is a non-Hausdorff two-point space.

On the other hand, the categorical quotient $\mathbb{C}/\!\!/\mathbb{C}^\times = \operatorname{Spec}(\mathbb{C}[z]^{\mathbb{C}^\times})$ is a point, since any $\mathbb{C}^\times$-invariant polynomial is constant (and there is only one closed orbit).

Similarly, if $\mathbb{C}^\times$ acts on $\mathbb{C}^n = \operatorname{Spec}(\mathbb{C}[z_1, \ldots, z_n])$ diagonally, then an invariant polynomial satisfies $f(\lambda z_1, \ldots, \lambda z_n) = f(z_1, \ldots, z_n)$ for all $\lambda \in \mathbb{C}^\times$. Such a polynomial must be constant, hence $\mathbb{C}^n/\!\!/\mathbb{C}^\times$ is a point.

In good cases categorical quotients of general varieties X may be constructed by 'gluing from affine pieces' as follows. Assume that G acts on X and $\pi\colon X \to Y$ is a morphism of varieties that is constant on G-orbits. If Y has an open affine cover $Y = \bigcup_\alpha V_\alpha$ such that $\pi^{-1}(V_\alpha)$ is affine and V_α is the categorical quotient (that is, $\pi|_{\pi^{-1}(V_\alpha)}\colon \pi^{-1}(V_\alpha) \to V_\alpha$ is the morphism dual to the inclusion of algebras $\mathbb{C}[\pi^{-1}(V_\alpha)]^G \to \mathbb{C}[\pi^{-1}(V_\alpha)]$), then Y is the categorical quotient $X/\!\!/G$.

Example 5.4.2. Let $\mathbb{C}^\times$ act on $\mathbb{C}^2\setminus\{0\}$ diagonally, where $\mathbb{C}^2 = \operatorname{Spec}(\mathbb{C}[z_0,z_1])$. We have an open affine cover $\mathbb{C}^2\setminus\{0\} = U_0 \cup U_1$, where

$$\begin{aligned} U_0 &= \mathbb{C}^2 \setminus \{z_0 = 0\} = \mathbb{C}^\times \times \mathbb{C} = \operatorname{Spec}(\mathbb{C}[z_0^{\pm 1}, z_1]),\\ U_1 &= \mathbb{C}^2 \setminus \{z_1 = 0\} = \mathbb{C} \times \mathbb{C}^\times = \operatorname{Spec}(\mathbb{C}[z_0, z_1^{\pm 1}]),\\ U_0 \cap U_1 &= \mathbb{C}^2 \setminus \{z_0 z_1 = 0\} = \mathbb{C}^\times \times \mathbb{C}^\times = \operatorname{Spec}(\mathbb{C}[z_0^{\pm 1}, z_1^{\pm 1}]). \end{aligned}$$

The algebras of $\mathbb{C}^\times$-invariant functions are

$$\mathbb{C}[z_0^{\pm1}, z_1]^{\mathbb{C}^\times} = \mathbb{C}[z_1/z_0], \quad \mathbb{C}[z_0, z_1^{\pm1}]^{\mathbb{C}^\times} = \mathbb{C}[z_0/z_1], \quad \mathbb{C}[z_0^{\pm1}, z_1^{\pm1}]^{\mathbb{C}^\times} = \mathbb{C}[(z_1/z_0)^{\pm1}].$$

It follows that $V_0 = U_0/\!\!/\mathbb{C}^\times \cong \mathbb{C}$ and $V_1 = U_1/\!\!/\mathbb{C}^\times \cong \mathbb{C}$ glue together along $V_0 \cap V_1 = (U_0 \cap U_1)/\!\!/\mathbb{C}^\times \cong \mathbb{C}^\times$ in the standard way to produce $\mathbb{C}P^1$. All $\mathbb{C}^\times$-orbits are closed in $\mathbb{C}^2\setminus\{0\}$, hence $\mathbb{C}P^1 = (\mathbb{C}^2\setminus\{0\})/\mathbb{C}^\times$ is the geometric quotient.

Similarly, $\mathbb{C}P^n = (\mathbb{C}^{n+1}\setminus\{0\})/\mathbb{C}^\times$ is the geometric quotient for the diagonal action of $\mathbb{C}^\times$.

Example 5.4.3. Now we let $\mathbb{C}^\times$ act on $\mathbb{C}^2\setminus\{0\}$ by $\lambda\cdot(z_0,z_1) = (\lambda z_0, \lambda^{-1} z_1)$. Using the same affine cover of $\mathbb{C}^2\setminus\{0\}$ as in the previous example, we obtain the following algebras of $\mathbb{C}^\times$-invariant functions:

$$\mathbb{C}[z_0^{\pm1}, z_1]^{\mathbb{C}^\times} = \mathbb{C}[z_0 z_1], \quad \mathbb{C}[z_0, z_1^{\pm1}]^{\mathbb{C}^\times} = \mathbb{C}[z_0 z_1], \quad \mathbb{C}[z_0^{\pm1}, z_1^{\pm1}]^{\mathbb{C}^\times} = \mathbb{C}[(z_0 z_1)^{\pm1}].$$

This times gluing together V_0 and V_1 along $V_0 \cap V_1 \cong \mathbb{C}^\times$ gives the variety obtained from two copies of $\mathbb{C}$ by identifying all nonzero points. This variety is nonseparated (the two zeros do not have nonintersecting neighbourhoods in the usual topology). Since we only consider separated varieties in our constructions, this is not a categorical quotient.

A toric variety V_Σ will be described as the categorical (or, in good cases, geometric) quotient of the 'total space' $U(\Sigma)$ by an action of an abelian algebraic group G. We now proceed to describe G and $U(\Sigma)$.

The total space $U(\Sigma)$ and the acting group G. Let Σ be a rational fan in the n-dimensional space $N_\mathbb{R}$ with m one-dimensional cones generated by primitive vectors $\boldsymbol{a}_1, \dots, \boldsymbol{a}_m$. We shall assume that the linear span of $\boldsymbol{a}_1, \dots, \boldsymbol{a}_m$ is the whole $N_\mathbb{R}$. (Equivalently, the toric variety V_Σ does not have torus factors, i.e. cannot be written as $V_\Sigma = V_{\Sigma'} \times \mathbb{C}^\times$. For the general case see [**66**, §5.1].)

We consider the map of lattices $\ell\colon \mathbb{Z}^m \to N$ sending the ith basis vector of $\mathbb{Z}^m$ to $\boldsymbol{a}_i \in N$. Our assumption implies that the corresponding map of algebraic tori,

$$\ell \otimes_\mathbb{Z} \mathbb{C}^\times \colon (\mathbb{C}^\times)^m \to T_N^{\mathrm{a}}$$

is surjective.

Define the group $G = G(\Sigma)$ as the kernel of the map $\ell\otimes_\mathbb{Z}\mathbb{C}^\times$ (which we continue to denote by ℓ for simplicity). We therefore have an exact sequence of groups

$$1 \longrightarrow G \longrightarrow (\mathbb{C}^\times)^m \xrightarrow{\ \ell\ } T_N^{\mathrm{a}} \longrightarrow 1. \tag{5.2}$$

Explicitly, G is given by

$$G = \Bigl\{(z_1, \dots, z_m) \in (\mathbb{C}^\times)^m \colon \prod_{i=1}^m z_i^{\langle \boldsymbol{u}, \boldsymbol{a}_i\rangle} = 1 \quad \text{for all } \boldsymbol{u} \in N^*\Bigr\}. \tag{5.3}$$

The group G is isomorphic to a product of $(\mathbb{C}^\times)^{m-n}$ and a finite abelian group. If Σ is a regular fan with at least one n-dimensional cone, then $G \cong (\mathbb{C}^\times)^{m-n}$.

Given a cone $\sigma \in \Sigma$, we set $g(\sigma) = \{i_1, \dots, i_k\} \subset [m]$ if σ is generated by $\boldsymbol{a}_{i_1} \dots, \boldsymbol{a}_{i_k}$, and consider the monomial $z^{\hat\sigma} = \prod_{j \notin g(\sigma)} z_j$. Define the affine variety

$$A(\Sigma) = \{\boldsymbol{z} \in \mathbb{C}^m \colon z^{\hat\sigma} = 0 \text{ for all } \ \sigma \in \Sigma\}$$

and its complement $U(\Sigma) = \mathbb{C}^m \setminus A(\Sigma)$, which is quasiaffine by definition. It has the affine cover

$$U(\Sigma) = \bigcup_{\sigma \in \Sigma} U(\sigma). \tag{5.4}$$

by affine varieties

$$U(\sigma) = \{\boldsymbol{z} \in \mathbb{C}^m \colon z^{\hat\sigma} \neq 0\} = \{\boldsymbol{z} \in \mathbb{C}^m \colon z_j \neq 0 \ \text{ for } \ j \notin g(\sigma)\} = (\mathbb{C}, \mathbb{C}^\times)^{g(\sigma)}$$

where we used the notation of Construction 4.1.9, so that $U(\sigma) \cong \mathbb{C}^k \times (\mathbb{C}^\times)^{m-k}$. Each subset $U(\sigma) \subset \mathbb{C}^m$ is invariant under the coordinatewise action of $(\mathbb{C}^\times)^m$ on $\mathbb{C}^m$, so that $U(\Sigma)$ is also invariant.

By definition, $A(\Sigma)$ is a union of coordinate subspaces, so we know from Proposition 4.6.2 that its complement $U(\Sigma)$ has the form

$$U(\mathcal{K}) = \mathbb{C}^m \setminus \bigcup_{\{i_1, \dots, i_k\} \notin \mathcal{K}} \bigl\{\boldsymbol{z} \in \mathbb{C}^m \colon z_{i_1} = \dots = z_{i_k} = 0\bigr\} = \mathcal{Z}_{\mathcal{K}}(\mathbb{C}, \mathbb{C}^\times)$$

for some simplicial complex $\mathcal{K}$ on $[m]$. What is this simplicial complex?

The answer is suggested by decomposition (5.4). We define the simplicial complex $\mathcal{K}_\Sigma$ generated by all subsets $g(\sigma) \subset [m]$:

$$\mathcal{K}_\Sigma = \{I \colon I \subset g(\sigma) \quad \text{for some } \sigma \in \Sigma\}.$$

If Σ is a simplicial fan, then each $I \subset g(\sigma)$ is $g(\tau)$ for some $\tau \in \Sigma$, and we obtain the 'underlying complex' of Σ defined in Example 2.2.7. In particular, if Σ is simplicial and complete, then $\mathcal{K}_\Sigma$ is a triangulation of S^{n-1}; and if Σ is a normal fan of a simple polytope, then $\mathcal{K}_\Sigma$ is the boundary complex of the polar simplicial polytope. If Σ is the normal fan of a non-simple polytope P (i.e. the fan over the faces of the polar polytope P^*), then $\mathcal{K}_\Sigma$ is obtained by replacing each face of P^* by a simplex with the same set of vertices; such a simplicial complex is not pure in general.

Proposition 5.4.4. *We have* $U(\Sigma) = U(\mathcal{K}_\Sigma)$.

Proof. We have $(\mathbb{C}, \mathbb{C}^\times)^I \subset (\mathbb{C}, \mathbb{C}^\times)^{g(\sigma)}$ whenever $I \subset g(\sigma)$, hence,

$$U(\Sigma) = \bigcup_{\sigma \in \Sigma} U(\sigma) = \bigcup_{\sigma \in \Sigma} (\mathbb{C}, \mathbb{C}^\times)^{g(\sigma)} = \bigcup_{\sigma \in \Sigma, I \subset g(\sigma)} (\mathbb{C}, \mathbb{C}^\times)^I = \bigcup_{I \in \mathcal{K}_\Sigma} (\mathbb{C}, \mathbb{C}^\times)^I = U(\mathcal{K}_\Sigma).$$

□

We observe that the subset $U(\Sigma) \subset \mathbb{C}^m$ depends only on the combinatorial structure of the fan Σ, while the subgroup $G \subset (\mathbb{C}^\times)^m$ depends on the geometric data, namely, the primitive generators of one-dimensional cones.

Toric variety as a quotient. Since $U(\Sigma) \subset \mathbb{C}^m$ is invariant under the coordinatewise action of $(\mathbb{C}^\times)^m$, we obtain a G-action on $U(\Sigma)$ by restriction.

THEOREM 5.4.5 (Cox [**65**, Theorem 2.1]). *Assume that the linear span of one-dimensional cones of* Σ *is the whole space* $N_\mathbb{R}$.
(a) *The toric variety* V_Σ *is naturally isomorphic to the categorical quotient* $U(\Sigma)/\!\!/G$.
(b) V_Σ *is the geometric quotient* $U(\Sigma)/G$ *if and only if the fan* Σ *is simplicial.*

PROOF. We first prove that the affine variety V_σ corresponding to a cone $\sigma \in \Sigma$ is the categorical quotient $U(\sigma)/\!\!/G$. The algebra of regular functions $\mathbb{C}[U(\sigma)]$ is isomorphic to $\mathbb{C}[z_i, z_j^{-1}\colon 1 \leqslant i \leqslant m,\ j \notin g(\sigma)]$ and is generated by Laurent monomials $\prod_{i=1}^m z_i^{k_i}$ with $k_i \geqslant 0$ for $i \in g(\sigma)$.

It follows easily from (5.3) that a monomial $\prod_{i=1}^m z_i^{k_i}$ is invariant under the G-action on $U(\sigma)$ if and only if it has the form $\prod_{i=1}^m z_i^{\langle \boldsymbol{u}, \boldsymbol{a}_i\rangle}$ for some $\boldsymbol{u} \in N^*$.

Conditions $\langle \boldsymbol{u}, \boldsymbol{a}_i\rangle \geqslant 0$ for $i \in g(\sigma)$ specify the dual cone $\check{\sigma} \subset N^*_\mathbb{R}$, see (2.1). Hence the invariant subalgebra $\mathbb{C}^G[U(\sigma)]$ is isomorphic to $\mathbb{C}[\check{\sigma} \cap N^*] = A_\sigma = \mathbb{C}[V_\sigma]$ (the isomorphism is given by $\prod_{i=1}^m y_i^{\langle \boldsymbol{u}, \boldsymbol{a}_i\rangle} \mapsto \chi^{\boldsymbol{u}}$). Thus, $U(\sigma)/\!\!/G \cong V_\sigma$.

The next step is to glue the isomorphisms $U(\sigma)/\!\!/G \cong V_\sigma$ together into an isomorphism $U(\Sigma)/\!\!/G \cong V_\Sigma$. To do this we need to check that the isomorphisms $\mathbb{C}^G[U(\sigma)] \to \mathbb{C}[V_\sigma]$ are compatible when we pass to the faces of σ. In other words, for each face $\tau \subset \sigma$ we need to establish the commutativity of the diagram

$$\begin{tikzcd}
\mathbb{C}^G[U(\sigma)] \arrow[r] \arrow[d, "\cong"'] & \mathbb{C}^G[U(\tau)] \arrow[d, "\cong"] \\
A_\sigma \arrow[r] & A_\tau .
\end{tikzcd} \tag{5.5}$$

By the definition of a face, we have $\tau = \sigma \cap \boldsymbol{u}^\perp$ for some $\boldsymbol{u} \in \check{\sigma} \cap N^*$, where $\boldsymbol{u}^\perp$ denotes the hyperplane in $N_\mathbb{R}$ normal to $\boldsymbol{u}$. Consider the monomial $\boldsymbol{z}(\boldsymbol{u}) = \prod_{i=1}^m z_i^{\langle \boldsymbol{u}, \boldsymbol{a}_i\rangle}$. Since $\tau = \sigma \cap \boldsymbol{u}^\perp$, the monomial $\boldsymbol{z}(\boldsymbol{u})$ has positive exponent of z_i for $i \in g(\sigma) \setminus g(\tau)$ and zero exponent of z_j for $j \in g(\tau)$. It follows that the algebra $\mathbb{C}[U(\tau)]$ is the localisation of $\mathbb{C}[U(\sigma)]$ by the ideal generated by $\boldsymbol{z}(\boldsymbol{u})$, i.e. $\mathbb{C}[U(\tau)] = \mathbb{C}[U(\sigma)]_{\boldsymbol{z}(\boldsymbol{u})}$. Since $\boldsymbol{z}(\boldsymbol{u})$ is a G-invariant monomial, the localisation commutes with passing to invariant subalgebras, i.e. $\mathbb{C}^G[U(\tau)] = \mathbb{C}^G[U(\sigma)]_{\boldsymbol{z}(\boldsymbol{u})}$. Similarly, $\mathbb{C}[V_\tau] = A_\tau = (A_\sigma)_{\chi^{\boldsymbol{u}}}$. Diagram (5.5) then takes the form

$$\begin{tikzcd}
\mathbb{C}^G[U(\sigma)] \arrow[r] \arrow[d, "\cong"'] & \mathbb{C}^G[U(\sigma)]_{\boldsymbol{z}(\boldsymbol{u})} \arrow[d, "\cong"] \\
\mathbb{C}[V_\sigma] \arrow[r] & \mathbb{C}[V_\sigma]_{\chi^{\boldsymbol{u}}} ,
\end{tikzcd}$$

where the vertical arrows are localisation homomorphism. This diagram is obviously commutative.

Now using the affine cover (5.4) and the compatibility of the isomorphisms on affine varieties we obtain the isomorphism $U(\Sigma)/\!\!/G \cong V_\Sigma = \bigcup_{\sigma\in\Sigma} V_\sigma$. Statement (a) is therefore proved.

To verify (b) we need to check that all orbits of the G-action on $U(\Sigma)$ are closed if and only if the fan Σ is simplicial.

Assume Σ is simplicial, and consider any G-orbit $G\boldsymbol{z}$, $\boldsymbol{z} \in U(\Sigma)$. We shall prove that $G\boldsymbol{z}$ is closed in the usual topology, which is sufficient since the closures of orbits in the usual and Zariski topologies coincide. We need to check that whenever a sequence $\{\boldsymbol{w}^{(k)}\colon k=1,2,\ldots\}$ of points of $G\boldsymbol{z}$ has a limit $\boldsymbol{w}\in U(\Sigma)$, this limit is in $G\boldsymbol{z}$. Write $\boldsymbol{w}^{(k)}=\boldsymbol{g}^{(k)}\boldsymbol{z}$ with $\boldsymbol{g}^{(k)}\in G$. Then it is enough to show that a subsequence of $\{\boldsymbol{g}^{(k)}\}$ converges to a point $\boldsymbol{g}\in G$, as in this case $\lim_{k\to\infty}\boldsymbol{w}^{(k)}=\boldsymbol{g}\boldsymbol{z}\in G\boldsymbol{z}$. We write

$$\boldsymbol{g}^{(k)}=\bigl(g_1^{(k)},\ldots,g_m^{(k)}\bigr)=\bigl(e^{\alpha_1^{(k)}+i\beta_1^{(k)}},\ldots,e^{\alpha_m^{(k)}+i\beta_m^{(k)}}\bigr)\in G\subset(\mathbb{C}^\times)^m$$

where $g_j^{(k)}\in\mathbb{C}^\times$ and $\alpha_j^{(k)},\beta_j^{(k)}\in\mathbb{R}$. Since $e^{i\beta_j^{(k)}}\in\mathbb{S}$ and the circle is compact, we may assume by passing to a subsequence that the sequence $\{e^{i\beta_j^{(k)}}\}$ has a limit $e^{i\beta_j}$ as $k\to\infty$ for each $j=1,\ldots,m$. It remains to consider the sequences $\{e^{\alpha_j^{(k)}}\}$.

By passing to a subsequence we may assume that each sequence $\{\alpha_j^{(k)}\}$, $j=1,\ldots,m$, has a finite or infinite limit (including $\pm\infty$). Let

$$I_+=\{j\colon\alpha_j^{(k)}\to+\infty\}\subset[m],\quad I_-=\{j\colon\alpha_j^{(k)}\to-\infty\}\subset[m].$$

Since the sequence $\{\boldsymbol{w}^{(k)}=\boldsymbol{g}^{(k)}\boldsymbol{z}\}$ is converging to $\boldsymbol{w}=(w_1,\ldots,w_m)\in U(\Sigma)$, we have $z_j=0$ for $j\in I_+$ and $w_j=0$ for $j\in I_-$. Then it follows from the decomposition $U(\Sigma)=\bigcup_{I\in\mathcal{K}_\Sigma}(\mathbb{C},\mathbb{C}^\times)^I$ that I_+ and I_- are simplices of $\mathcal{K}_\Sigma$. Let σ_+,σ_- be the corresponding cones of Σ (here we use the fact that Σ is simplicial). Then $\sigma_+\cap\sigma_-=\{\mathbf{0}\}$ by definition of a fan. By Lemma 2.1.2, there is a linear function $\boldsymbol{u}\in N^*$ such that $\langle\boldsymbol{u},\boldsymbol{a}\rangle>0$ for any nonzero $\boldsymbol{a}\in\sigma_+$, and $\langle\boldsymbol{u},\boldsymbol{a}\rangle<0$ for any nonzero $\boldsymbol{a}\in\sigma_-$. Now, since $\boldsymbol{g}^{(k)}\in G$, it follows from (5.3) that

$$\sum_{j=1}^m\alpha_j^{(k)}\langle\boldsymbol{u},\boldsymbol{a}_j\rangle=0. \tag{5.6}$$

This implies that both I_+ and I_- are empty, as otherwise the sum above tends to infinity. Thus, each sequence $\{\alpha_j^{(k)}\}$ has a finite limit α_j, and a subsequence of $\{\boldsymbol{g}^{(k)}\}$ converges to $(e^{\alpha_1+i\beta_1},\ldots,e^{\alpha_m+i\beta_m})$. Passing to the limit in (5.6) and in the similar equation for $\beta_j^{(k)}$ as $k\to\infty$ we obtain that $(e^{\alpha_1+i\beta_1},\ldots,e^{\alpha_m+i\beta_m})\in G$.

The fact that the G-action on $U(\Sigma)$ with non-simplicial Σ has non-closed orbits is left as an exercise (alternatively, see [**65**, §2] or [**66**, Theorem 5.1.11]. □

The quotient torus $T_N^{\mathrm{a}}=(\mathbb{C}^\times)^m/G$ acts on $V_\Sigma=U(\Sigma)/\!\!/G$ with a dense orbit.

Proposition 5.4.6.

(a) *If Σ is a simplicial fan, then the G-action on $U(\Sigma)$ is almost free (i.e., all stabiliser subgroups are finite);*

(b) *If Σ is regular, then the G-action on $U(\Sigma)$ is free.*

Proof. The stabiliser of a point $\boldsymbol{z}\in\mathbb{C}^m$ under the action of $(\mathbb{C}^\times)^m$ is

$$(\mathbb{C}^\times)^{\omega(\boldsymbol{z})}=\{(t_1,\ldots,t_m)\in(\mathbb{C}^\times)^m\colon t_i=1\text{ if }z_i\neq0\},$$

where $\omega(\boldsymbol{z})$ be the set of zero coordinates of $\boldsymbol{z}$. The stabiliser of $\boldsymbol{z}$ under the G-action is $G_{\boldsymbol{z}}=(\mathbb{C}^\times)^{\omega(\boldsymbol{z})}\cap G$. Since G is the kernel of the map $\ell\colon(\mathbb{C}^\times)^m\to T_N^{\mathrm{a}}$ induced by the map of lattices $\mathbb{Z}^m\to N$, the subgroup $G_{\boldsymbol{z}}$ is the kernel of the composite map

$$(\mathbb{C}^\times)^{\omega(\boldsymbol{z})}\hookrightarrow(\mathbb{C}^\times)^m\xrightarrow{\ell}T_N^{\mathrm{a}}. \tag{5.7}$$

This homomorphism of tori is induced by the map of lattices $\mathbb{Z}^{\omega(\boldsymbol{z})}\to\mathbb{Z}^m\to N$, where $\mathbb{Z}^{\omega(\boldsymbol{z})}\to\mathbb{Z}^m$ is the inclusion of a coordinate sublattice.

Now let Σ be a simplicial fan and $\boldsymbol{z} \in U(\Sigma)$. Then $\omega(\boldsymbol{z}) = g(\sigma)$ for a cone $\sigma \in \Sigma$. Therefore, the set of primitive generators $\{\boldsymbol{a}_i : i \in \omega(\boldsymbol{z})\}$ is linearly independent. Hence, the map $\mathbb{Z}^{\omega(\boldsymbol{z})} \to \mathbb{Z}^m \to N$ taking $\boldsymbol{e}_i$ to $\boldsymbol{a}_i$ is a monomorphism, which implies that the kernel of (5.7) is a finite group.

If the fan Σ is regular, then $\{\boldsymbol{a}_i : i \in \omega(\boldsymbol{z})\}$ is a part of basis of N. In this case (5.7) is a monomorphism and $G_{\boldsymbol{z}} = \{1\}$. □

Remark. Closedness of orbits is a necessary condition for the topological quotient $U(\Sigma)/G$ to be Hausdorff (Exercise B.3.2). The proof of Theorem 5.4.5 (b) above uses Separation Lemma (Lemma 2.1.2); this is another example of situation when convex-geometric separation translates into Hausdorffness.

We may consider the following more general setup. Let $\boldsymbol{a}_1, \ldots, \boldsymbol{a}_m$ be a set of primitive vectors in $N_{\mathbb{R}} \cong \mathbb{R}^n$, and let $\mathcal{K}$ be an $(n-1)$-dimensional simplicial complex on the vertex set $[m]$. Assume further that, for any $I \in \mathcal{K}$, the set of vectors $\{\boldsymbol{a}_i : i \in I\}$ is linearly independent. The latter set spans a simplicial cone, which we denote by σ_I. The data $\{\mathcal{K}, \boldsymbol{a}_1, \ldots, \boldsymbol{a}_m\}$ defines the coordinate subspace arrangement complement $U(\mathcal{K}) \subset \mathbb{C}^m$ and the group G (5.3). Furthermore, the action of G on $U(\mathcal{K})$ is almost free (and it is free if all cones σ_I are regular; this is proved in the same way as Proposition 5.4.6). However, the quotient $U(\mathcal{K})/G$ is Hausdorff precisely when the cones $\{\sigma_I : I \in \mathcal{K}\}$ form a fan Σ. In this case $\mathcal{K} = \mathcal{K}_\Sigma$ and $U(\mathcal{K}) = U(\Sigma)$.

To see that the quotient $U(\Sigma)/G$ is Hausdorff in the case when Σ is a simplicial fan, we do not need to use the algebraic criterion of separatedness as in the proof of Lemma 5.1.4. Instead, we may modify the argument for the closedness of orbits in the proof of Theorem 5.4.5 and show that action of G on $U(\Sigma)$ is *proper* (Exercise 5.4.14), which guarantees that $U(\Sigma)/G$ is Hausdorff.

Example 5.4.7. Let V_σ be the affine toric variety corresponding to an n-dimensional simplicial cone σ. We may write $V_\sigma = V_\Sigma$ where Σ is the simplicial fan consisting of all faces of σ. Then $m = n$, $U(\Sigma) = \mathbb{C}^n$, and $\ell \colon \mathbb{Z}^n \to N$ is the inclusion of a sublattice of finite rank generated by $\boldsymbol{a}_1, \ldots, \boldsymbol{a}_n$. Therefore, G is a finite group and $V_\sigma = \mathbb{C}^n/G = \operatorname{Spec} \mathbb{C}[z_1, \ldots, z_n]^G$.

In particular, if we consider the cone σ generated by $2\boldsymbol{e}_1 - \boldsymbol{e}_2$ and $\boldsymbol{e}_2$ in $\mathbb{R}^2$ (see Example 5.1.6), then G is $\mathbb{Z}_2$ embedded as $\{(1,1), (-1,-1)\}$ in $(\mathbb{C}^\times)^2$. The quotient construction realises the quadratic cone $V_\sigma = \operatorname{Spec} \mathbb{C}[z_1, z_2]^G = \operatorname{Spec} \mathbb{C}[z_1^2, z_1 z_2, z_2^2]$ as a quotient of $\mathbb{C}^2$ by $\mathbb{Z}_2$.

Example 5.4.8. Consider the complete fan of Example 5.1.7. Then

$$U(\Sigma) = \mathbb{C}^3 \setminus \{z_1 = z_2 = z_3 = 0\} = \mathbb{C}^3 \setminus \{0\}$$

The subgroup G defined by (5.2) is the diagonal $\mathbb{C}^\times$ in $(\mathbb{C}^\times)^3$. We therefore obtain $V_\Sigma = U(\Sigma)/G = \mathbb{C}P^2$.

Example 5.4.9. Consider the fan Σ in $\mathbb{R}^2$ with 3 one-dimensional cones generated by the vectors $\boldsymbol{e}_1$, $\boldsymbol{e}_2$ and $-\boldsymbol{e}_1 - \boldsymbol{e}_2$. This fan is not complete, but its 1-dimensional cones generate $\mathbb{R}^2$, so we may apply Theorem 5.4.5. The simplicial complex $\mathcal{K}_\Sigma$ consists of 3 disjoint points. The space $U(\Sigma) = U(\mathcal{K}_\Sigma)$ is therefore the complement to 3 coordinate lines in $\mathbb{C}^3$:

$$U(\Sigma) = \mathbb{C}^3 \setminus \bigl(\{z_1 = z_2 = 0\} \cup \{z_1 = z_3 = 0\} \cup \{z_2 = z_3 = 0\}\bigr)$$

The group G is the diagonal $\mathbb{C}^\times$ in $(\mathbb{C}^\times)^3$. Hence $V_\Sigma = U(\Sigma)/G$ is a quasiprojective variety obtained by removing three points from $\mathbb{C}P^2$.

Exercises.

5.4.10. The one-dimensional cones of a fan $\Sigma \subset N_\mathbb{R}$ span $N_\mathbb{R}$ if and only if the toric variety V_Σ does not have $\mathbb{C}^\times$-factors.

5.4.11. If Σ is a regular fan with at least one n-dimensional cone, then $G \cong (\mathbb{C}^\times)^{m-n}$.

5.4.12. A G-invariant Laurent monomial has the form $\prod_{i=1}^m z_i^{\langle \boldsymbol{u}, \boldsymbol{a}_i\rangle}$ for some $\boldsymbol{u} \in N^*$.

5.4.13. If the fan Σ is non-simplicial, then there exists a non-closed orbit of the G-action on $U(\Sigma)$.

5.4.14. A G-action on X is called *proper* if the map $G \times X \to X \times X$, $(g, x) \mapsto (gx, x)$ is proper, i.e. the preimage of a compact subset is compact. Modify the argument in the proof of Theorem 5.4.5 (b) to show that the G-action on $U(\Sigma)$ is proper whenever Σ is simplicial. (Hint: show that if sequences $\{\boldsymbol{z}^{(k)}\}$ and $\{\boldsymbol{g}^{(k)}\boldsymbol{z}^{(k)}\}$ have limits in $U(\Sigma)$, then a subsequence of $\{\boldsymbol{g}^{(k)}\}$ has a limit in G.)

5.4.15. Let Σ be a regular fan whose one-dimensional cones span $N_\mathbb{R}$. Observe that $U(\Sigma)$ is 2-connected (see Exercise 4.6.11). Show by considering the exact homotopy sequence of the principal G-bundle $U(\Sigma) \to V_\Sigma$ that the nonsingular toric variety V_Σ is simply connected, and $H_2(V_\Sigma; \mathbb{Z})$ is naturally identified with the kernel of the map $\ell\colon \mathbb{Z}^m \to N$. Hence $H^2(V_\Sigma; \mathbb{Z}) = \mathbb{Z}^m/N^*$, which coincides with the *Picard group* of V_Σ, see [**96**, §3.4].

5.5. Hamiltonian actions and symplectic reduction

Here we describe projective toric manifolds as *symplectic quotients* of Hamiltonian torus actions on $\mathbb{C}^m$. This approach may be viewed as a symplectic geometry version of the algebraic quotient construction from the previous section, although historically the symplectic construction preceded the algebraic one [**8**], [**109**].

Symplectic reduction. We briefly review the background material in symplectic geometry, referring the reader to monographs by Audin [**8**] and Guillemin [**109**] for a detailed treatment.

A *symplectic manifold* is a pair (W, ω) consisting of a smooth (but not necessarily closed) manifold W and a closed differential 2-form ω which is nondegenerate at each point. The dimension of a symplectic manifold W is necessarily even.

Assume now that a (compact) torus T acts on W preserving the symplectic form ω. We denote the Lie algebra of the torus T by $\mathfrak{t}$ (since T is commutative, its Lie algebra is trivial, but the construction can be generalised to noncommutative Lie groups). Given an element $\boldsymbol{v} \in \mathfrak{t}$, we denote by $X_{\boldsymbol{v}}$ the corresponding T-invariant vector field on W. The torus action is called *Hamiltonian* if the 1-form $\omega(X_{\boldsymbol{v}}, \cdot\,)$ is exact for any $\boldsymbol{v} \in \mathfrak{t}$. In other words, an action is Hamiltonian if for any $\boldsymbol{v} \in \mathfrak{t}$ there exist a function $H_{\boldsymbol{v}}$ on W (called a *Hamiltonian*) satisfying the condition

$$\omega(X_{\boldsymbol{v}}, Y) = dH_{\boldsymbol{v}}(Y)$$

for any vector field Y on W. The function $H_{\boldsymbol{v}}$ is defined up to constant. Choose a basis $\{\boldsymbol{e}_i\}$ in $\mathfrak{t}$ and the corresponding Hamiltonians $\{H_{\boldsymbol{e}_i}\}$. Then the *moment map*

$$\mu\colon W \to \mathfrak{t}^*, \qquad (x, \boldsymbol{e}_i) \mapsto H_{\boldsymbol{e}_i}(x)$$

(where $x \in W$) is defined. Observe that changing the Hamiltonians $\{H_{\boldsymbol{e}_i}\}$ by constants results in shifting the image of μ by a vector in $\mathfrak{t}^*$. According to a theorem of Atiyah [**5**] and Guillemin–Sternberg [**110**], the image $\mu(W)$ of the moment map is convex, and if W is compact, then $\mu(W)$ is a convex polytope in $\mathfrak{t}^*$.

Example 5.5.1. The most basic example is $W = \mathbb{C}^m$ with symplectic form

$$\omega = i\sum_{k=1}^{m} dz_k \wedge d\overline{z}_k = 2\sum_{k=1}^{m} dx_k \wedge dy_k,$$

where $z_k = x_k + iy_k$. The coordinatewise action of the torus $\mathbb{T}^m$ on $\mathbb{C}^m$ is Hamiltonian. The moment map $\mu\colon \mathbb{C}^m \to \mathbb{R}^m$ is given by $\mu(z_1, \ldots, z_m) = (|z_1|^2, \ldots, |z_m|^2)$ (an exercise). The image of the moment map μ is the positive orthant $\mathbb{R}^m_{\geqslant}$.

Construction 5.5.2 (symplectic reduction). Assume given a Hamiltonian action of a torus T on a symplectic manifold W. Assume further that the moment map $\mu\colon W \to \mathfrak{t}^*$ is *proper*, that is $\mu^{-1}(V)$ is compact for each compact subset $V \subset \mathfrak{t}^*$ (this is always the case if W itself is compact). Let $\boldsymbol{u} \in \mathfrak{t}^*$ be a *regular value* of the moment map, i.e. the differential $\mathcal{T}_x W \to \mathfrak{t}^*$ is surjective for all $x \in \mu^{-1}(\boldsymbol{u})$. Then the level set $\mu^{-1}(\boldsymbol{u})$ is a smooth compact T-invariant submanifold in W. Furthermore, the T-action on $\mu^{-1}(\boldsymbol{u})$ is almost free (an exercise).

Assume now that the T-action on $\mu^{-1}(\boldsymbol{u})$ is free. The restriction of the symplectic form ω to $\mu^{-1}(\boldsymbol{u})$ may be degenerate. However, the quotient manifold $\mu^{-1}(\boldsymbol{u})/T$ is endowed with is a unique symplectic form ω' such that

$$p^*\omega' = i^*\omega,$$

where $i\colon \mu^{-1}(\boldsymbol{u}) \to W$ is the inclusion and $p\colon \mu^{-1}(\boldsymbol{u}) \to \mu^{-1}(\boldsymbol{u})/T$ the projection.

We therefore obtain a new symplectic manifold $(\mu^{-1}(\boldsymbol{u})/T, \omega')$ which is referred to as the *symplectic reduction*, or *symplectic quotient* of (W, ω) by T.

The construction of symplectic reduction works also under milder assumptions on the action (see [**79**] and more references there), but the generality described here will be enough for our purposes.

The toric case. The algebraic quotient construction describes a toric manifold V_Σ as a quotient of a noncompact set $U(\Sigma)$ by a noncompact group G. Using symplectic reduction, a projective toric manifold V_P corresponding to a simple lattice polytope P can be obtained as the quotient of a compact submanifold $\mathcal{Z}_P \subset U(\Sigma_P)$ by a free action of a compact torus.

Let Σ be a complete regular fan in $N_\mathbb{R} \cong \mathbb{R}^n$ with m one-dimensional cones generated by $\boldsymbol{a}_1, \ldots, \boldsymbol{a}_m$. Consider the exact sequence of maximal compact subgroups (tori) corresponding to exact sequence of algebraic tori (5.2):

$$1 \longrightarrow K \longrightarrow \mathbb{T}^m \xrightarrow{\ell} T_N \longrightarrow 1, \tag{5.8}$$

where $T_N = N \otimes_\mathbb{Z} \mathbb{S} \cong \mathbb{T}^n$, $\ell\colon \mathbb{T}^m \to T_N$ is the map of tori corresponding to the map of lattices $\mathbb{Z}^m \to N$, $\boldsymbol{e}_i \mapsto \boldsymbol{a}_i$, and $K = \operatorname{Ker} \ell$. The group K is isomorphic to $\mathbb{T}^{m-n}$ because Σ is complete and regular.

We let $K \subset \mathbb{T}^m$ act on $\mathbb{C}^m$ by restriction of the coordinatewise action of $\mathbb{T}^m$. This K-action on $\mathbb{C}^m$ is also Hamiltonian, and the corresponding moment map is given by the composition

$$\mu_\Sigma\colon \mathbb{C}^m \longrightarrow \mathbb{R}^m \longrightarrow \mathfrak{k}^*, \tag{5.9}$$

where $\mathbb{R}^m \to \mathfrak{k}^*$ is the map of dual Lie algebras corresponding to the inclusion $K \to \mathbb{T}^m$. Choose a basis in $\mathfrak{k}^* \cong \mathbb{R}^{m-n}$, so that the linear map $\mathbb{R}^m \to \mathfrak{k}^*$ is defined by an $(m-n) \times m$-matrix $C = (c_{jk})$. Moment map (5.9) is then given by

$$(z_1, \ldots, z_m) \longmapsto \Bigl(\sum_{k=1}^m c_{1k}|z_k|^2, \ldots, \sum_{k=1}^m c_{m-n,k}|z_k|^2\Bigr),$$

and its level set $\mu_\Sigma^{-1}(\boldsymbol{d})$ corresponding to a value $\boldsymbol{d} = (d_1, \ldots, d_{m-n}) \in \mathfrak{k}^*$ is the intersection of $m-n$ Hermitian quadrics in $\mathbb{C}^m$:

$$\sum_{k=1}^m c_{jk}|z_k|^2 = d_j \qquad \text{for } j = 1, \ldots, m-n. \tag{5.10}$$

To apply the symplectic reduction we need to identify the regular values of the moment map μ_Σ. We recall from Section 2.1 that a polytope (1.1) is called Delzant if its normal fan is regular. If $\Sigma = \Sigma_P$ is the normal fan of a Delzant polytope P, then the $(m-n) \times m$-matrix C above is the one considered in Construction 1.1.5. Namely, the rows of C form a basis of linear dependencies between the vectors $\boldsymbol{a}_i$.

Given a polytope (1.1), we denote by $\mathcal{Z}_P$ the intersection of quadrics (5.10) corresponding to $\boldsymbol{d} = C\boldsymbol{b}_P$, where $\boldsymbol{b}_P = (b_1, \ldots, b_m)^t$:

$$\mathcal{Z}_P = \mu_\Sigma^{-1}(C\boldsymbol{b}_P) = \Bigl\{\boldsymbol{z} \in \mathbb{C}^m \colon \sum_{k=1}^m c_{jk}\bigl(|z_k|^2 - b_k\bigr) = 0 \ \text{ for } j = 1, \ldots, m-n\Bigr\}. \tag{5.11}$$

Proposition 5.5.3. *Assume that $\Sigma = \Sigma_P$ is the normal fan of a Delzant polytope P given by* (1.1). *Then $\boldsymbol{d} = C\boldsymbol{b}_P$ is a regular value of the moment map μ_Σ.*

Proof. We only sketch the proof here, as the statement will be proved in a more general context in Section 6.1. We need to check that this intersection is nondegenerate at each point $\boldsymbol{z} \in \mathcal{Z}_P$, i.e. that $\mathcal{Z}_P$ is a smooth submanifold in $\mathbb{C}^m$. This means that the $m-n$ gradient vectors of the left hand sides of quadratic equations (5.10) are linearly independent at each $\boldsymbol{z} \in \mathcal{Z}_P$. This can be shown to be equivalent to that the polytope P defined by (1.1) is simple. □

As we shall see in Section 6.2, the manifold $\mathcal{Z}_P$ is $\mathbb{T}^m$-equivariantly homeomorphic to the moment-angle manifold $\mathcal{Z}_{\mathcal{K}_P}$ (where $\mathcal{K}_P$ is the nerve complex of P, or the underlying simplicial complex of the normal fan Σ_P). Furthermore, we have $\mathcal{Z}_P \subset U(\Sigma_P)$. (The reader may either wait until Chapter 6, or view these two statements as exercises.)

We therefore may consider the symplectic quotient of $\mathbb{C}^m$ by $K \cong \mathbb{T}^{m-n}$, the $2n$-dimensional symplectic manifold $\mu_\Sigma^{-1}(C\boldsymbol{b}_P)/K = \mathcal{Z}_P/K$.

Theorem 5.5.4. *Let P be a lattice Delzant polytope with the normal fan Σ_P, and let V_P be the corresponding projective toric manifold. The inclusion $\mathcal{Z}_P \subset U(\Sigma_P)$ induces a diffeomorphism*

$$\mathcal{Z}_P/K \xrightarrow{\cong} U(\Sigma_P)/G = V_P.$$

Therefore, any projective toric manifold V_P is obtained as the symplectic quotient of $\mathbb{C}^m$ by an action of a torus $K \cong \mathbb{T}^{m-n}$.

Proof. We sketch the proof given in [**8**, Prop. VI.3.1.1]; a different proof of a more general statement will be given in Section 6.5.

Consider the function

$$f\colon \mathbb{C}^m \to \mathbb{R}, \quad f(\boldsymbol{z}) = \|\mu_\Sigma(\boldsymbol{z}) - C\boldsymbol{b}_P\|^2.$$

It is nonnegative and minimal on the set $\mathcal{Z}_P = \mu_\Sigma^{-1}(C\boldsymbol{b}_P)$. The only critical points of f in $U(\Sigma_P)$ are $\boldsymbol{z} \in \mathcal{Z}_P$. Hence, for any $\boldsymbol{z} \in U(\Sigma_P)$, the gradient trajectory descending from $\boldsymbol{z}$ will reach a point in $\mathcal{Z}_P$. Furthermore, any gradient trajectory is contained in a G-orbit. We therefore obtain that each G-orbit of $U(\Sigma_P)$ intersects $\mathcal{Z}_P$. Finally, it can be shown that each G-orbit intersects $\mathcal{Z}_P$ at a unique K-orbit, i.e. for each $\boldsymbol{z} \in \mathcal{Z}_P$ we have that

$$G \cdot \boldsymbol{z} \cap \mathcal{Z}_P = K \cdot \boldsymbol{z}.$$

The statement follows. □

The toric manifold V_P therefore acquires a symplectic structure as the symplectic quotient $\mu_\Sigma^{-1}(C\boldsymbol{b}_P)/K$. On the other hand, the projective embedding of V_P defined by the lattice polytope P provides a symplectic form on V_P by restriction of the standard symplectic form on the complex projective space. It can be shown [**109**, Appendix 2] that the diffeomorphism from Theorem 5.5.4 preserves the cohomology class of the symplectic form, or equivalently, the two symplectic structures on V_P are T_N-equivariantly symplectomorphic.

The symplectic quotient $\mu_\Sigma^{-1}(C\boldsymbol{b}_P)/K$ has a residual action of the quotient n-torus $T_N = \mathbb{T}^m/K$, which is obviously Hamiltonian. This action is identified, via Theorem 5.5.4, with the action of the maximal compact subgroup $T_N \subset T_N^{\mathrm{a}}$ on the toric variety V_P. We denote by $\mu_V\colon V_P \to \mathfrak{t}^*$ the moment map for the Hamiltonian action of T_N on V_P, where $\mathfrak{t} \cong \mathbb{R}^n$ is the Lie algebra of T_N. We obtain from (5.8) that the map ℓ^* embeds $\mathfrak{t}^*$ as a subspace in $\mathbb{R}^m$, and observe that ℓ^* is the map A_P from Construction 1.1.5.

Proposition 5.5.5. *The image of the moment map $\mu_V\colon V_P \to \mathfrak{t}^*$ is the polytope P, up to translation.*

Proof. Let ω be the standard symplectic form on $\mathbb{C}^m$ and $\mu\colon \mathbb{C}^m \to \mathbb{R}^m$ be the moment map for the standard action of $\mathbb{T}^m$ (see Example 5.5.1). Let $p\colon \mathcal{Z}_P \to V_P$ be the quotient projection by the action of K, and let $i\colon \mathcal{Z}_P \to \mathbb{C}^m$ be the inclusion, so that the symplectic form ω' on V_P satisfies $p^*\omega' = i^*\omega$. Let $H_{\boldsymbol{e}_i}\colon \mathbb{C}^m \to \mathbb{R}$ be the Hamiltonian of the $\mathbb{T}^m$-action on $\mathbb{C}^m$ corresponding to the ith basis vector $\boldsymbol{e}_i$ (explicitly, $H_{\boldsymbol{e}_i}(\boldsymbol{z}) = |z_i|^2$), and let $H_{\boldsymbol{a}_i}\colon V_P \to \mathbb{R}$ be the Hamiltonian of the T_N-action on V_P corresponding to $\boldsymbol{a}_i \in \mathfrak{t}$. Denote by $X_{\boldsymbol{e}_i}$ the vector field on $\mathcal{Z}_P$ generated by $\boldsymbol{e}_i$, and denote by $Y_{\boldsymbol{a}_i}$ the vector field on V_P generated by $\boldsymbol{a}_i$. Observe that $p_*X_{\boldsymbol{e}_i} = Y_{\boldsymbol{a}_i}$. For any vector field Z on $\mathcal{Z}_P$ we have

$$\begin{aligned} dH_{\boldsymbol{e}_i}(Z) = i^*\omega(X_{\boldsymbol{e}_i}, Z) &= p^*\omega'(X_{\boldsymbol{e}_i}, Z) \\ &= \omega'(Y_{\boldsymbol{a}_i}, p_*Z) = dH_{\boldsymbol{a}_i}(p_*Z) = d(p^*H_{\boldsymbol{a}_i})(Z), \end{aligned}$$

hence $H_{\boldsymbol{e}_i} = p^*H_{\boldsymbol{a}_i}$ or $H_{\boldsymbol{e}_i}(\boldsymbol{z}) = H_{\boldsymbol{a}_i}(p(\boldsymbol{z}))$ up to constant. By definition of the moment map this implies that $\mu_V(V_P)$ is identified with $\mu(\mathcal{Z}_P) \subset \mathbb{R}^m$ up to shift by a vector in $\mathfrak{t}^* \subset \mathbb{R}^m$. The subset $\mu(\mathcal{Z}_P) \subset \mathbb{R}^m$ consists of $\boldsymbol{y} \in \mathbb{R}^m$ with nonnegative coordinates satisfying $C\boldsymbol{y} = C\boldsymbol{b}_P$ (see (5.11)). According to (1.7), the latter specifies the image of P under the affine map i_P whose linear part is exactly $A_P\colon \mathfrak{t}^* \to \mathbb{R}^m$. □

Example 5.5.6. Let $P = \Delta^n$ be the standard simplex (see Example 1.1.6). The cones of the normal fan Σ_P are spanned by the proper subsets of the set of $n+1$ vectors $\{\boldsymbol{e}_1, \ldots, \boldsymbol{e}_n, -\boldsymbol{e}_1 - \cdots - \boldsymbol{e}_n\}$. The groups $G \cong \mathbb{C}^\times$ and $K \cong \mathbb{S}^1$ are the diagonal subgroups in $(\mathbb{C}^\times)^{n+1}$ and $\mathbb{T}^{n+1}$ respectively, and $U(\Sigma_P) = \mathbb{C}^{n+1} \setminus \{0\}$. The matrix C is a row of $n+1$ units. The moment map (5.9) is given by $\mu_\Sigma(z_1, \cdots, z_{n+1}) = |z_1|^2 + \ldots + |z_{n+1}|^2$. Since $Cb_P = 1$, the manifold $\mathcal{Z}_P = \mu_P^{-1}(1)$ is the unit sphere $\mathbb{S}^{2n+1} \subset \mathbb{C}^{n+1}$, and $\mathbb{S}^{2n+1}/K \cong (\mathbb{C}^{n+1} \setminus \{0\})/G = V_P$ is the complex projective space $\mathbb{C}P^n$.

Exercises.

5.5.7. The moment map $\mu\colon \mathbb{C}^m \to \mathbb{R}^m$ for the coordinatewise action of $\mathbb{T}^m$ on $\mathbb{C}^m$ is given by $\mu(z_1, \ldots, z_m) = (|z_1|^2, \ldots, |z_m|^2)$.

5.5.8. The T-action on $\mu^{-1}(\boldsymbol{u})$ is almost free.

5.5.9. The level set $\mathcal{Z}_P = \mu_\Sigma^{-1}(C\boldsymbol{b}_P)$ is $\mathbb{T}^m$-equivariantly homeomorphic to the moment-angle manifold $\mathcal{Z}_{\mathcal{K}_P}$.

5.5.10. Show that $\mathcal{Z}_P \subset U(\Sigma_P)$.

5.6. Quotient space by the torus action

Here we take a closer look at the action of the compact torus T_N on a toric manifold V_Σ. The topological properties of the quotient projection $\pi\colon V_\Sigma \to V_\Sigma/T_N$ will be taken as the starting point for subsequent topological generalisations of toric manifolds in Chapter 7.

We first observe that if V_P is a projective toric manifold, then $V_P = \mathcal{Z}_P/K$ by the construction of the previous section, therefore the quotient of V_P by the action of $T_N = \mathbb{T}^m/K$ coincides with the quotient of $\mathcal{Z}_P$ by $\mathbb{T}^m$. Both quotients are identified with the simple Delzant polytope P; in fact, the moment map $\mu_V\colon V_P \to P$ is the quotient projection (see Proposition 5.5.5). In the non-projective smooth case, there is no moment map, and there is no canonical way to identify the quotient V_Σ/T_N with a convex polytope. However, there is a face decomposition (stratification) of V_Σ/T_N according to orbit types, and this face structure is very similar to that of a simple polytope.

As usual, for each $I \subset [m]$ we denote by $\mathbb{T}^I = \prod_{i\in I} \mathbb{T}$ the corresponding coordinate subgroup in $\mathbb{T}^m$.

We consider the map $\mathbb{R}_\geqslant \times \mathbb{T} \to \mathbb{C}$ defined by $(y,t) \mapsto yt$. Taking product we obtain a map $\mathbb{R}^m_\geqslant \times \mathbb{T}^m \to \mathbb{C}^m$. The preimage of a point $\boldsymbol{z} \in \mathbb{C}^m$ under this map is $\boldsymbol{y} \times \mathbb{T}^{\omega(\boldsymbol{z})}$, where $y_i = |z_i|$ for $1 \leqslant i \leqslant m$ and $\omega(\boldsymbol{z}) = \{i\colon z_i = 0\} \subset [m]$ is the set of zero coordinates of $\boldsymbol{z}$. Therefore, $\mathbb{C}^m$ can be identified with the quotient space

$$\mathbb{R}^m_\geqslant \times \mathbb{T}^m/{\sim} \quad \text{where } (\boldsymbol{y}, \boldsymbol{t}_1) \sim (\boldsymbol{y}, \boldsymbol{t}_2) \text{ if } \boldsymbol{t}_1^{-1}\boldsymbol{t}_2 \in \mathbb{T}^{\omega(\boldsymbol{y})}. \tag{5.12}$$

Now let P be a simple Delzant polytope. Following Davis–Januszkiewicz [**70**], with may describe $\mathcal{Z}_P$ and V_P as similar quotient spaces.

Given $\boldsymbol{x} \in P$, set $I_{\boldsymbol{x}} = \{i \in [m]\colon \boldsymbol{x} \in F_i\}$ (the set of facets containing $\boldsymbol{x}$).

Proposition 5.6.1. *$\mathcal{Z}_P$ is $\mathbb{T}^m$-equivariantly homeomorphic to the quotient*

$$P \times \mathbb{T}^m/\!\sim \quad \textit{where } (\boldsymbol{x}, \boldsymbol{t}_1) \sim (\boldsymbol{x}, \boldsymbol{t}_2) \ \textit{ if } \ \boldsymbol{t}_1^{-1}\boldsymbol{t}_2 \in \mathbb{T}^{I_{\boldsymbol{x}}}.$$

Proof. It follows from (1.7) and (5.11) that $\mathcal{Z}_P/\mathbb{T}^m = \mu(\mathcal{Z}_P) = i_P(P) \subset \mathbb{R}^m_\geqslant$. Since $\mathbb{C}^m = \mathbb{R}^m_\geqslant \times \mathbb{T}^m/\!\sim$, we have $\mathcal{Z}_P = i_P(P) \times \mathbb{T}^m/\!\sim$ by restriction, where $\sim$ is the equivalence relation from (5.12). It remains to note that a point $\boldsymbol{x} \in P$ is mapped by i_P to $\boldsymbol{y} \in \mathbb{R}^m_\geqslant$ with $I_{\boldsymbol{x}} = \omega(\boldsymbol{y})$. ☐

To obtain a quotient presentation of V_P, we consider the map $\ell\colon \mathbb{T}^m \to T_N$ from (5.8), and for each $\boldsymbol{x} \in P$ define the subtorus $T_{\boldsymbol{x}} = \ell(\mathbb{T}^{I_{\boldsymbol{x}}}) \subset T_N$. If $\boldsymbol{x}$ is a vertex then $T_{\boldsymbol{x}} = T_N$, and if $\boldsymbol{x}$ is an interior point of P then $T_{\boldsymbol{x}} = \{1\}$.

Proposition 5.6.2. *A projective toric manifold V_P is T_N-equivariantly homeomorphic to the quotient*

$$P \times T_N/\!\approx \quad \textit{where } (\boldsymbol{x}, t_1) \approx (\boldsymbol{x}, t_2) \ \textit{ if } \ t_1^{-1}t_2 \in T_{\boldsymbol{x}}. \tag{5.13}$$

Proof. Using Proposition 5.6.1 we obtain

$$V_P = \mathcal{Z}_P/K = \bigl(P \times (\mathbb{T}^m/K)\bigr)/\!\sim\; = \bigl(P \times \ell(\mathbb{T}^m)\bigr)/\!\sim\; = P \times T_N/\!\approx. \qquad \square$$

The projection $\pi\colon V_P = P \times T_N/\!\approx\, \to P$ is the quotient map for the T_N-action, and its fibre $\pi^{-1}(\boldsymbol{x}) = T_{\boldsymbol{x}}$ is the stabiliser of the T_N-orbit corresponding to $\boldsymbol{x}$. The T_N-action on V_P is therefore free over the interior of the polytope, the vertices correspond to the fixed points, and points in the relative interior of a codimension-k face correspond to orbits with the same k-dimensional stabiliser.

Construction 5.6.3. Let Σ be a simplicial fan and V_Σ the corresponding toric variety. Consider the affine cover $\{V_\sigma\colon \sigma \in \Sigma\}$ (see Construction 5.1.3). The orbit space $(V_\sigma)_\geqslant = V_\sigma/T_N$ by the action of the compact torus can be identified with the set of semigroup homomorphisms from S_σ to the semigroup $\mathbb{R}_\geqslant$ of nonnegative real numbers:

$$V_\sigma/T_N = \mathrm{Hom}_{\mathrm{sg}}(S_\sigma, \mathbb{R}_\geqslant), \quad \sigma \in \Sigma$$

(the details of this construction can be found in [**96**, §4.1]).

If the fan Σ is regular, then $V_\sigma \cong \mathbb{C}^k \times (\mathbb{C}^\times)^{n-k}$ where $k = \dim \sigma$, see Example 5.1.5. It follows easily that the T_N-action on the nonsingular toric variety V_Σ is locally standard, and the cover $\{V_\sigma\colon \sigma \in \Sigma\}$ provides an atlas of standard charts (see Section D.1). Furthermore, $(V_\sigma)_\geqslant \cong \mathbb{R}^k_\geqslant \times \mathbb{R}^{n-k}$ and the orbit space $Q = V_\Sigma/T_N$ is a manifold with corners, with atlas $\{(V_\sigma)_\geqslant\colon \sigma \in \Sigma\}$.

In the singular case the varieties V_σ may be not isomorphic to $\mathbb{C}^k \times (\mathbb{C}^\times)^{n-k}$ and the T_N-action on V_Σ may fail to be locally standard. Nevertheless, the cover $\{(V_\sigma)_\geqslant\colon \sigma \in \Sigma\}$ defines a structure of a manifold with faces on $Q = V_\Sigma/T_N$.

If Σ is a complete simplicial fan, then the face decomposition of the manifold with corners $Q = V_\Sigma/T_N$ is Poincaré dual to the sphere triangulation defined by the simplicial complex $\mathcal{K}_\Sigma$. There is also a projection $Q \times T_N \to V_\Sigma$ defining a homeomorphism $V_\Sigma \cong Q \times T_N/\!\approx$, by analogy with (5.13).

We finish by summarising the observations of this section as follows:

Proposition 5.6.4.

(a) *The action of the compact torus T_N on a nonsingular toric variety V is locally standard.*

(b) *If* V_P *is a projective toric manifold then the quotient* V_P/T_N *is diffeomorphic to* P *as a manifold with corners.*

If a toric manifold V_Σ is not projective, then the manifold with corners $Q = V_\Sigma/T_N$ may be not homeomorphic to any simple polytope (although no such examples are known, see Problem 5.2.3).

CHAPTER 6

Geometric structures on moment-angle manifolds

6.1. Intersections of quadrics

6.2. Moment-angle manifolds from polytopes

6.3. LVM-manifolds

6.4. Moment-angle manifolds from complete simplicial fans

6.5. Complex geometry of moment-angle manifolds

6.6. Dolbeault cohomology and Hodge numbers

6.7. Hamiltonian-minimal Lagrangian submanifolds

6.8. Moment-angle complexes as Kempf–Ness sets

CHAPTER 7

Half-dimensional torus actions

7.1. Basic cohomological theory

7.2. Torus manifolds

7.2.1. Characteristic submanifolds.

7.2.2. Orbit quotients and manifolds with corners.

7.2.3. Cohomology ring calculations.

7.2.4. Torus manifolds over homology polytopes.

7.2.5. Torus manifolds over face-acyclic manifolds with corners.

7.3. Quasitoric manifolds

7.3.1. Definition and main constructions.

7.3.2. Omniorientations and combinatorial quasitoric data.

7.3.3. Canonical smooth and stably complex structures.

7.3.4. Weights and signs at fixed points.

7.3.5. Cohomology ring and characteristic classes.

7.3.6. Equivariant almost complex structures.

7.3.7. Relationship with toric manifolds.

7.4. Topological toric manifolds

7.5. Bounded flag manifolds

7.6. Bott towers

7.7. Weight graphs

CHAPTER 8

Homotopy theory of moment-angle complexes

8.1. Colimit and homotopy colimit decompositions

8.2. Rational homotopy theory

8.3. Explicit homotopy types

8.4. Stable decompositions of polyhedral products

8.5. Loop spaces

CHAPTER 9

Complex cobordism in toric topology

9.1. Hirzebruch genera of combinatorial quasitoric data

9.2. Toric generators of cobordism ring

9.3. Quasitoric representatives in complex cobordism classes

9.4. Toric genera

APPENDIX A

Commutative and homological algebra

Here we review some basic algebraic notions and results in a way suited for topological applications. In order to make algebraic constructions compatible with topological ones we sometimes use a notation which may seem unusual to a reader with an algebraic background. This in particular concerns the way we treat gradings and resolutions.

We fix a ground ring $\mathbf{k}$, which is always assumed to be a field or the ring $\mathbb{Z}$ of integers. In the latter case by a '$\mathbf{k}$-vector space of dimension d' we mean an abelian group of rank d.

A.1. Algebras and modules

A $\mathbf{k}$-*algebra* (or simply *algebra*) A is a ring which is also a $\mathbf{k}$-vector space, and whose multiplication $A\times A\to A$ is $\mathbf{k}$-bilinear. (The latter condition is void if $\mathbf{k}=\mathbb{Z}$, so $\mathbb{Z}$-algebras are ordinary rings.) All our algebras will be commutative and with unit 1, unless explicitly stated otherwise. The basic example is $A=\mathbf{k}[v_1,\dots,v_m]$, the *polynomial algebra* in m generators, for which we shall often use a shortened notation $\mathbf{k}[m]$.

An algebra A is *finitely generated* if there are finitely many elements $a_1,\dots,a_n$ of A such that every element of A can be written as a polynomial in $a_1,\dots,a_n$ with coefficients in $\mathbf{k}$. Therefore, a finitely generated algebra is the quotient of a polynomial algebra by an ideal.

An A-*module* is a $\mathbf{k}$-vector space M on which A acts linearly, that is, there is a map $A\times M\to M$ which is $\mathbf{k}$-linear in each argument and satisfies $1m=m$ and $(ab)m=a(bm)$ for all $a,b\in A$, $m\in M$. Any ideal I of A is an A-module. If $A=\mathbf{k}$, then an A-module is a $\mathbf{k}$-vector space.

An A-module M is *finitely generated* if there exist $x_1,\dots,x_n$ in M such that every element x of M can be written (not necessarily uniquely) as $x=a_1x_1+\dots+a_nx_n,\ \ a_i\in A$.

An algebra A is $\mathbb{Z}$-*graded* (or simply *graded*) if it is represented as a direct sum $A=\bigoplus_{i\in\mathbb{Z}}A^i$ such that $A^i\cdot A^j\subset A^{i+j}$. Elements $a\in A^i$ are said to be *homogeneous* of degree i, denoted $\deg a=i$. The set of homogeneous elements of A is denoted by $\mathcal{H}(A)=\bigcup_i A^i$. An ideal I of A is *homogeneous* if it is generated by homogeneous elements. In most cases our graded algebras will be either *nonpositively graded* (i.e. $A^i=0$ for $i>0$) or *nonnegatively graded* (i.e. $A^i=0$ for $i<0$); the latter is also called an $\mathbb{N}$-graded algebra. A nonnegatively graded algebra A is *connected* if $A^0=\mathbf{k}$. For a nonnegatively graded algebra A, define the *positive ideal* by $A^+=\bigoplus_{i>0}A^i$; if A is connected and $\mathbf{k}$ is a field then A^+ is a maximal ideal.

If A is a graded algebra, then an A-module M is *graded* if $M=\bigoplus_{i\in\mathbb{Z}}M^i$ such that $A^i\cdot M^j\subset M^{i+j}$. An A-module map $f\colon M\to N$ between two graded modules is *degree-preserving* (or of *degree* 0) if $f(M^i) \subset N^i$, and is of *degree* k if $f(M^i) \subset N^{i+k}$ for all i.

Graded algebras arising in topology are often *graded-commutative* (or *skew-commutative*) rather than commutative in the usual sense. This means that

$$ab = (-1)^{ij}\, ba \quad \text{for any } a \in A^i, b \in A^j.$$

If the characteristic of $\mathbf{k}$ is not 2, then the square of an odd-degree element in a graded-commutative algebra is zero. To avoid confusion we double the grading in commutative algebras A; the resulting graded algebras $A = \bigoplus_{i\in\mathbb{Z}} A^{2i}$ are commutative in either sense.

For example, we make the polynomial algebra $\mathbf{k}[v_1, \ldots, v_m]$ graded by setting $\deg v_i = 2$. It then becomes a *free* graded commutative algebra on m generators of degree two (free means no relations apart from the graded commutativity). The *exterior algebra* $\Lambda[u_1, \ldots, u_m]$ has relations $u_i^2 = 0$ and $u_i u_j = -u_j u_i$. We shall assume $\deg u_i = 1$ unless otherwise specified. An exterior algebra is a free graded commutative algebra if the characteristic of $\mathbf{k}$ is not 2.

Bigraded (i.e. $\mathbb{Z}\oplus\mathbb{Z}$-graded) and *multigraded* ($\mathbb{Z}^m$-graded) algebras A are defined similarly; their homogeneous elements $a \in A$ have *bidegree* $\operatorname{bideg} a = (i, j) \in \mathbb{Z}\oplus\mathbb{Z}$ or *multidegree* $\operatorname{mdeg} a = \boldsymbol{i} \in \mathbb{Z}^m$ respectively.

The *tensor product* $M \otimes_A N$ of A-modules M and N is the quotient of a free A-module on the set of generators $M \times N$ by the submodule generated by all elements of the following types:

$$(x + x', y) - (x, y) - (x', y), \quad (x, y + y') - (x, y) - (x, y'),$$
$$(ax, y) - a(x, y), \quad (x, ay) - a(x, y),$$

where $x, x' \in M$, $y, y' \in N$, $a \in A$. For each basis element (x, y), its image in $M \otimes_A N$ is denoted by $x \otimes y$.

We shall denote the tensor product $M \otimes_{\mathbf{k}} N$ of $\mathbf{k}$-vector spaces by simply $M \otimes N$. For example, if $M = N = \mathbf{k}[v]$, then $M \otimes N = \mathbf{k}[v_1, v_2]$.

The tensor product $A \otimes B$ of graded-commutative algebras A and B is a graded commutative algebra, with the multiplication defined on homogeneous elements by

$$(a \otimes b) \cdot (a' \otimes b') = (-1)^{\deg b \deg a'} aa' \otimes bb'.$$

An A-module F is *free* if it is isomorphic to a direct sum $\bigoplus_{i\in I} F_i$, where each F_i is isomorphic to A as an A-module. If both A and F are graded then every F_i is isomorphic to $A^{[j]}$ for some j, where $A^{[j]}$ is the graded A-module with $(A^{[j]})^k = A^{k-j}$. A *basis* of a free A-module F is a set $\mathcal{S}$ of elements of F such that each $x \in F$ can be uniquely written as a finite linear combination of elements of $\mathcal{S}$ with coefficients in A. If A is finitely generated then all bases have the same cardinality (an exercise), called the *rank* of F. If $\mathcal{S}$ is a basis of a free A-module F, then for any A-module M a set map $\mathcal{S} \to M$ extends to an A-module homomorphism $F \to M$.

A module P is *projective* if for any epimorphism of modules $p\colon M \to N$ and homomorphism $f\colon P \to N$, there is a homomorphism $f'\colon P \to M$ such that $pf' = f$. This is described by the following commutative diagram:

$$\begin{tikzcd}
M \arrow[r, "p"] & N \arrow[r] & 0 \\
& P \arrow[u, "f"'] \arrow[ul, dashed, "f'"] &
\end{tikzcd}$$

Equivalently P is projective if it is a direct summand of a free module (an exercise). In particular, free modules are projective.

A sequence of homomorphisms of A-modules

$$\cdots \longrightarrow M_1 \xrightarrow{f_1} M_2 \xrightarrow{f_2} M_3 \xrightarrow{f_3} M_4 \longrightarrow \cdots$$

is called an *exact sequence* if $\operatorname{Im} f_i = \operatorname{Ker} f_{i+1}$ for all i.

A *chain complex* is a sequence $C_* = \{C_i, \partial_i\}$ of A-modules C_i and homomorphisms $\partial_i\colon C_i \to C_{i-1}$ such that $\partial_i \partial_{i+1} = 0$. This is usually written as

$$\cdots \longrightarrow C_{i+1} \xrightarrow{\partial_{i+1}} C_i \xrightarrow{\partial_i} C_{i-1} \longrightarrow \cdots$$

The condition $\partial_i \partial_{i+1} = 0$ implies that $\operatorname{Im} \partial_{i+1} \subset \operatorname{Ker} \partial_i$. The elements of $\operatorname{Ker} \partial$ are called *cycles*, and the elements of $\operatorname{Im} \partial$ are *boundaries*. The *ith homology group* (or *homology module*) of C_* is defined by

$$H_i[C_*] = \operatorname{Ker} \partial_i / \operatorname{Im} \partial_{i+1}.$$

A *cochain complex* is a sequence $C^* = \{C^i, d^i\}$ of A-modules C^i and homomorphisms $d^i\colon C^i \to C^{i+1}$ such that $d^i d^{i-1} = 0$. This is usually written as

$$\cdots \longrightarrow C^{i-1} \xrightarrow{d^{i-1}} C^i \xrightarrow{d^i} C^{i+1} \longrightarrow \cdots .$$

The elements of $\operatorname{Ker} d$ are called *cocycles*, and the elements of $\operatorname{Im} d$ are *coboundaries*. The *ith cohomology group* (or *cohomology module*) of C^* is defined by

$$H^i[C^*] = \operatorname{Ker} d^i / \operatorname{Im} d^{i-1}.$$

A cochain complex may be also viewed as a graded **k**-vector space $C^* = \bigoplus_i C^i$ in which every graded component C^i is an A-module, together with an A-linear map $d\colon C^* \to C^*$ raising the degree by 1 and satisfying the condition $d^2 = 0$.

Note that a chain complex may be turned to a cochain complex by inverting the grading (i.e. turning the ith graded component into the $(-i)$th).

Let $f, g\colon C^* \to D^*$ be two maps of cochain complexes (i.e. both f and g commute with the differentials). A *cochain homotopy* between f and g is a set of maps $s = \{s^i\colon C^i \to D^{i-1}\}$ satisfying the identities

$$ds + sd = f - g$$

(more precisely, $d^{i-1}s^i + s^{i+1}d^i = f^i - g^i$). This is described by the following commutative diagram

$$\begin{tikzcd}
\cdots \arrow[r] & C^{i-1} \arrow[r, "d^{i-1}"] \arrow[d] \arrow[dl, "s^{i-1}"'] & C^i \arrow[r, "d^i"] \arrow[d, "f^i-g^i"] \arrow[dl, "s^i"'] & C^{i+1} \arrow[r] \arrow[d] \arrow[dl, "s^{i+1}"'] & \cdots \\
\cdots \arrow[r] & D^{i-1} \arrow[r, "d^{i-1}"'] & D^i \arrow[r, "d^i"'] & D^{i+1} \arrow[r] & \cdots
\end{tikzcd}$$

If there is a cochain homotopy between f and g, then f and g induce the same map in cohomology (an exercise). A *chain homotopy* between maps of chain complexes is defined similarly.

A *differential graded algebra* is a graded algebra A together with a **k**-linear map $d\colon A \to A$, called the *differential*, which raises the degree by one, and satisfies the identity $d^2 = 0$ (so that $\{A^i, d^i\}$ is a cochain complex) and the *Leibniz identity*

$$d(a \cdot b) = da \cdot b + (-1)^i a \cdot db \quad \text{for } a \in A^i, b \in A. \tag{A.1}$$

The cohomology $H[A, d] = \operatorname{Ker} d / \operatorname{Im} d$ of a differential graded algebra A is a graded algebra (an exercise). Differential graded algebras whose differential lowers the degree by one are also considered, in which case homology is a graded algebra.

Exercises.

A.1.1. If A is a finitely generated algebra, then all bases of a free A-module have the same cardinality.

A.1.2. A module is projective if and only if it is a direct summand of a free module.

A.1.3. Cochain homotopic maps between cochain complexes induce the same maps in cohomology.

A.1.4. The cohomology of a differential graded algebra is a graded algebra.

A.2. Homological theory of graded rings and modules

From now on we assume that A is a commutative finitely generated $\mathbf{k}$-algebra with unit, graded by nonnegative even numbers (i.e. $A = \bigoplus_{i \geqslant 0} A^i$) and connected (i.e. $A^0 = \mathbf{k}$). The basic example to keep in mind is $A = \mathbf{k}[m] = \mathbf{k}[v_1, \ldots, v_m]$ with $\deg v_i = 2$, however we shall need a greater generality occasionally. We also assume that all A-modules M are nonnegatively graded and finitely generated, and all module maps are degree-preserving, unless the contrary is explicitly stated.

A *free* (respectively, *projective*) *resolution* of M is an exact sequence of A-modules

$$\cdots \xrightarrow{d} R^{-i} \xrightarrow{d} \cdots \xrightarrow{d} R^{-1} \xrightarrow{d} R^0 \to M \to 0 \tag{A.2}$$

in which all R^{-i} are free (respectively, projective) A-modules. A free resolution exists for every M (an exercise, or see constructions below). The minimal number p for which there exists a projective resolution (A.2) with $R^{-i} = 0$ for $i > p$ is called the *projective* (or *homological*) *dimension* of the module M; we shall denote it by $\operatorname{pdim}_A M$ or shortly $\operatorname{pdim} M$. If such p does not exist, we set $\operatorname{pdim} M = \infty$. The module $M_i = \operatorname{Ker}[d \colon R^{-i+1} \to R^{-i+2}]$ is called the *ith syzygy module* for M.

If $\mathbf{k}$ is a field and A is as above, then an A-module is projective if and only if it is free (see Exercise A.2.15), and we therefore may not distinguish between free and projective resolutions in this case.

We can convert a resolution (A.2) into a bigraded $\mathbf{k}$-vector space $R = \bigoplus_{i,j} R^{-i,j}$ where $R^{-i,j} = (R^{-i})^j$ is the jth graded component of the module R^{-i}, and the $(-i, j)$th component of d acts as $d^{-i,j} \colon R^{-i,j} \to R^{-i+1,j}$. We refer to the first grading of R as *external*; it comes from the indexing of the terms in the resolution and is therefore nonpositive by our convention. The second, *internal*, grading of R comes from the grading in the modules R^{-i} and is therefore even and nonnegative. The *total* degree of an element of R is defined as the sum of its external and internal degrees. We can view A as a bigraded algebra with trivial first grading (i.e. $A^{i,j} = 0$ for $i \neq 0$ and $A^{0,j} = A^j$); then R becomes a bigraded A-module.

If the drop the term M in resolution (A.2), then the resulting cochain complex is exactly R (with respect to its external grading), and the bigraded cohomology

$H[R,d]$ satisfies that

$$\begin{aligned} H^{-i,j}[R,d] &= \operatorname{Ker} d^{-i,j} / \operatorname{Im} d^{-i-1,j} = 0 \quad \text{for } i > 0,\\ H^{0,j}[R,d] &= M^j. \end{aligned}$$

We may view M as a trivial cochain complex $0 \to M \to 0$, or as a bigraded module with trivial external grading, i.e. $M^{i,j} = 0$ for $i \neq 0$ and $M^{0,j} = M^j$. Then resolution (A.2) can be interpreted as a map of cochain complexes of A-modules:

$$\tag{A.3} \begin{tikzcd} \cdots \arrow[r] & R^{-i} \arrow[r, "d"] \arrow[d] & \cdots \arrow[r, "d"] & R^{-1} \arrow[r, "d"] \arrow[d] & R^0 \arrow[r] \arrow[d] & 0 \\ \cdots \arrow[r] & 0 \arrow[r] & \cdots \arrow[r] & 0 \arrow[r] & M \arrow[r] & 0, \end{tikzcd}$$

or shortly as a map $[R,d] \to [M,0]$, inducing an isomorphism in cohomology.

The *Poincaré series* of a graded $\mathbf{k}$-vector space $V = \bigoplus_i V^i$ whose graded components are finite-dimensional is given by

$$F(V;\lambda) = \sum_i (\dim_{\mathbf{k}} V^i)\lambda^i.$$

Proposition A.2.1. *Let* (A.2) *be a finite free resolution of an A-module, in which R^{-i} is a free module of rank q_i on generators of degrees $d_{1i}, \ldots, d_{q_i i}$, for $i \geqslant 0$. Then*

$$F(M;\lambda) = F(A;\lambda) \sum_{i \geqslant 0} (-1)^i (\lambda^{d_{1i}} + \cdots + \lambda^{d_{q_i i}}).$$

Proof. Since $H^{-i,j}[R,d] = 0$ for $i > 0$ and $H^{0,j}[R,d] = M^j$, we obtain

$$\sum_{i \geqslant 0} (-1)^i \dim_{\mathbf{k}} R^{-i,j} = \dim_{\mathbf{k}} M^j$$

by a basic property of the Euler characteristic. Multiplying by λ^j and summing up over j we obtain

$$\sum_{i \geqslant 0} (-1)^i F(R^{-i};\lambda) = F(M;\lambda).$$

Since each R^{-i} is a free A-module, its Poincaré series is given by $F(R^{-i};\lambda) = F(A;\lambda)(\lambda^{d_{1i}} + \cdots + \lambda^{d_{q_i i}})$, which implies the required formula. □

Construction A.2.2 (minimal resolution). Let $\mathbf{k}$ be a field, and let $M = \bigoplus_{i \geqslant 0} M^i$ be a graded A-module, which is not necessarily finitely generated, but whose every graded component M^i is finite-dimensional as a $\mathbf{k}$-vector space. There is the following canonical way of constructing a free resolution for M.

Take the lowest degree i in which $M^i \neq 0$ and choose a $\mathbf{k}$-vector space basis in M^i. Span an A-submodule M_1 by this basis and then take the lowest degree in which $M \neq M_1$. In this degree choose a $\mathbf{k}$-vector space basis in the complement of M_1, and span a module M_2 by this basis and M_1. Continuing this process we obtain a system of generators for M which has a finite number of elements in each degree, and has the property that images of the generators form a basis of the $\mathbf{k}$-vector space $M \otimes_A \mathbf{k} = M/(A^+ \cdot M)$. A system of generators of M obtained in this way is referred to as *minimal* (or as a *minimal basis*).

Now choose a minimal generating set in M and span by its elements a free A-module $R^0_{\min}$. Then we have an epimorphism $R^0_{\min} \to M$. Next we choose a minimal basis in the kernel of this epimorphism, and span by it a free module $R^{-1}_{\min}$. Then choose a minimal basis in the kernel of the map $R^{-1}_{\min} \to R^0_{\min}$, and so on. On the ith step we choose a minimal basis in the kernel of the map $d\colon R^{-i+1}_{\min} \to R^{-i+2}_{\min}$ constructed on the previous step, and span a free module $R^{-i}_{\min}$ by this basis. As a result we obtain a free resolution of M, which is referred to as *minimal.* A minimal resolution is unique up to an isomorphism.

Proposition A.2.3. *For a minimal resolution of M, the induced maps*

$$R^{-i}_{\min} \otimes_A \mathbf{k} \to R^{-i+1}_{\min} \otimes_A \mathbf{k}$$

are zero for $i \geqslant 1$.

Proof. By construction, the map $R^0_{\min} \otimes_{\mathbf{k}} A \to M \otimes_{\mathbf{k}} A$ is an isomorphism, which implies that the kernel of the map $R^0_{\min} \to M$ is contained in $A^+ \cdot R^0_{\min}$. Similarly, for each $i \geqslant 1$ the kernel of the map $d\colon R^{-i}_{\min} \to R^{-i+1}_{\min}$ is contained in $A^+ \cdot R^{-i}_{\min}$, and therefore the image of the same map is contained in $A^+ \cdot R^{-i+1}_{\min}$. This implies that the induced maps $R^{-i}_{\min} \otimes_A \mathbf{k} \to R^{-i+1}_{\min} \otimes_A \mathbf{k}$ are zero. □

Remark. If $\mathbf{k} = \mathbb{Z}$ then the above described inductive procedure still gives a minimal basis for an A-module M, but the kernel of the map $d\colon R^0_{\min} \to M$ may not be contained in $A^+ \cdot R^0_{\min}$, and the induced map $R^{-1}_{\min} \otimes_A \mathbb{Z} \to R^0_{\min} \otimes_A \mathbb{Z}$ may be nonzero.

Construction A.2.4 (Koszul resolution). Let $A = \mathbf{k}[v_1, \ldots, v_m]$ and $M = \mathbf{k}$ with the A-module structure given by the augmentation map sending each v_i to zero. We turn the tensor product

$$E = E_m = \Lambda[u_1, \ldots, u_m] \otimes \mathbf{k}[v_1, \ldots, v_m]$$

into a *bigraded differential algebra* by setting

$$\begin{aligned} \operatorname{bideg} u_i &= (-1, 2), \quad \operatorname{bideg} v_i = (0, 2), \\ du_i &= v_i, \quad dv_i = 0 \end{aligned} \tag{A.4}$$

and requiring d to satisfy the Leibniz identity (A.1). Then $[E, d]$ together with the augmentation map $\varepsilon\colon E \to \mathbf{k}$ defines a cochain complex of $\mathbf{k}[m]$-modules

$$\begin{aligned} 0 \to \Lambda^m[u_1, \ldots, u_m] \otimes \mathbf{k}[v_1, \ldots, v_m] &\xrightarrow{d} \cdots \\ \xrightarrow{d} \Lambda^1[u_1, \ldots, u_m] \otimes \mathbf{k}[v_1, \ldots, v_m] &\xrightarrow{d} \mathbf{k}[v_1, \ldots, v_m] \xrightarrow{\varepsilon} \mathbf{k} \to 0, \end{aligned} \tag{A.5}$$

where $\Lambda^i[u_1, \ldots, u_m]$ is the subspace of $\Lambda[u_1, \ldots, u_m]$ generated by monomials of length i. We shall show that the complex above is an exact sequence, or equivalently, that $\varepsilon\colon [E, d] \to [\mathbf{k}, 0]$ is a quasi-isomorphism. There is an obvious inclusion $\eta\colon \mathbf{k} \to E$ such that $\varepsilon\eta = \mathrm{id}$. To finish the proof we shall construct a cochain homotopy between id and $\eta\varepsilon$, that is, a set of $\mathbf{k}$-linear maps $s = \{s^{-i,2j}\colon E^{-i,2j} \to E^{-i-1,2j}\}$ satisfying the identity

$$ds + sd = \mathrm{id} - \eta\varepsilon. \tag{A.6}$$

For $m = 1$ we define the map $s_1\colon E_1^{0,*} = \mathbf{k}[v] \to E_1^{-1,*}$ by the formula

$$s_1(a_0 + a_1 v + \cdots + a_j v^j) = u(a_1 + a_2 v + \cdots + a_j v^{j-1}).$$

Then for $f = a_0 + a_1 v + \cdots + a_j v^j \in E_1^{0,*}$ we have $ds_1 f = f - a_0 = f - \eta\varepsilon f$ and $s_1 df = 0$. On the other hand, for $uf \in E_1^{-1,*}$ we have $s_1 d(uf) = uf$ and

$ds_1(uf) = 0$. In any case (A.6) holds. Now we may assume by induction that for $m = k-1$ the required cochain homotopy $s_{k-1}\colon E_{k-1} \to E_{k-1}$ is already constructed. Since $E_k = E_{k-1} \otimes E_1$, $\varepsilon_k = \varepsilon_{k-1} \otimes \varepsilon_1$ and $\eta_k = \eta_{k-1} \otimes \eta_1$, a direct calculation shows that the map

$$s_k = s_{k-1} \otimes \mathrm{id} + \eta_{k-1}\varepsilon_{k-1} \otimes s_1$$

is a cochain homotopy between id and $\eta_k\varepsilon_k$.

Since $\Lambda^i[u_1,\ldots,u_m] \otimes \mathbf{k}[m]$ is a free $\mathbf{k}[m]$-module, (A.5) is a free resolution for the $\mathbf{k}[m]$-module $\mathbf{k}$. It is known as the *Koszul resolution.* It can be shown to be minimal (an exercise).

Let (A.2) be a projective resolution of an A-module M, and N is another A-module. Applying the functor $\otimes_A N$ to (A.3) we obtain a homomorphism of cochain complexes

$$[R \otimes_A N, d] \to [M \otimes_A N, 0],$$

which does not induce a cohomology isomorphism in general. The $(-i)$th graded cohomology module of the cochain complex

$$\cdots \to R^{-i} \otimes_A N \to \cdots \to R^{-1} \otimes_A N \to R^0 \otimes_A N \to 0 \tag{A.7}$$

is denoted by $\mathrm{Tor}_A^{-i}(M,N)$. We shall also consider the bigraded A-module

$$\mathrm{Tor}_A(M,N) = \bigoplus_{i,j\geqslant 0} \mathrm{Tor}_A^{-i,j}(M,N)$$

where $\mathrm{Tor}_A^{-i,j}(M,N)$ is the jth graded component of $\mathrm{Tor}_A^{-i}(M,N)$

The following properties of $\mathrm{Tor}_A^{-i}(M,N)$ are well-known (see e.g. [**145**]).

Theorem A.2.5.

(a) *The module* $\mathrm{Tor}_A^{-i}(M,N)$ *does not depend, up to isomorphism, on a choice of resolution* (A.2)*;*

(b) $\mathrm{Tor}_A^{-i}(\,\cdot\,,N)$ *and* $\mathrm{Tor}_A^{-i}(M,\,\cdot\,)$ *are covariant functors;*

(c) $\mathrm{Tor}_A^{0}(M,N) = M \otimes_A N$*;*

(d) $\mathrm{Tor}_A^{-i}(M,N) \cong \mathrm{Tor}_A^{-i}(N,M)$*;*

(e) *An exact sequence of A-modules*

$$0 \longrightarrow M_1 \longrightarrow M_2 \longrightarrow M_3 \longrightarrow 0$$

induces the following long exact sequence:

$$\begin{aligned}\cdots \longrightarrow \mathrm{Tor}_A^{-i}(M_1,N) \longrightarrow \mathrm{Tor}_A^{-i}(M_2,N) \longrightarrow \mathrm{Tor}_A^{-i}(M_3,N) \longrightarrow \cdots\\ \cdots \longrightarrow \mathrm{Tor}_A^{-1}(M_1,N) \longrightarrow \mathrm{Tor}_A^{-1}(M_2,N) \longrightarrow \mathrm{Tor}_A^{-1}(M_3,N)\\ \longrightarrow \mathrm{Tor}_A^{0}(M_1,N) \longrightarrow \mathrm{Tor}_A^{0}(M_2,N) \longrightarrow \mathrm{Tor}_A^{0}(M_3,N) \longrightarrow 0.\end{aligned}$$

In the case $N = \mathbf{k}$ the Tor-modules can be read from a minimal resolution of M as follows:

Proposition A.2.6. *Let* $\mathbf{k}$ *be a field, and let* (A.2) *be a minimal resolution of an A-module M. Then*

$$\mathrm{Tor}_A^{-i}(M,\mathbf{k}) \cong R_{\min}^{-i} \otimes_A \mathbf{k},$$

$$\dim_{\mathbf{k}} \mathrm{Tor}_A^{-i}(M,\mathbf{k}) = \operatorname{rank} R_{\min}^{-i}.$$

Proof. Indeed, the differentials in the cochain complex

$$\cdots \longrightarrow R^{-i}_{\min} \otimes_A \mathbf{k} \longrightarrow \cdots \longrightarrow R^{-1}_{\min} \otimes_A \mathbf{k} \longrightarrow R^0_{\min} \otimes_A \mathbf{k} \longrightarrow 0$$

are all trivial by Proposition A.2.3. □

Corollary A.2.7. *Let* $\mathbf{k}$ *be a field, and let* M *be a* A*-module. Then*

$$\operatorname{pdim} M = \max\bigl\{i\colon\ \operatorname{Tor}_A^{-i}(M, \mathbf{k}) \neq 0\bigr\}.$$

Corollary A.2.8. *If* $\mathbf{k}$ *is a field, then* $\operatorname{pdim} M \leqslant m$ *for any* $\mathbf{k}[v_1, \ldots, v_m]$*-module* M*.*

Proof. By the previous corollary and Theorem A.2.5 (d),

$$\operatorname{pdim} M = \max\bigl\{i\colon\ \operatorname{Tor}^{-i}_{\mathbf{k}[m]}(M, \mathbf{k}) \neq 0\bigr\} = \max\bigl\{i\colon\ \operatorname{Tor}^{-i}_{\mathbf{k}[m]}(\mathbf{k}, M) \neq 0\bigr\}.$$

Using the Koszul resolution for the $\mathbf{k}[m]$-module $\mathbf{k}$ we obtain

$$\begin{aligned}\operatorname{Tor}^{-i}_{\mathbf{k}[m]}(\mathbf{k}, M) = H^{-i}\bigl[\Lambda[u_1, \ldots, u_m] \otimes \mathbf{k}[m] \otimes_{\mathbf{k}[m]} M, d\bigr]&\\ = H^{-i}\bigl[\Lambda[u_1, \ldots, u_m] \otimes M, d\bigr].&\end{aligned}$$

Therefore,

$$\operatorname{pdim} M = \max\bigl\{i\colon\ \operatorname{Tor}^{-i}_{\mathbf{k}[m]}(\mathbf{k}, M) \neq 0\bigr\} \leqslant \max\bigl\{i\colon \Lambda^i[u_1, \ldots, u_m] \otimes M \neq 0\bigr\} = m.$$

□

Example A.2.9. Let $A = \mathbf{k}[v_1, \ldots, v_m]$ and $M = N = \mathbf{k}$. By the minimality of the Koszul resolution,

$$\operatorname{Tor}_{\mathbf{k}[v_1,\ldots,v_m]}(\mathbf{k}, \mathbf{k}) = \Lambda[u_1, \ldots, u_m]$$

and $\operatorname{pdim}_A \mathbf{k} = m$.

We note that in general there is no canonical way to define a multiplication in $\operatorname{Tor}_A(M, N)$, even if both M and N are A-algebras rather than just A-modules. However, in the particular case when $A = \mathbf{k}[m]$, M is an algebra with a unit and $N = \mathbf{k}$ there is the following canonical way to define a product in $\operatorname{Tor}_A(M, N)$, extending the previous example. We consider the differential bigraded algebra $[\Lambda[u_1, \ldots, u_m] \otimes M, d]$ whose bigrading and differential are defined similarly to (A.4):

$$\begin{aligned}&\operatorname{bideg} u_i = (-1, 2), \quad \operatorname{bideg} x = (0, \deg x) \quad \text{for } x \in M,\\ &du_i = v_i \cdot 1, \quad dx = 0\end{aligned} \tag{A.8}$$

(here $v_i \cdot 1$ is the element of M obtained by applying $v_i \in \mathbf{k}[m]$ to $1 \in M$, and we identify u_i with $u_i \otimes 1$ and x with $1 \otimes x$ for simplicity). Using the fact that the cohomology of a differential graded algebra is a graded algebra we obtain:

Lemma A.2.10. *Let* M *be a graded* $\mathbf{k}[v_1, \ldots, v_m]$*-algebra. Then* $\operatorname{Tor}_{\mathbf{k}[m]}(M, \mathbf{k})$ *is a bigraded* $\mathbf{k}$*-algebra whose product is defined via the isomorphism*

$$\operatorname{Tor}_{\mathbf{k}[v_1,\ldots,v_m]}\bigl(M, \mathbf{k}\bigr) \cong H\bigl[\Lambda[u_1, \ldots, u_m] \otimes M, d\bigr].$$

Proof. Using the Koszul resolution in the definition of $\operatorname{Tor}_{\mathbf{k}[m]}(\mathbf{k}, M)$ and Theorem A.2.5 (d) we calculate

$$\begin{aligned}&\operatorname{Tor}_{\mathbf{k}[m]}(M, \mathbf{k}) \cong \operatorname{Tor}_{\mathbf{k}[m]}(\mathbf{k}, M)\\ &\qquad = H\bigl[\Lambda[u_1, \ldots, u_m] \otimes \mathbf{k}[m] \otimes_{\mathbf{k}[m]} M, d\bigr] \cong H\bigl[\Lambda[u_1, \ldots, u_m] \otimes M, d\bigr].\end{aligned}$$ □

The algebra $\big[\Lambda[u_1,\dots,u_m]\otimes M, d\big]$ is known as the *Koszul algebra* (or *Koszul complex*) of M.

REMARK. Lemma A.2.10 holds also in the case when M does not have unit (e.g., when it is a graded ideal in $\mathbf{k}[m]$). In this case formula (A.8) for the differential needs to be updated as follows:

$$d(u_i x) = v_i \cdot x, \quad dx = 0 \quad \text{for } x \in M.$$

We finish this discussion of $\mathrm{Tor}_{\mathbf{k}[m]}(M,\mathbf{k})$ by mentioning an important conjecture of commutative homological algebra.

CONJECTURE A.2.11 (Horrocks, see [**36**, p. 453]). *Let M be a graded $\mathbf{k}[v_1,\dots,v_m]$-module such that $\dim_{\mathbf{k}} M<\infty$, where $\mathbf{k}$ is a field. Then*

$$\dim_{\mathbf{k}} \mathrm{Tor}^{-i}_{\mathbf{k}[v_1,\dots,v_m]}(M,\mathbf{k}) \geqslant \tbinom{m}{i}.$$

It is sometimes formulated in a weaker form:

CONJECTURE A.2.12 (weak Horrocks' Conjecture). *Let M be a graded $\mathbf{k}[v_1,\dots,v_m]$-module such that $\dim_{\mathbf{k}} M<\infty$, where $\mathbf{k}$ is a field. Then*

$$\dim_{\mathbf{k}} \mathrm{Tor}_{\mathbf{k}[v_1,\dots,v_m]}(M,\mathbf{k}) \geqslant 2^m.$$

If the algebra A is not necessarily commutative, then $\mathrm{Tor}_A(M,N)$ is defined for a right A-module M and a left A-module N in the same way as above. However, in this case $\mathrm{Tor}_A(M,N)$ is no longer an A-module, and is just a $\mathbf{k}$-vector space. If both M and N are A-bimodules, then $\mathrm{Tor}_A(M,N)$ is an A-bimodule itself.

The construction of Tor can also be extended to the case of differential graded modules and algebras, see Section B.4.

In the standard notation adopted in the algebraic literature, the modules in a resolution (A.2) are numbered by nonnegative rather than nonpositive integers:

$$\cdots \xrightarrow{d} R^i \xrightarrow{d} \cdots \xrightarrow{d} R^1 \xrightarrow{d} R^0 \to M \to 0.$$

In this notation, the ith Tor-module is denoted by $\mathrm{Tor}_i^A(M,N)$, $\ i\geqslant 0$ (note that (A.7) becomes a chain complex, and $\mathrm{Tor}_*^A(M,N)$ is its homology). Therefore, the two notations are related by

$$\mathrm{Tor}_A^{-i}(M,N) = \mathrm{Tor}_i^A(M,N).$$

Applying the functor $\mathrm{Hom}_A(\quad, N)$ to (A.3) (with R^{-i} replaced by R^i) we obtain the cochain complex

$$0 \to \mathrm{Hom}_A(R^0,N) \to \mathrm{Hom}_A(R^1,N) \to \cdots \to \mathrm{Hom}_A(R^i,N) \to \cdots .$$

Its ith cohomology module is denoted by $\mathrm{Ext}^i_A(M,N)$.

The properties of the functor Ext are similar to those given by Theorem A.2.5 for Tor, with the exception of (d):

THEOREM A.2.13.

(a) *The module $\mathrm{Ext}^i_A(M,N)$ does not depend, up to isomorphism, on a choice of resolution* (A.2)*;*

(b) $\mathrm{Ext}^i_A(\,\cdot\,,N)$ *is a contravariant functor, and* $\mathrm{Ext}^i_A(M,\,\cdot\,)$ *is a covariant functor;*

(c) $\mathrm{Ext}^0_A(M,N)=\mathrm{Hom}_A(M,N)$*;*

(d) *An exact sequence of A-modules*

$$0 \longrightarrow M_1 \longrightarrow M_2 \longrightarrow M_3 \longrightarrow 0$$

induces the following long exact sequence:

$$\begin{aligned}0 &\longrightarrow \mathrm{Ext}^0_A(M_3,N) \longrightarrow \mathrm{Ext}^0_A(M_2,N) \longrightarrow \mathrm{Ext}^0_A(M_1,N)\\ &\longrightarrow \mathrm{Ext}^1_A(M_3,N) \longrightarrow \mathrm{Ext}^1_A(M_2,N) \longrightarrow \mathrm{Ext}^1_A(M_1,N) \longrightarrow \cdots\\ \cdots &\longrightarrow \mathrm{Ext}^i_A(M_3,N) \longrightarrow \mathrm{Ext}^i_A(M_2,N) \longrightarrow \mathrm{Ext}^i_A(M_1,N) \longrightarrow \cdots;\end{aligned}$$

(e) *An exact sequence of A-modules*

$$0 \longrightarrow N_1 \longrightarrow N_2 \longrightarrow N_3 \longrightarrow 0$$

induces the following long exact sequence:

$$\begin{aligned}0 &\longrightarrow \mathrm{Ext}^0_A(M,N_1) \longrightarrow \mathrm{Ext}^0_A(M,N_2) \longrightarrow \mathrm{Ext}^0_A(M,N_3)\\ &\longrightarrow \mathrm{Ext}^1_A(M,N_1) \longrightarrow \mathrm{Ext}^1_A(M,N_2) \longrightarrow \mathrm{Ext}^1_A(M,N_3) \longrightarrow \cdots\\ \cdots &\longrightarrow \mathrm{Ext}^i_A(M,N_1) \longrightarrow \mathrm{Ext}^i_A(M,N_2) \longrightarrow \mathrm{Ext}^i_A(M,N_3) \longrightarrow \cdots.\end{aligned}$$

Exercises.

A.2.14. Show that a free resolution exists for every A-module M. (Hint: use the fact that every module is the quotient of a free module.)

A.2.15. If $\mathbf{k}$ is a field and $A = \mathbf{k}[m]$, then every projective graded A-module is free (hint: see [**145**, Lemma VII.6.2]). This is also true in the ungraded case, but is much harder to prove (a theorem of Quillen and Suslin, settling the famous problem of Serre). More generally, if A is a finitely generated nonnegatively graded commutative connected algebra over a field $\mathbf{k}$, then every projective A-module is free (see [**84**, Theorem A3.2]). Give an example of a projective module over a ring which is not free.

A.2.16. The Koszul resolution is minimal.

A.3. Regular sequences and Cohen–Macaulay algebras

Cohen–Macaulay algebras and modules play an important role in commutative algebra, algebraic geometry and combinatorics. Their definition uses the notion of a regular sequence (see Definition A.3.1 below), which also plays an important role in algebraic topology, namely in the construction of new cohomology theories (see [**138**] and Appendix Section F.3). In the case of finitely generated algebras over a field $\mathbf{k}$, an algebra is Cohen–Macaulay if and only if it is a free module of finite rank over a polynomial subalgebra.

Here we consider nonnegatively evenly graded finitely generated commutative connected algebras A over a field $\mathbf{k}$ and finitely generated nonnegatively graded A-modules M (the case $\mathbf{k} = \mathbb{Z}$ requires extra care, and is treated separately in some particular cases in the main chapters of the book). The positive part A^+ is the unique homogeneous maximal ideal of A, and the results we discuss here are parallel to those from the homological theory of Noetherian local rings (we refer to [**35**, Chapters 1–2] or [**84**, Chapter 19] for the details).

Given a sequence of elements $\boldsymbol{t} = (t_1, \dots, t_k)$ of A, we denote by $A/\boldsymbol{t}$ the quotient algebra of A by the ideal generated by $\boldsymbol{t}$, and denote by $M/\boldsymbol{t}M$ the quotient module of M by the submodule $t_1M + \cdots + t_kM$. An element $t \in A$ is called a *zero divisor* on M if $tx = 0$ for some nonzero $x \in M$. An element $t \in A$ is not a zero divisor on M if and only if the map $M \xrightarrow{t} M$ given by multiplication by t is injective.

Definition A.3.1. Let M be an A-module. A homogeneous sequence $\boldsymbol{t} = (t_1, \dots, t_k) \in \mathcal{H}(A^+)$ is called an *M-regular sequence* if t_{i+1} is not a zero divisor on $M/(t_1M + \cdots + t_iM)$ for $0 \leqslant i < k$. We often refer to A-regular sequences simply as *regular*.

The importance of regular sequences in homological algebra builds on the fundamental fact that an exact sequence of modules remains exact after taking quotients by a regular sequence:

Proposition A.3.2. *Assume given an exact sequence of A-modules:*

$$\cdots \longrightarrow S^i \xrightarrow{f_i} S^{i-1} \xrightarrow{f_{i-1}} \cdots \xrightarrow{f_1} S^0 \xrightarrow{f_0} M \to 0$$

If $\boldsymbol{t}$ is an M-regular and S^i-regular sequence for all $i \geqslant 0$, then the sequence of $A/\boldsymbol{t}$-modules

$$\cdots \to S^i/\boldsymbol{t}S^i \xrightarrow{\overline{f}_i} S^{i-1}/\boldsymbol{t}S^{i-1} \xrightarrow{\overline{f}_{i-1}} \cdots \xrightarrow{\overline{f}_1} S^0/\boldsymbol{t}S^0 \xrightarrow{\overline{f}_0} M/\boldsymbol{t}M \to 0$$

is also exact.

Proof. Using induction we reduce the statement to the case when $\boldsymbol{t}$ consists of a single element t. Since

$$S^i/tS^i = S^i \otimes_A (A/t),$$

and $\otimes_A(A/t)$ is a right exact functor, it is enough to verify exactness of the quotient sequence starting from the term S^1/tS^1.

Consider the following fragment of the quotient sequence ($i \geqslant 1$):

$$S^{i+1}/tS^{i+1} \xrightarrow{\overline{f}_{i+1}} S^i/tS^i \xrightarrow{\overline{f}_i} S^{i-1}/tS^{i-1} \xrightarrow{\overline{f}_{i-1}} S^{i-2}/tS^{i-2}$$

(where we denote $S^{-1} = M$). For any element $x \in S^i$ we denote by $\overline{x}$ its residue class in S^i/tS^i. Let $\overline{f}_i(\overline{x}) = 0$, then $f_i(x) = ty$ for some $y \in S^{i-1}$ and $tf_{i-1}(y) = 0$. Since t is S^{i-2}-regular, we have $f_{i-1}(y) = 0$. Hence, there is $x' \in S^i$ such that $y = f_i(x')$. This implies that $f_i(x - tx') = 0$. Therefore, $x - tx' \in f_{i+1}(S^{i+1})$ and $\overline{x} \in \overline{f}_{i+1}(S^{i+1}/tS^{i+1})$. Thus, the quotient sequence is exact. $\square$

The following proposition is often used as the definition of regular sequences:

Proposition A.3.3. *A sequence $t_1, \dots, t_k \in \mathcal{H}(A^+)$ is M-regular if and only if M is a free (not necessarily finitely generated) $\mathbf{k}[t_1, \dots, t_k]$-module.*

Proof. If M is a free $\mathbf{k}[t_1, \dots, t_k]$-module, then $M/(t_1M + \cdots + t_{i-1}M)$ is a free $\mathbf{k}[t_i, \dots, t_k]$-module, which implies that t_i is $M/(t_1M + \cdots + t_{i-1}M)$-regular for $1 \leqslant i \leqslant k$. Therefore, $t_1, \dots, t_k$ is an M-regular sequence.

Conversely, let $\boldsymbol{t} = (t_1, \dots, t_k)$ be an M-regular sequence. Consider a minimal resolution $[R_{\min}, d]$ for the $\mathbf{k}[\boldsymbol{t}]$-module M. Then, by Proposition A.3.2, the sequence of $\mathbf{k}$-modules

$$\cdots \longrightarrow R^{-1}_{\min}/\boldsymbol{t}R^{-1}_{\min} \longrightarrow R^0_{\min}/\boldsymbol{t}R^0_{\min} \longrightarrow M/\boldsymbol{t}M \longrightarrow 0$$

is exact. Note that $R^{-i}_{\min}/\boldsymbol{t}R^{-i}_{\min} = R^{-i}_{\min} \otimes_{\mathbf{k}[\boldsymbol{t}]} \mathbf{k}$. Since the resolution is minimal, the map $R^0_{\min} \otimes_{\mathbf{k}[\boldsymbol{t}]} \mathbf{k} \to M \otimes_{\mathbf{k}[\boldsymbol{t}]} \mathbf{k}$ is an isomorphism. Hence, $R^{-i}_{\min} \otimes_{\mathbf{k}[\boldsymbol{t}]} \mathbf{k} = 0$ for

$i > 0$, which implies that $R^{-i}_{\min} = 0$. Thus, $R^0_{\min} \to M$ is an isomorphism, i.e. M is a free $\mathbf{k}[\boldsymbol{t}]$-module. ☐

The following is a direct corollary of Proposition A.3.13:

PROPOSITION A.3.4. *The property of being a regular sequence does not depend on the order of elements in* $\boldsymbol{t} = (t_1, \dots, t_k)$.

LEMMA A.3.5. *Let* $\boldsymbol{t}$ *be a sequence of elements of* A *which is* A*-regular and* M*-regular. Then*

$$\mathrm{Tor}^{-i}_A(M, \mathbf{k}) = \mathrm{Tor}^{-i}_{A/\boldsymbol{t}}(M/\boldsymbol{t}M, \mathbf{k}).$$

PROOF. Applying Proposition A.3.2 to a minimal resolution of M, we obtain a minimal resolution of the $A/\boldsymbol{t}$-module $M/\boldsymbol{t}M$. The rest follows from Proposition A.2.6. ☐

An M-regular sequence is *maximal* if it is not contained in an M-regular sequence of greater length.

THEOREM A.3.6 (D. Rees). *All maximal* M*-regular sequences in* A *have the same length given by*

$$\mathrm{depth}_A M = \min\bigl\{i\colon \mathrm{Ext}^i_A(\mathbf{k}, M) \ne 0\bigr\}. \tag{A.9}$$

This number given by (A.9) is referred to as the *depth* of M; the simplified notation $\mathrm{depth}\, M$ will be used whenever it creates no confusion. The proof of Theorem A.3.6 uses the following fact:

LEMMA A.3.7. *Let* $\boldsymbol{t} = (t_1, \dots, t_n) \in \mathcal{H}(A^+)$ *be a* M*-regular sequence. Then*

$$\mathrm{Ext}^n_A(\mathbf{k}, M) \cong \mathrm{Hom}_A(\mathbf{k}, M/\boldsymbol{t}M).$$

PROOF. We use induction on n. The case $n = 0$ is tautological. It follows from Lemma A.3.4 that t_n is an M-regular element, so we have the exact sequence

$$0 \longrightarrow M \xrightarrow{\cdot t_n} M \longrightarrow M/t_nM \longrightarrow 0.$$

The map $\mathrm{Ext}^i_A(\mathbf{k}, M) \to \mathrm{Ext}^i_A(\mathbf{k}, M)$ induced by multiplication by t_n is zero (an exercise). Therefore, the second long exact sequence for Ext (see Theorem A.2.13 (e)) induced by the short exact sequence above splits into short exact sequences of the form

$$0 \longrightarrow \mathrm{Ext}^{n-1}_A(\mathbf{k}, M) \longrightarrow \mathrm{Ext}^{n-1}_A(\mathbf{k}, M/t_nM) \longrightarrow \mathrm{Ext}^n_A(\mathbf{k}, M) \longrightarrow 0.$$

Let $\boldsymbol{t}' = (t_1, \dots, t_{n-1})$. By induction,

$$\mathrm{Ext}^{n-1}_A(\mathbf{k}, M) \cong \mathrm{Hom}_A(\mathbf{k}, M/\boldsymbol{t}'M) = 0,$$

where the latter identity follows from Exercise A.3.17, since t_n is $M/\boldsymbol{t}'M$-regular. Now the exact sequence above implies that

$$\mathrm{Ext}^n_A(\mathbf{k}, M) \cong \mathrm{Ext}^{n-1}_A(\mathbf{k}, M/t_nM) \cong \mathrm{Hom}_A(\mathbf{k}, M/\boldsymbol{t}M),$$

where the latter identity follows by induction. ☐

PROOF OF THEOREM A.3.6. Let $\boldsymbol{t} = (t_1, \dots, t_n)$ be a maximal M-regular sequence. Then, by Lemma A.3.7 and Exercise A.3.17,

$$\mathrm{Ext}^n_A(\mathbf{k}, M) \cong \mathrm{Hom}_A(\mathbf{k}, M/\boldsymbol{t}M) \ne 0,$$

as A does not contain an $M/\boldsymbol{t}M$-regular element. On the other hand,

$$\operatorname{Ext}^i_A(\mathbf{k}, M) \cong \operatorname{Hom}_A\big(\mathbf{k}, M/(t_1M + \cdots + t_iM)\big) = 0$$

for $i < n$, since t_{i+1} is $M/(t_1M + \cdots + t_iM)$-regular. $\square$

The following fundamental result relates the depth to the projective dimension of a module.

Theorem A.3.8 (Auslander–Buchsbaum). *Let $M \neq 0$ be an A-module such that* $\operatorname{pdim} M < \infty$. *Then*

$$\operatorname{pdim} M + \operatorname{depth} M = \operatorname{depth} A.$$

Proof. First let $\operatorname{depth} A = 0$. Assume that $\operatorname{pdim} M = p > 0$. Consider the minimal resolution for M (which is finite by hypothesis):

$$0 \to R^{-p}_{\min} \xrightarrow{d_p} R^{-p+1}_{\min} \longrightarrow \cdots \longrightarrow R^0_{\min} \longrightarrow M \to 0.$$

Since $\operatorname{depth} A = 0$, we have $\operatorname{Hom}_A(\mathbf{k}, A) = \operatorname{Ext}^0_A(\mathbf{k}, A) \neq 0$ by Theorem A.3.6. Therefore, there is a monomorphism of A-modules $i\colon \mathbf{k} \to A$. In the commutative diagram

$$\begin{tikzcd}
R^{-p} \otimes_A \mathbf{k} \arrow[r, "d_p \otimes_A \mathbf{k}"] \arrow[d, "\mathrm{id} \otimes_A i"'] & R^{-p+1} \otimes_A \mathbf{k} \arrow[d, "\mathrm{id} \otimes_A i"] \\
R^{-p} \arrow[r, "d_p"] & R^{-p+1}
\end{tikzcd}$$

the maps d_p and $\mathrm{id} \otimes_A i$ are injective (the latter because the module R^{-p} is free). Hence, $d_p \otimes_A \mathbf{k}$ is also injective, which contradicts minimality of the resolution. Hence, $\operatorname{pdim} M = 0$, which implies that M is a free A-module and $\operatorname{depth} M = \operatorname{depth} A = 0$.

Now let $\operatorname{depth} A > 0$. Assume that $\operatorname{depth} M = 0$. Consider the first syzygy module $M_1 = \operatorname{Ker}[R^0 \to M]$ for M. It follows from (A.9) and the exact sequence for Ext that $\operatorname{depth} M_1 = 1$. Since $\operatorname{pdim} M_1 = \operatorname{pdim} M - 1$, it is enough to prove the Auslender–Buchsbaum formula for the module M_1. Hence, we may assume that $\operatorname{depth} M > 0$. This implies that there is an element $t \in A$ which is A-regular and M-regular (an exercise). Then

$$\operatorname{depth}_{A/t} A/t = \operatorname{depth}_A A - 1, \quad \operatorname{depth}_{A/t} M/tM = \operatorname{depth}_A M - 1$$

by the definition of depth, and

$$\operatorname{pdim}_{A/t} M/tM = \operatorname{pdim}_A M$$

by Corollary A.2.7 and Lemma A.3.5. Now the proof is finished by induction on $\operatorname{depth} A$. $\square$

The (Krull) *dimension* of A, denoted $\dim A$, is the maximal number of elements of A algebraically independent over $\mathbf{k}$. The *dimension* of an A-module M is defined as $\dim M = \dim(A/\operatorname{Ann} M)$, where

$$\operatorname{Ann} M = \{a \in A\colon aM = 0\}$$

is the *annihilator* of M.

Definition A.3.9. A sequence $t_1, \ldots, t_n$ of algebraically independent homogeneous elements of A is called a *homogeneous system of parameters* (briefly *hsop*) for M if $\dim M/(t_1M+\cdots+t_nM) = 0$. Equivalently, $t_1, \ldots, t_n$ is an hsop if $n = \dim M$ and M is a finitely-generated $\mathbf{k}[t_1, \ldots, t_n]$-module.

The following result (due to Hilbert) is a graded version of the well-known *Noether normalisation lemma*:

Theorem A.3.10 ([**35**, Theorem 1.5.17]). *An hsop exists for any A-module M. If $\mathbf{k}$ is an infinite field and A is generated by degree-two (i.e. linear) elements, then a degree-two hsop can be chosen for M.*

An hsop consisting of linear elements is referred to as a *linear system of parameters* (briefly *lsop*).

It is easy to see that a regular sequence consists of algebraically independent elements, which implies that $\operatorname{depth} M \leqslant \dim M$.

Definition A.3.11. M is a *Cohen–Macaulay A-module* if $\operatorname{depth} M = \dim M$, that is, if A contains an M-regular sequence $t_1, \ldots, t_n$ of length $n = \dim M$. If A is a Cohen–Macaulay A-module, then it is called a *Cohen–Macaulay algebra.*

By Proposition A.3.13, A is a Cohen–Macaulay algebra if and only if it is a free finitely generated module over a polynomial subalgebra.

Proposition A.3.12. *Let M be a Cohen–Macaulay A-module. Then a sequence $\boldsymbol{t} = (t_1, \ldots, t_k) \in \mathcal{H}(A^+)$ is M-regular if and only if it is a part of an hsop for M.*

Proof. Let $\dim M = n$. Assume that $\boldsymbol{t}$ is an M-regular sequence. The fact that t_i is an $M/(t_1M + \cdots + t_{i-1}M)$-regular element implies that

$$\dim M/(t_1M + \cdots + t_iM) = \dim M/(t_1M + \cdots + t_{i-1}M) - 1$$

for $i = 1, \ldots, k$ (an exercise). Therefore, $\dim M/\boldsymbol{t}M = n - k$, i.e. $\boldsymbol{t}$ is a part of an hsop for M.

For the other direction, see [**35**, Theorem 2.1.2 (c)]. □

In particular, any hsop in a Cohen–Macaulay algebra A is regular.

Proposition A.3.13. *Algebra A is Cohen–Macaulay if and only if it is a free finitely generated module over a polynomial subalgebra.*

Proof. Assume that A is Cohen–Macaulay and $\dim A = n$. Then there is a regular sequence $\boldsymbol{t} = (t_1, \ldots, t_n)$ in A. By the previous proposition, $\boldsymbol{t}$ is an hsop, so that $\dim A/\boldsymbol{t} = 0$ and therefore A is a finitely generated $\mathbf{k}[t_1, \ldots, t_n]$-module. This module is also free by Proposition A.3.3.

On the other hand, if A is free finitely generated over $\mathbf{k}[t_1, \ldots, t_n]$ where $t_1, \ldots, t_n \in A$, then $\dim A = n$ and $t_1, \ldots, t_n$ is a regular sequence by Proposition A.3.3, so that A is Cohen–Macaulay. □

Proposition A.3.14. *If A is Cohen–Macaulay with an lsop $\boldsymbol{t} = (t_1, \ldots, t_n)$, then there is the following formula for the Poincaré series of A:*

$$F(A;\lambda) = \frac{F\bigl(A/(t_1, \ldots, t_n);\lambda\bigr)}{(1-\lambda^2)^n},$$

where $F(A/(t_1, \ldots, t_n);\lambda)$ is a polynomial with nonnegative integer coefficients.

Proof. Since A is a free finitely generated module over $\mathbf{k}[t_1,\ldots,t_n]$, we have an isomorphism of $\mathbf{k}$-vector spaces $A\cong(A/\boldsymbol{t})\otimes\mathbf{k}[t_1,\ldots,t_n]$. Calculating the Poincaré series of both sides yields the required formula. $\square$

Remark. If A is generated by its elements $a_1,\ldots,a_n$ of positive degrees $d_1,\ldots,d_n$ respectively, then it may be shown that the Poincaré series of A is a rational function of the form

$$F(A;\lambda)=\frac{P(\lambda)}{(1-\lambda^{d_1})(1-\lambda^{d_2})\cdots(1-\lambda^{d_n})},$$

where $P(\lambda)$ is a polynomial with integer coefficients. However, in general the polynomial $P(\lambda)$ cannot be given explicitly, and some of its coefficients may be negative.

Exercises.

A.3.15. An M-regular sequence consists of algebraically independent elements.

A.3.16. The map $\operatorname{Ext}^i_A(\mathbf{k},M)\to\operatorname{Ext}^i_A(\mathbf{k},M)$ induced by multiplication by an element $x\in\mathcal{H}(A^+)$ is zero.

A.3.17. The following conditions are equivalent for an A-module M:

(a) Every element of $\mathcal{H}(A^+)$ is a zero divisor on M, i.e. $\operatorname{depth}M=0$;
(b) $\operatorname{Hom}_A(\mathbf{k},M)\neq 0$.

(Hint: show that if $\mathcal{H}(A^+)$ consists of zero divisors on M then the ideal A^+ annihilates a homogeneous element of M, see [**84**, Corollary 3.2].)

A.3.18. Let $\operatorname{depth}A>0$, let M be an A-module with $\operatorname{depth}M=0$, and let $M_1=\operatorname{Ker}[R^0\to M]$ be the first syzygy module for M. Then $\operatorname{depth}M_1=1$.

A.3.19. If $\operatorname{depth}A>0$ and $\operatorname{depth}M>0$, then there exists an element $t\in A$ which is A-regular and M-regular.

A.3.20. The Auslander–Buchsbaum formula (Theorem A.3.8) does not hold if $\operatorname{pdim}M=\infty$.

A.3.21. Show that $\dim A=0$ if and only if A is finite-dimensional as a $\mathbf{k}$-vector space. Is it true that $\operatorname{depth}A=0$ implies that $\dim_{\mathbf{k}}A$ is finite?

A.3.22. Give an example of an algebra A over a field $\mathbf{k}$ of finite characteristic which is generated by linear elements, but does not have an lsop.

A.3.23. A regular sequence consists of algebraically independent elements.

A.3.24. If $t\in\mathcal{H}(A^+)$ is an M-regular element, then $\dim M/tM=\dim M-1$.

A.3.25. Let $\mathbf{k}=\mathbb{Z}$. Show that if A is a free finitely generated module over a polynomial subalgebra $\mathbb{Z}[t_1,\ldots,t_k]$ then t_{i+1} is not a zero divisor on $A/(t_1,\ldots,t_i)$ for $0\leqslant i<k$, but the converse is not true. Therefore, the two possible definitions of a regular sequence over $\mathbb{Z}$ do not agree. (The reason why Proposition A.3.13 fails over $\mathbb{Z}$ is that minimal resolutions do not have the required good properties, see the remark after Construction A.2.2.)

A.4. Formality and Massey products

Here we develop the algebraic formalism used in rational homotopy theory. We work with differential graded algebras $A = \bigoplus_{i \geqslant 0} A^i$ over a field $\mathbf{k}$ of zero characteristic (usually $\mathbb{R}$ or $\mathbb{Q}$), and refer to such A as *dg-algebras* for short. We do not assume A to be finitely generated. Commutativity of dg-algebras is always understood in the graded sense.

A dg-algebra A is called *connected* if $H^0[A,d] = \mathbf{k}$. (Note that a connected dg-algebra is not necessarily connected as a graded algebra.) A connected dg-algebra A is *simply connected* if $H^1[A,d] = 0$.

A *homomorphism* between dg-algebras (A, d_A) and (B, d_B) is a $\mathbf{k}$-linear map $f\colon A \to B$ which preserves the degree, i.e. $f(A^i) \subset B^i$, and satisfies $f(ab) = f(a)f(b)$ and $d_B f(a) = f(d_A a)$ for all $a, b \in A$. Such a homomorphism induces a homomorphism $\widetilde{f}\colon H[A, d_A] \to H[B, d_B]$ of cohomology algebras. We refer to f as a *quasi-isomorphism* if $\widetilde{f}$ is an isomorphism. The equivalence relation generated by quasi-isomorphisms of dg-algebras is referred to as the *weak equivalence*. Since quasi-isomorphism are often not invertible, a weak equivalence between A and B implies only the existence of a zigzag of quasi-isomorphisms of the form

$$A \leftarrow A_1 \rightarrow A_2 \leftarrow A_3 \rightarrow \ldots \leftarrow A_k \rightarrow B.$$

Such a 'long' zigzag of quasi-isomorphisms can be reduced to a 'short' zigzag of the form $A \leftarrow M \rightarrow B$ using the notion of a minimal model.

Definition A.4.1. A commutative dg-algebra $M = \bigoplus_{i \geqslant 0} M^i$ is called *minimal* (in the sense of Sullivan) if the following three conditions are satisfied:

(a) $M^0 = \mathbf{k}$ and $d(M^0) = 0$;

(b) M is a free commutative dg-algebra, i.e.

$$M = \Lambda[x_k\colon \deg x_k \text{ is odd}] \otimes \mathbf{k}[x_k\colon \deg x_k \text{ is even}],$$

there are only finitely many generators in each degree, and

$$\deg x_k \leqslant \deg x_l \quad \text{for } k \leqslant l;$$

(c) dx_k is a polynomial on generators $x_1, \ldots, x_{k-1}$, for each $k \geqslant 1$ (this is called the *nilpotence condition* on d).

Clearly, a minimal dg-algebra M is simply connected if and only if $M^1 = 0$. In this case $\deg x_k \geqslant 2$ for each k, and the nilpotence condition is equivalent to the *decomposability* of d, i.e.

$$d(M) \subset M^+ \cdot M^+,$$

where M^+ is the subspace generated by elements of positive degree. For non-simply connected dg-algebras decomposability does not imply nilpotence: the algebra $\Lambda[x,y]$, $\deg x = \deg y = 1$, with $dx = 0$, $dy = xy$ is not minimal.

Definition A.4.2. A minimal dg-algebra M is called a *minimal model* for a commutative dg-algebra A if there is a quasi-isomorphism $h\colon M \to A$.

Theorem A.4.3.

(a) *For each connected commutative dg-algebra A satisfying the condition $\dim H^i[A] < \infty$ for all i, there exists a unique (up to an isomorphism) minimal model $M(A)$.*

(b) *A homomorphism of commutative dg-algebras* $f\colon A \to B$ *lifts to a homomorphism* $\widehat{f}\colon M(A) \to M(B)$ *of their minimal models closing the commutative diagram*

$$\begin{tikzcd}
M(A) \arrow[r, "\widehat{f}"] \arrow[d, "h_A"'] & M(B) \arrow[d, "h_B"] \\
A \arrow[r, "f"] & B
\end{tikzcd}.$$

(c) *If* f *is a quasi-isomorphism, then* $\widehat{f}$ *is an isomorphism.*

Remark. Minimal models can be also defined for dg-algebras A which do not satisfy the finiteness condition $\dim H^i[A] < \infty$, but we shall not need this.

This theorem is due to Sullivan (simply connected case) and Halperin (general). A proof can be found in [**140**, Theorem II.6] or [**89**, §12].

Corollary A.4.4. *A weak equivalence between two commutative dg-algebras* A, B *satisfying the condition of Theorem A.4.3 can be represented by a 'short' zigzag* $A \leftarrow M \to B$ *of quasi-isomorphisms, where* M *is a minimal model for* A *(or* B*).*

Definition A.4.5. A dg-algebra is A called *formal* if it is weakly equivalent to its cohomology $H[A]$ (viewed as a dg-algebra with zero differential).

Corollary A.4.6. *A commutative dg-algebra* A *with a minimal model* $M(A)$ *is formal if and only if* $M(A)$ *in formal. In this case there is a zigzag of quasi-isomorphisms* $A \leftarrow M(A) \to H[A]$.

If A is formal and has a minimal model $M(A)$, then a minimal model can be recovered from the cohomology algebra $H[A]$ using an inductive procedure.

Remark. Even if A is formal, the zigzag $A \leftarrow M(A) \to H[A]$ usually cannot be reduced to a single quasi-isomorphism $A \to H[A]$ or $H[A] \to A$.

Example A.4.7. Let M be a dg-algebra with three generators a_1, a_2, a_3 of degree 1 and differential given by

$$da_1 = da_2 = 0, \quad da_3 = a_1 a_2.$$

This M is minimal, but nor formal. Indeed the first degree cohomology $H^1[M]$ is generated by the classes α_1, α_2 corresponding to the cocycles a_1, a_2, and we have $\alpha_1\alpha_2 = 0$. Assume there is a quasi-isomorphism $f\colon M \to H[M]$; then we have $f(a_3) = k_1\alpha_1 + k_2\alpha_2$ for some $k_1, k_2 \in \mathbf{k}$. This implies that $f(a_1a_3) = 0$, which is impossible since a_1a_3 represents a nontrivial cohomology class.

We next review Massey products, which provide a simple and effective tool for establishing nonformality of a dg-algebra. Massey products constitute a series of higher-order operations (or *brackets*) in the cohomology of a dg-algebra, with the second-order operation coinciding with the cohomology multiplication, while the higher-order brackets are only defined for certain tuples of cohomology classes. We shall only consider triple (third-order) Massey products here.

Construction A.4.8 (triple Massey product). Let A be a dga, and let $\alpha_1, \alpha_2, \alpha_3$ be three cohomology classes such that $\alpha_1\alpha_2 = \alpha_2\alpha_3 = 0$ in $H[A]$. Choose their representing cocycles $a_i \in A^{k_i}$, $i = 1, 2, 3$. Since the pairwise cohomology products vanish, there are elements $a_{12} \in A^{k_1+k_2-1}$ and $a_{23} \in A^{k_2+k_3-1}$ such that

$$da_{12} = a_1 a_2 \quad \text{and} \quad da_{23} = a_2 a_3.$$

Then one easily checks that

$$(-1)^{k_1+1} a_1 a_{23} + a_{12} a_3$$

is a cocycle in $A^{k_1+k_2+k_3-1}$. Its cohomology class is called a (triple) *Massey product* of α_1, α_2 and α_3, and denoted by $\langle \alpha_1, \alpha_2, \alpha_3 \rangle$.

More precisely, the Massey product $\langle \alpha_1, \alpha_2, \alpha_3 \rangle$ is the set of all elements in $H^{k_1+k_2+k_3-1}[A]$ obtained in this way. Since there are choices of a_{12} and a_{23} involved, the set $\langle \alpha_1, \alpha_2, \alpha_3 \rangle$ may consist of more than one element. In fact, a_{12} is defined up to addition of a cocycle in $A^{k_1+k_2-1}$, and a_{23} is defined up a cocycle in $A^{k_2+k_3-1}$. Therefore, any two elements in the set $\langle \alpha_1, \alpha_2, \alpha_3 \rangle$ differ by an element of the subset

$$\alpha_1 \cdot H^{k_2+k_3-1}[A] + \alpha_3 \cdot H^{k_1+k_2-1}[A] \subset H^{k_1+k_2+k_3-1}[A],$$

which is called the *indeterminacy* of the Massey product $\langle \alpha_1, \alpha_2, \alpha_3 \rangle$.

A Massey product $\langle \alpha_1, \alpha_2, \alpha_3 \rangle$ is called *trivial* (or *vanishing*) if it contains zero. Clearly, a Massey product $\langle \alpha_1, \alpha_2, \alpha_3 \rangle$ is trivial if and only if its image in the quotient algebra $H[A]/(\alpha_1, \alpha_3)$ is zero.

Proposition A.4.9. *Let $f\colon A \to B$ be a quasi-isomorphism of dg-algebras. Then all Massey products in $H[A]$ are trivial if and only they are all trivial in $H[B]$.*

Proof. Assume that all Massey products in $H[B]$ are trivial. Let $\langle \alpha_1, \alpha_2, \alpha_3 \rangle$ be a Massey product in $H[A]$. We define elements $a_1, a_2, a_3, a_{12}, a_{23} \in A$ as in Construction A.4.8, and set $b_i = f(a_i)$, $b_{ij} = f(a_{ij})$. Let β_i denote the cohomology class corresponding to the cocycle b_i. Since $\beta_1\beta_2 = \widetilde{f}(\alpha_1\alpha_2) = 0$ and $\beta_2\beta_3 = 0$, the Massey product $\langle \beta_1, \beta_2, \beta_3 \rangle$ is defined. By the assumption, $0 \in \langle \beta_1, \beta_2, \beta_3 \rangle$. This means that we may choose $b'_{12}, b'_{23} \in B$ in such a way that

$$b_1 b_2 = db'_{12}, \quad b_2 b_3 = db'_{23}, \quad \text{and} \quad (-1)^{k_1+1} b_1 b'_{23} + b'_{12} b_3 = db_{123}$$

for some $b_{123} \in B^{k_1+k_2+k_3-2}$. Since $db_{12} = b_1 b_2$, we have $d(b'_{12} - b_{12}) = 0$. Since $f\colon A \to B$ is a quasi-isomorphism, there is a cocycle $c_{12} \in A$ such that $f(c_{12}) = b'_{12} - b_{12}$, and similarly there is a cocycle $c_{23} \in A$ such that $f(c_{23}) = b'_{23} - b_{23}$. Set

$$a'_{12} = a_{12} + c_{12}, \quad a'_{23} = a_{23} + c_{23}.$$

Then $da'_{12} = da_{12} = a_1 a_2$ and similarly $da'_{23} = a_2 a_3$. Therefore, the cohomology class of $c = (-1)^{k_1+1} a_1 a'_{23} + a'_{12} a_3$ is a Massey product of $\alpha_1, \alpha_2, \alpha_3$. Then

$$f(c) = (-1)^{k_1+1} b_1 b'_{23} + b'_{12} b_3 = db_{123}.$$

Using again the fact that $f\colon A \to B$ is a quasi-isomorphism, we obtain that c is a coboundary, i.e. $c = da_{123}$ for some $a_{123} \in A$. Therefore, the Massey product $\langle \alpha_1, \alpha_2, \alpha_3 \rangle$ is trivial.

The fact that the triviality of Massey products in $H[A]$ implies their triviality in $H[B]$ is proved similarly (an exercise). □

Corollary A.4.10. *If a dg-algebra A is formal, then all Massey products in $H[A]$ are trivial.*

Proof. Apply Proposition A.4.9 to a zigzag $A \leftarrow \cdots \rightarrow H[A]$ of quasi-isomorphisms and use that fact that all Massey products for the dg-algebra $H[A]$ with zero differential are trivial. $\square$

Exercises.

A.4.11. Let $A \to B$ be a quasi-isomorphism of dg-algebras. Show that the triviality of Massey products in $H[A]$ implies their triviality in $H[B]$.

A.4.12. Find a nontrivial Massey product in the cohomology of the dg-algebra of Example A.4.7.

APPENDIX B

Algebraic topology

B.1. Homotopy and homology

Here we collect the basic constructions and facts, with almost no proofs. For the details the reader is referred to standard sources on algebraic topology, such as **[91]**, **[115]** or **[172]**. Unlike the other appendices, we do not include exercises here, as there would be too many of them.

All spaces here are topological spaces, and all maps are continuous. We denote by $\mathbf{k}$ a commutative ring with unit (usually $\mathbb{Z}$ or a field; in fact an abelian group is enough until we start considering the multiplication in cohomology).

Basic homotopy theory. Two maps $f, g\colon X \to Y$ of spaces are *homotopic* (denoted by $f \simeq g$) if there is a map $F\colon X \times \mathbb{I} \to Y$ (where $\mathbb{I} = [0,1]$ is the unit segment) such that $F|_{X\times 0} = f$ and $F|_{X\times 1} = g$. We denote the map $F|_{X\times t}\colon X \to Y$ by F_t, for $t \in \mathbb{I}$. The homotopy is an equivalence relation, and we denote by $[X, Y]$ the set of homotopy classes of maps from X to Y.

Two spaces X and Y are *homotopy equivalent* if there are maps $f\colon X \to Y$ and $g\colon Y \to X$ such that $g \circ f$ and $f \circ g$ are homotopic to the identity maps of X and Y respectively. The *homotopy type* of a space X is the class of spaces homotopy equivalent to X.

A space X is *contractible* if it is homotopy equivalent to a point.

A *pair* (X, A) of spaces consists of a space X and its subspace A. A *map of pairs* $f\colon (X, A) \to (Y, B)$ is a continuous map $f\colon X \to Y$ such that $f(A) \subset B$.

A *pointed space* (or *based space*) is a pair (X, pt) where pt is a point of X. We denote by X_+ the pointed space $(X \sqcup pt, pt)$, where $X \sqcup pt$ is X with a disjoint point added. A *map of pointed spaces* (or *pointed map*) is a map of pairs $(X, pt) \to (Y, pt)$. We denote a pointed space (X, pt) simply by X whenever the choice of the basepoint is clear or irrelevant. Given two pointed spaces (X, pt) and (Y, pt), their *wedge* (or *bouquet*) is defined as the pointed space $X \vee Y$ obtained by attaching X and Y at the basepoints. Then $X \vee Y$ is contained as a pointed subspace in the product $X \times Y$, and the quotient $X \wedge Y = (X \times Y)/(X \vee Y)$ is called the *smashed product* of X and Y.

For any pointed space (X, pt) the set of homotopy classes of pointed maps $(S^k, pt) \to (X, pt)$ (where S^k is a k-dimensional sphere, $k \geqslant 0$) is a group for $k > 0$, which is called the *kth homotopy group* of (X, pt) and denoted by $\pi_k(X, pt)$ or simply $\pi_k(X)$. We have that $\pi_0(X)$ is the set of path connected components of X. The group $\pi_1(X)$ is called the *fundamental group* of X. The groups $\pi_k(X)$ are abelian for $k > 1$. A map $f\colon X \to Y$ induces a homomorphism $f_*\colon \pi_k(X) \to \pi_k(Y)$ for each k, and the homomorphisms induced by homotopic maps are the same.

A *locally trivial fibration* (or a *fibre bundle*) is a quadruple (E, B, F, p) where E, B, F are spaces and p is a map $E \to B$ such that for any point $x \in B$ there is a neighbourhood $U \subset B$ and a homeomorphism $\varphi\colon p^{-1}(U) \xrightarrow{\cong} U \times F$ closing the commutative diagram

$$\begin{tikzcd}
p^{-1}(U) \arrow[rr, "\varphi"] \arrow[dr, "p"'] & & U \times F \arrow[dl] \\
& B &
\end{tikzcd}$$

The space E is called the *total space*, B the *base*, and F the *fibre* of the fibre bundle. The terms ‘locally trivial fibration’ and ‘fibre bundle’ are also often used for the map $p\colon E \to B$.

A *cell complex* (or a *CW-complex*) is a Hausdorff topological space X represented as a union $\bigcup e_i^q$ of pairwise nonintersecting subsets e_i^q, called *cells*, in such a way that for each cell e_i^q there is a map of a closed q-disk D^q to X (the *characteristic map* of e_i^q) whose restriction to the interior of D^q is a homeomorphism onto e_i^q. Furthermore, the following two conditions are assumed:

(C) the boundary $\bar{e}_i^q \setminus e_i^q$ of a cell is contained in a union of finitely many cells of dimensions $< q$;

(W) a subset $Y \subset X$ is closed if and only the intersection $Y \cap \bar{e}_i^q$ is closed for ever cell e_i^q (i.e. the topology of X is the weakest topology in which all characteristic maps are continuous).

The union of cells of X of dimension $\leqslant n$ is called the *nth skeleton* of X and denoted by $\mathrm{sk}^n X$ or by X^n. A cell complex X can be obtained from its 0th skeleton $\mathrm{sk}^0 X$ (which is a discrete set) by iterating the operation of *attaching a cell*: a space Z is obtained from Y by attaching an n-cell along a map $f\colon S^{n-1} \to Y$ if Z is the pushout of the form

$$\begin{tikzcd}
S^{n-1} \arrow[r, "f"] \arrow[d] & Y \arrow[d] \\
D^n \arrow[r] & Z
\end{tikzcd}$$

We use the notation $Z = Y \cup_f D^n$.

A *cell subcomplex* of a cell complex X is a closed subspace which is a union of cells of X. Each skeleton of X is a cell subcomplex.

A map $f\colon X \to Y$ between cell complexes is called a *cellular map* if $f(\mathrm{sk}^n X) \subset \mathrm{sk}^n Y$ for all n.

Theorem B.1.1 (Cellular approximation). *A map between cell complexes is homotopic to a cellular map.*

For any natural n and any group π (which is assumed to be abelian if $n > 1$) there exists a connected cell complex $K(\pi, n)$ such that $\pi_n\bigl(K(\pi, n)\bigr) \cong \pi$ and $\pi_k\bigl(K(\pi, n)\bigr) = 0$ for $k \neq n$. (A cell complex $K(\pi, n)$ can be constructed by taking the wedge of n-spheres corresponding to a set of generators of π and then killing the higher homotopy groups by attaching cells.) The space $K(\pi, n)$ is called the *Eilenberg–Mac Lane space* (corresponding to n and π), and it is unique up to homotopy equivalence. Examples of Eilenberg–Mac Lane spaces include the circle

$S^1 = K(\mathbb{Z},1)$, the infinite-dimensional real projective space $\mathbb{R}P^\infty = K(\mathbb{Z}_2,1)$ and the infinite-dimensional complex projective space $\mathbb{C}P^\infty = K(\mathbb{Z},2)$.

A space X which is homotopy equivalent to $K(\pi,1)$ for some π is called *aspherical.* Any surface (a closed 2-dimensional manifold) which not a 2-sphere or a real projective plane is aspherical.

A locally trivial fibration $p\colon E \to B$ satisfies the following *covering homotopy property* (*CHP* for short) with respect to maps of cell complexes Y: for any homotopy $F\colon Y\times\mathbb{I} \to B$ and any map $\widetilde{f}\colon Y \to E$ such that $p\circ\widetilde{f} = F_0$ there is a covering homotopy $\widetilde{F}\colon Y\times\mathbb{I} \to E$, satisfying $\widetilde{F}_0 = \widetilde{f}$ and $p\circ\widetilde{F} = F$. This is described by the commutative diagram

$$\begin{tikzcd}
Y \arrow[r, "\widetilde{f}"] \arrow[d, "i_0"'] & E \arrow[d, "p"] \\
Y\times\mathbb{I} \arrow[r, "F"] \arrow[ur, dashed, "\widetilde{F}"] & B
\end{tikzcd}$$

A map $p\colon E \to B$ satisfying CHP with respect to maps of cell complexes $Y \to B$ is called a *Serre fibration.* A map $p\colon E \to B$ satisfying CHP with respect to all maps $Y \to B$ is called a *Hurewitz fibration.* The difference between Serre and Hurewitz fibrations is not important for our constructions and will be ignored; we shall use the term *fibration* for both of them. A fibre bundle is a fibration, but not every fibration is a fibre bundle. All fibres $p^{-1}(b)$, $b\in B$, of a fibration are homotopy equivalent if B is connected.

Theorem B.1.2 (Exact sequence of fibration). *For a fibration $p\colon E\to B$ with fibre F there exists a long exact sequence*

$$\cdots \to \pi_k(F) \xrightarrow{i_*} \pi_k(E) \xrightarrow{p_*} \pi_k(B) \xrightarrow{\partial} \pi_{k-1}(F) \to \cdots$$

where the map i_* is induced by the inclusion of the fibre $i\colon F\to E$, the map p_* is induced by the projection $p\colon E\to B$, and ∂ is the connecting homomorphism.

The connecting homomorphism ∂ is defined as follows. Take an element $\gamma\in\pi_k(B)$ and choose a representative $g\colon S^k\to B$ in the homotopy class γ. By considering the composition $S^{k-1}\times\mathbb{I}\to S^k \xrightarrow{g} B$ (where the first map contracts the top and bottom bases of the cylinder to the north and south poles of the sphere), we may view the map g as a homotopy $F\colon S^{k-1}\times\mathbb{I}\to B$ of the trivial map $F_0\colon S^{k-1}\to pt$ with F_1 also being trivial. Using the CHP we lift F to a homotopy $\widetilde{F}\colon S^{k-1}\times\mathbb{I}\to E$ with $\widetilde{F}_0$ still trivial, but $\widetilde{F}_1\colon S^{k-1}\to E$ being trivial only after projecting onto B. The latter condition means that $\widetilde{F}_1$ is in fact a map $S^{k-1}\to F$, homotopy class of which we take for $\partial\gamma$.

The *path space* of a space X is the space PX of pointed maps $(\mathbb{I},0)\to(X,pt)$. The *loop space* ΩX is the space of pointed maps $f\colon(\mathbb{I},0)\to(X,pt)$ such that $f(1)=pt$. The map $p\colon PX\to X$ given by $p(f)=f(1)$ is a fibration, with fibre (homotopy equivalent to) ΩX.

Proposition B.1.3. *For any map $f\colon X\to Y$ there exist a homotopy equivalence $h\colon X\to\widetilde{X}$ and a fibration $p\colon\widetilde{X}\to Y$ such that $f=p\circ h$. Furthermore, this* *decomposition is functorial in the sense that a commutative diagram of maps*

$$\begin{tikzcd}
X \arrow[r] \arrow[d, "f"'] & X' \arrow[d, "f'"'] \\
Y \arrow[r] & Y'
\end{tikzcd}$$

induces a commutative diagram

$$\begin{tikzcd}
X \arrow[rr] \arrow[dr, "h"] \arrow[dd, "f"'] & & X' \arrow[dr, "h'"] \arrow[dd, "f'"'] & \\
& \widetilde{X} \arrow[rr] \arrow[dl, "p"] & & \widetilde{X'} \arrow[dl, "p'"] \\
Y \arrow[rr] & & Y' &
\end{tikzcd}$$

Proof. Let $\widetilde{X}$ be the set of pairs (x, g) consisting of a point $x \in X$ and a path $g\colon \mathbb{I} \to Y$ with $g(0) = f(x)$.This is described by the pullback diagram

$$\begin{tikzcd}
\widetilde{X} \arrow[r] \arrow[d] & Y^{\mathbb{I}} \arrow[d, "p_0"] \\
X \arrow[r, "f"] & Y
\end{tikzcd}$$

where $Y^{\mathbb{I}}$ is the space of all paths $g\colon \mathbb{I} \to Y$ and the map p_0 takes g to $g(0)$.

Then the homotopy equivalence $h\colon X \to \widetilde{X}$ is given by $h(x) = (x, c_{f(x)})$, where $c_{f(x)}\colon \mathbb{I} \to Y$ is the constant path $t \mapsto f(x)$, and the fibration $p\colon \widetilde{X} \to Y$ is given by $p(x, h) = h(1)$. The functoriality property follows by inspection. □

A space homotopy equivalent to the fibre of the fibration $p\colon \widetilde{X} \to Y$ from Proposition B.1.3 is referred to as the *homotopy fibre* of the map $f\colon X \to Y$, and denoted by $\operatorname{hofib} f$. The functoriality of the construction of $\widetilde{X}$ implies that the homotopy fibre is well-defined: for any other decomposition $f = p' \circ h'$ into a composition of a fibration p' and a homotopy equivalence h' the homotopy fibre of f is homotopy equivalent to the fibre of p'. The homotopy fibre of the inclusion $pt \to X$ is the loop space ΩX.

An inclusion $i\colon A \to X$ of a cell subcomplex in a cell complex X satisfies the following *homotopy extension property* (*HEP* for short): for any map $\widehat{f}\colon X \to Y$, a homotopy $F\colon A \times \mathbb{I} \to Y$ such that $F_0 = \widehat{f}|_A$ can be extended to a homotopy $\widehat{F}\colon X \times \mathbb{I} \to Y$ such that $\widehat{F}_0 = \widehat{f}$ and $\widehat{F}|_{A\times\mathbb{I}} = F$. This is described by the commutative diagram

$$\begin{tikzcd}
A \arrow[r, "F'"] \arrow[d, "i"'] & Y^{\mathbb{I}} \arrow[d, "p_0"] \\
X \arrow[r, "Y"] \arrow[ur, dashed, "\widehat{F}'"] & Y
\end{tikzcd}$$

where F' is the *adjoint* of F (i.e. $F'(a) = h$ where $h(t) = F(a, t) \in Y$), the map p_0 takes h to $h(0)$, and $\widehat{F}'$ is the adjoint of $\widehat{F}$.

A map $i\colon A \to X$ satisfying HEP is called a *cofibration*, and $X/i(A)$ is its *cofibre*. A pair (X, A) for which $i\colon A \to X$ is a cofibration is called a *Borsuk pair.*

The *cone* over a space X is the quotient space $\operatorname{cone} X = (X \times \mathbb{I})/(X \times 1)$. The *suspension* ΣX is the quotient $(\operatorname{cone} X)/(X \times 0)$. There are inclusions $X \hookrightarrow \operatorname{cone} X \hookrightarrow \Sigma X$ of closed subspaces, where first map is given by $x \mapsto (x, 0)$ and the second by $(x, t) \mapsto \bigl(x, (t + 1)/2\bigr)$ for $x \in X$, $t \in \mathbb{I}$. The inclusion $i\colon X \to \operatorname{cone} X$ is a cofibration, with cofibre ΣX.

PROPOSITION B.1.4. *For any map $f\colon X \to Y$ there exist a cofibration $i\colon X \to \widehat{Y}$ and a homotopy equivalence $h\colon \widehat{Y} \to Y$ such that $f = h \circ i$. This decomposition is functorial.*

PROOF. Let $\widehat{Y}$ be the quotient of $(X \times \mathbb{I}) \sqcup Y$ obtained by identifying $x \times 0 \in X \times \mathbb{I}$ with $f(x) \in Y$. This is described by the pushout diagram

$$\begin{tikzcd}
X \arrow[r, "f"] \arrow[d, "i_0"'] & Y \arrow[d] \\
X \times \mathbb{I} \arrow[r] & \widehat{Y}
\end{tikzcd}$$

where the map i_0 takes x to $x \times 0$.

The cofibration $i\colon X \to \widehat{Y}$ is given by $i(x) = x \times 1$, and the homotopy equivalence $h\colon \widehat{Y} \to Y$ is given by $h(x \times t) = f(x)$ and $h(y) = y$. □

The space $\widehat{Y}$ from Proposition B.1.4 is known as the *mapping cylinder* of the map $f\colon X \to Y$. The space $\widehat{Y}/i(X) = \widehat{Y}/(X \times 1)$ is known as the *mapping cone* of $f\colon X \to Y$. A space homotopy equivalent to the mapping cone of f is called the *homotopy cofibre* of the map $f\colon X \to Y$. The homotopy cofibre of the projection $X \to pt$ is the suspension ΣX.

Simplicial homology and cohomology. Let $\mathcal{K}$ be a simplicial complex on the set $[m] = \{1, \dots, m\}$. An *oriented q-simplex* σ of $\mathcal{K}$ is a k-simplex $I = \{i_1, \dots, i_{q+1}\} \in \mathcal{K}$ together with an equivalence class of total orderings of the set I, two orderings being equivalent if they differ by an even permutation. Denote by $[I]$ the oriented q-simplex with the equivalence class of orderings containing $i_1 < \dots < i_{q+1}$.

Define the *qth simplicial chain group (or module) $C_q(\mathcal{K}; \mathbf{k})$ with coefficients in* $\mathbf{k}$ as the free $\mathbf{k}$-module with basis consisting of oriented q-simplices of $\mathcal{K}$, modulo the relations $\sigma + \bar{\sigma} = 0$ whenever σ and $\bar{\sigma}$ are differently oriented q-simplices corresponding to the same simplex of $\mathcal{K}$. Therefore, $C_q(\mathcal{K}, \mathbf{k})$ is a free $\mathbf{k}$-module with rank equal to $f_q(\mathcal{K})$ (the number of q-simplices of $\mathcal{K}$) for $q \geqslant 0$, and $C_q(\mathcal{K}; \mathbf{k}) = 0$ for $q < 0$.

Define the *simplicial chain boundary homomorphisms*

$$\begin{aligned}
&\partial_q\colon C_q(\mathcal{K}; \mathbf{k}) \to C_{q-1}(\mathcal{K}; \mathbf{k}), \quad q \geqslant 1,\\
&\partial_q[i_1, \dots, i_{q+1}] = \sum_{j=1}^{q+1} (-1)^{j-1} [i_1, \dots, \widehat{i_j}, \dots, i_{q+1}],
\end{aligned}$$

where $\widehat{i_j}$ denotes that i_j is missing. It is easily checked that $\partial_q \partial_{q+1} = 0$, so that $C_*(\mathcal{K}; \mathbf{k}) = \{C_q(\mathcal{K}; \mathbf{k}), \partial_q\}$ is a chain complex, called the *simplicial chain complex* of $\mathcal{K}$ (with coefficients in $\mathbf{k}$).

The *qth simplicial homology group of $\mathcal{K}$ with coefficients in* $\mathbf{k}$, denoted by $H_q(\mathcal{K};\mathbf{k})$, is defined as the qth homology group of the simplicial chain complex $C_*(\mathcal{K};\mathbf{k})$.

The *Euler characteristic* $\chi(\mathcal{K})$ of $\mathcal{K}$ is defined by

$$\chi(\mathcal{K})=\sum_{q\geqslant 0}(-1)^q\operatorname{rank}H_q(\mathcal{K};\mathbf{k}).$$

It is also given by

$$\chi(\mathcal{K})=f_0-f_1+f_2-\cdots,$$

where f_i is the number of i-simplices of $\mathcal{K}$, and therefore $\chi(\mathcal{K})$ is independent of $\mathbf{k}$.

The *augmented simplicial chain complex* $\widetilde{C}_*(\mathcal{K};\mathbf{k})$ is obtained by taking into account the empty simplex $\varnothing\in\mathcal{K}$. That is, $\widetilde{C}_q(\mathcal{K};\mathbf{k})=C_q(\mathcal{K};\mathbf{k})$ for $q\neq -1$ and $\widetilde{C}_{-1}(\mathcal{K};\mathbf{k})\cong\mathbf{k}$ is a free $\mathbf{k}$-module with basis $[\varnothing]$, so that $\widetilde{C}_*(\mathcal{K};\mathbf{k})$ is written as

$$\cdots\to C_q(\mathcal{K};\mathbf{k})\xrightarrow{\partial_q}\cdots\longrightarrow C_1(\mathcal{K};\mathbf{k})\xrightarrow{\partial_1}C_0(\mathcal{K};\mathbf{k})\xrightarrow{\varepsilon}\widetilde{C}_{-1}(\mathcal{K};\mathbf{k})\to 0,$$

where $\varepsilon=\partial_0$ is the *augmentation* taking each vertex $[i]$ to $[\varnothing]$.

The qth homology group of $\widetilde{C}_*(\mathcal{K};\mathbf{k})$ is called the *qth reduced simplicial homology group of $\mathcal{K}$ with coefficients in* $\mathbf{k}$ and is denoted by $\widetilde{H}_q(\mathcal{K};\mathbf{k})$. We have that $H_q(\mathcal{K};\mathbf{k})=\widetilde{H}_q(\mathcal{K};\mathbf{k})$ for $q\geqslant 1$, and $H_0(\mathcal{K};\mathbf{k})\cong\widetilde{H}_0(\mathcal{K};\mathbf{k})\oplus\mathbf{k}$ unless $\mathcal{K}$ consists of $\varnothing$ only, in which case $\widetilde{H}_{-1}(\varnothing;\mathbf{k})\cong\mathbf{k}$.

The *simplicial cochain complex* of $\mathcal{K}$ with coefficients in $\mathbf{k}$ is defined to be

$$C^*(\mathcal{K};\mathbf{k})=\operatorname{Hom}_{\mathbf{k}}\bigl(C_*(\mathcal{K};\mathbf{k}),\mathbf{k}\bigr).$$

In explicit terms, $C^*(\mathcal{K};\mathbf{k})=\{C^q(\mathcal{K};\mathbf{k}),d_q\}$. Here $C^q(\mathcal{K};\mathbf{k})$ is the *qth simplicial cochain group (or module) of $\mathcal{K}$ with coefficients in* $\mathbf{k}$; it is a free $\mathbf{k}$-module with basis consisting of simplicial cochains α_I corresponding to q-simplices $I\in\mathcal{K}$; the cochain α_I takes value 1 on the oriented simplex $[I]$ and vanishes on all other oriented simplices. The value of the *cochain differential*

$$d_q=\partial^*_{q+1}\colon C^q(\mathcal{K};\mathbf{k})\to C^{q+1}(\mathcal{K};\mathbf{k})$$

on the basis elements is given by

$$d\alpha_I=\sum_{j\in[m]\setminus I,\,j\cup I\in\mathcal{K}}\varepsilon(j,j\cup I)\alpha_{j\cup I},$$

where the sign is given by $\varepsilon(j,j\cup I)=(-1)^{r-1}$ if j is the rth element of the set $j\cup I\subset[m]$, written in increasing order.

The *qth simplicial cohomology group of $\mathcal{K}$ with coefficients in* $\mathbf{k}$, denoted by $H^q(\mathcal{K};\mathbf{k})$, is defined as the qth cohomology group of the simplicial cochain complex $C^*(\mathcal{K};\mathbf{k})$.

The *qth reduced simplicial cohomology group of $\mathcal{K}$ with coefficients in* $\mathbf{k}$, denoted by $\widetilde{H}^q(\mathcal{K};\mathbf{k})$, is defined as the qth cohomology group of the cochain complex

$$0\to\mathbf{k}\xrightarrow{d_{-1}}C^0(\mathcal{K};\mathbf{k})\xrightarrow{d_0}C^1(\mathcal{K};\mathbf{k})\xrightarrow{d_1}\cdots\longrightarrow C^q(\mathcal{K};\mathbf{k})\xrightarrow{d_q}\cdots$$

obtained by applying the functor Hom to the augmented chain complex $\widetilde{C}_*(\mathcal{K};\mathbf{k})$. The map d_{-1} takes $1=\alpha_\varnothing$ to the sum of $\alpha_{\{i\}}$ corresponding to all vertices $\{i\}\in\mathcal{K}$. We have that $H^q(\mathcal{K};\mathbf{k})=\widetilde{H}^q(\mathcal{K};\mathbf{k})$ for $q\geqslant 1$, and $H^0(\mathcal{K};\mathbf{k})\cong\widetilde{H}^0(\mathcal{K};\mathbf{k})\oplus\mathbf{k}$ unless $\mathcal{K}$ consists of $\varnothing$ only, in which case $\widetilde{H}^{-1}(\varnothing;\mathbf{k})\cong\mathbf{k}$.

The first nontrivial cohomology group $H^q(X;\mathbb{Z})$ with $q>0$ is torsion-free.

Singular homology and cohomology. Let X be a topological space. A *singular q-simplex* of X is a continuous map $f\colon \Delta^q \to X$. The *qth singular chain group $C_q(X;\mathbf{k})$ of X with coefficients in* $\mathbf{k}$ is the free $\mathbf{k}$-module generated by all singular q-simplices.

For each $i = 1,\dots,q+1$ there is a linear map $\varphi_q^i\colon \Delta^{q-1} \to \Delta^q$ which sends Δ^{q-1} to the face of Δ^q opposite its ith vertex, preserving the order of vertices. The *ith face* of a singular simplex $f\colon \Delta^q \to X$, denoted by $f^{(i)}$ is defined to be the singular $(q-1)$-simplex given by the composition

$$f^{(i)} = f \circ \varphi_q^i\colon \Delta^{q-1} \to \Delta^q \to X.$$

The *singular chain boundary homomorphisms* are given by

$$\partial_q\colon C_q(X;\mathbf{k}) \to C_{q-1}(X;\mathbf{k}), \quad q \geqslant 1,$$
$$\partial_q f = \sum_{i=1}^{q+1} (-1)^{i-1} f^{(i)}.$$

Then $\partial_q\partial_{q+1} = 0$, so that $C_*(X;\mathbf{k}) = \{C_q(X;\mathbf{k}), \partial_q\}$ is a chain complex, called the *singular chain complex* of X (with coefficients in $\mathbf{k}$).

The *qth singular homology group of X with coefficients in* $\mathbf{k}$, denoted by $H_q(X;\mathbf{k})$, is the qth homology group of the singular chain complex $C_*(X;\mathbf{k})$.

Assume that a space X has only finite number of nontrivial homology groups (with $\mathbb{Z}$ coefficients), and each of these groups has finite rank. Then the *Euler characteristic* $\chi(X)$ of X is defined by

$$\chi(X) = \sum_{q \geqslant 0} (-1)^q \operatorname{rank} H_q(X;\mathbb{Z}).$$

Proposition B.1.5. *The group $H_0(X;\mathbf{k})$ is a free $\mathbf{k}$-module of rank equal to the number of path connected components of X.*

The *augmented singular chain complex* $\widetilde{C}_*(X;\mathbf{k})$ is defined by $\widetilde{C}_q(X;\mathbf{k}) = C_q(X;\mathbf{k})$ for $q \neq -1$ and $\widetilde{C}_{-1}(X;\mathbf{k}) = \mathbf{k}$; the *augmentation* $\varepsilon = \partial_0\colon C_0(X;\mathbf{k}) \to \mathbf{k}$ is given by $\varepsilon(f) = 1$ for all singular 0-simplices f.

The qth homology group of $\widetilde{C}_*(X;\mathbf{k})$ is called the *qth reduced singular homology group of $\mathcal{K}$ with coefficients in* $\mathbf{k}$ and is denoted by $\widetilde{H}_q(X;\mathbf{k})$. We have $H_q(X;\mathbf{k}) = \widetilde{H}_q(X;\mathbf{k})$ for $q \geqslant 1$, and $H_0(X;\mathbf{k}) \cong \widetilde{H}_0(X;\mathbf{k}) \oplus \mathbf{k}$ if X is nonempty.

A nonempty space X is *acyclic* if $\widetilde{H}_q(X;\mathbb{Z}) = 0$ for all q.

We shall drop the coefficient group $\mathbf{k}$ in the notation of chains and homology occasionally.

For any simplicial complex $\mathcal{K}$, there is an obvious inclusion $i\colon C_q(\mathcal{K}) \to C_q(|\mathcal{K}|)$ of the simplicial chain groups into the singular chain groups of the geometric realisation $\mathcal{K}$, defining an inclusion of chain complexes.

Theorem B.1.6. *The map $i\colon C_q(\mathcal{K}) \to C_q(|\mathcal{K}|)$ induces an isomorphism $H_q(\mathcal{K}) \xrightarrow{\cong} H_q(|\mathcal{K}|)$ between the simplicial homology groups of $\mathcal{K}$ and the singular homology groups of $|\mathcal{K}|$.*

If (X,A) is pair of spaces, then $C_*(A)$ is a chain subcomplex in $C_*(X)$ (that is, $C_q(A)$ is a submodule of $C_q(X)$ and $\partial_q C_q(A) \subset C_{q-1}(A)$). The quotient complex

$$C_*(X,A) = \bigl\{C_q(X,A) = C_q(X)/C_q(A),\ \ \bar\partial_q\colon C_q(X,A) \to C_{q-1}(X,A)\bigr\}$$

is called the *singular chain complex of pair* (X, A). Its qth homology group $H_q(X, A)$ is called the *qth singular homology group of* (X, A), or *qth relative singular homology group of X modulo A*. Note that $H_q(X) = H_q(X, \varnothing)$ and $\widetilde{H}_q(X) = H_q(X, pt)$.

The *singular cochain complex* $C^*(X; \mathbf{k}) = \{C^q(X; \mathbf{k}), d_q\}$ is defined to be

$$C^*(X; \mathbf{k}) = \mathrm{Hom}_{\mathbf{k}}\bigl(C_*(X; \mathbf{k}), \mathbf{k}\bigr).$$

Its qth cohomology group is called the *qth singular cohomology group of X* and denoted by $H^q(X; \mathbf{k})$. The *reduced singular cohomology groups* $\widetilde{H}^q(X; \mathbf{k})$ and the *relative cohomology groups* $H^q(X, A)$ are defined similarly.

The ranks of the groups $H_q(X; \mathbb{Z})$ and $H^q(X; \mathbb{Z})$ coincide, and the number $b^q(X) = \operatorname{rank} H^q(X; \mathbb{Z})$ is called the *qth Betti number* of X. The Betti numbers with coefficients in a field $\mathbf{k}$ are defined similarly.

The (co)homology groups have the following fundamental properties.

Theorem B.1.7 (Functoriality and homotopy invariance). *A map* $f\colon X \to Y$ *induces homomorphisms* $f_*\colon H_q(X; \mathbf{k}) \to H_q(Y; \mathbf{k})$ *and* $f^*\colon H^q(Y; \mathbf{k}) \to H^q(X; \mathbf{k})$. *If two maps* $f, g\colon X \to Y$ *are homotopic, then* $f_* = g_*$ *and* $f^* = g^*$.

We omit the coefficient group $\mathbf{k}$ in the notation for the rest of this subsection.

Theorem B.1.8 (Exact sequences of pairs). *For any pair* (X, A) *there are long exact sequences*

$$\cdots \to H_q(A) \xrightarrow{i_*} H_q(X) \xrightarrow{j_*} H_q(X, A) \xrightarrow{\partial} H_{q-1}(A) \to \cdots,$$

$$\cdots \to H^q(X, A) \xrightarrow{j^*} H^q(X) \xrightarrow{i^*} H^q(A) \xrightarrow{d} H^{q+1}(X, A) \to \cdots$$

Furthermore, if the inclusion $A \to X$ *is a cofibration (e.g., if it is an inclusion of a cell subcomplex), then the quotient projection* $(X, A) \to (X/A, pt)$ *induces isomorphisms*

$$H_q(X, A) \cong H_q(X/A, pt) = \widetilde{H}_q(X/A), \quad H^q(X, A) \cong H^q(X/A, pt) = \widetilde{H}^q(X/A).$$

In the homology exact sequence of pair, the map i_* is induced by the inclusion $A \to X$, the map j_* is induced by the map of pairs $(X, \varnothing) \to (X, A)$, and the *connecting homomorphism* ∂ is the homology homomorphism corresponding the boundary homomorphism sending a relative cocycle $c \in C_q(X, A)$ to $\partial c \in C_q(A)$. The maps in the cohomology exact sequence are defined dually.

Let (X, A) be a pair and $B \subset A$ be a subspace. The inclusion of pairs $(X \backslash B, A \backslash B) \to (X, A)$ induces maps

$$H_q(X \backslash B, A \backslash B) \to H_q(X, A),$$

which are referred to as the *excision homomorphisms*.

Theorem B.1.9 (Excision). *If the closure of B is contained in the interior of A, then the excision homomorphisms* $H_q(X \backslash B, A \backslash B) \to H_q(X, A)$ *are isomorphisms.*

Theorem B.1.10 (Mayer–Vietoris exact sequences). *Let X be a space,* $A \subset X$, $B \subset X$ *and* $A \cup B = X$. *Assume that the excision homomorphisms*

$$H_q(B, A \cap B) \to H_q(X, A) \quad \text{and} \quad H_q(A, A \cap B) \to H_q(X, B)$$

are isomorphisms for all q. Then there are exact sequences

$$\cdots \to H_q(A\cap B) \xrightarrow{\beta} H_q(A)\oplus H_q(B) \xrightarrow{\alpha} H_q(X) \xrightarrow{\delta} H_{q-1}(A\cap B) \to \cdots,$$

$$\cdots \to H^q(X) \xrightarrow{\alpha} H^q(A)\oplus H^q(B) \xrightarrow{\beta} H^q(A\cap B) \xrightarrow{\delta} H^{q+1}(X) \to \cdots.$$

In the homology Mayer–Vietoris sequence, the map β is the difference of the homomorphism induced by the inclusions $A\cap B\to A$ and $A\cap B\to B$, the map α is the sum of the homomorphism induced by the inclusions $A\to X$ and $B\to X$, and the connecting homomorphism δ is the composite

$$H_q(X) \xrightarrow{j_*} H_q(X,A) = H_q(B, A\cap B) \xrightarrow{\partial} H_{q-1}(A\cap B).$$

The first key calculation of homology groups follows directly from the general properties above:

Proposition B.1.11. *The n-disk D^n is acyclic, and the reduced homology of an n-sphere S^n is given by $\widetilde{H}_n(S^n;\mathbf{k})\cong\mathbf{k}$ and $\widetilde{H}_i(S^n;\mathbf{k})=0$ for $i\neq n$.*

Here is another direct corollary:

Theorem B.1.12 (Suspension isomorphism). *For any space X and any $q>0$ there are isomorphisms*

$$\widetilde{H}_q(\Sigma X)\cong\widetilde{H}_{q-1}(X) \quad \textit{and} \quad \widetilde{H}^q(\Sigma X)\cong\widetilde{H}^{q-1}(X).$$

A closed connected topological n-dimensional manifold X is *orientable* over $\mathbf{k}$ if $H_n(X;\mathbf{k})\cong\mathbf{k}$. Every compact connected manifold is oriented over $\mathbb{Z}_2$ or any field of characteristic two.

Theorem B.1.13 (Poincaré duality). *If a closed connected n-manifold is orientable over $\mathbf{k}$, then $H_q(X;\mathbf{k})\cong H^{n-q}(X;\mathbf{k})$.*

Cellular homology, and cohomology multiplication. Let X be a cell complex with skeletons $\operatorname{sk}^n X = X^n$ for $n=0,1,2,\dots$

We start with the case $\mathbf{k}=\mathbb{Z}$, and omit this coefficient group in the notation. The group $\mathcal{C}_q(X)=H_q(X^q,X^{q-1})$ is called the *qth cellular chain group*. This is a free abelian group with basis of q-dimensional cells of X, and therefore a cellular chain may be viewed as an integral combination of cells of X.

The *cellular boundary homomorphism* $\partial_q\colon\mathcal{C}_q(X)\to\mathcal{C}_{q-1}(X)$ is defined as the composition

$$\partial_q\colon \mathcal{C}_q(X)=H_q(X^q,X^{q-1})\to H_{q-1}(X^{q-1})\to H_{q-1}(X^{q-1},X^{q-2})=\mathcal{C}_{q-1}(X)$$

of homomorphisms from the homology exact sequences of pairs (X^q,X^{q-1}) and (X^{q-1},X^{q-2}). The resulting chain complex

$$\cdots\to\mathcal{C}_q(X)\xrightarrow{\partial_q}\cdots\longrightarrow\mathcal{C}_1(X)\xrightarrow{\partial_1}\mathcal{C}_0(X)\to 0$$

is called the *cellular chain complex* of X. Its qth homology group, which we denote by $\mathcal{H}_q(X)$ for a moment, is called the *qth cellular homology group* of X.

Theorem B.1.14. *For any cell complex X, there is a canonical isomorphism $\mathcal{H}_q(X)\cong H_q(X)$ between the cellular and singular homology groups.*

The cellular homology groups $\mathcal{H}_q(X;\mathbf{k})$ with coefficients in $\mathbf{k}$ and the cohomology groups $\mathcal{H}^q(X;\mathbf{k})$ are defined similarly; they a canonically isomorphic to the appropriate singular homology and cohomology groups.

We shall therefore not distinguish between the singular and cellular homology and cohomology groups of cell complexes.

The product $X_1 \times X_2$ of cell complexes is a cell complex with cells of the form $e_1 \times e_2$, where e_1 is a cell of X_1 and e_2 is a cell of X_2.

Given two cellular cochains $c_1 \in \mathcal{C}^{q_1}(X)$ and $c_2 \in \mathcal{C}^{q_2}(X)$, define their *product* as the cochain $c_1 \times c_2 \in \mathcal{C}^{q_1+q_2}(X \times X)$ whose value on a cell $e_1 \times e_2$ of $X \times X$ is given by $(-1)^{q_1q_2}c_1(e_1)c_2(e_2)$. This product satisfies the identity

$$\delta(c_1 \times c_2) = \delta c_1 \times c_2 + (-1)^{q_1} c_1 \times \delta c_2,$$

where δ is the cellular cochain differential, and therefore defines a map of cochain complexes

$$\mathcal{C}^*(X) \otimes \mathcal{C}^*(X) \to \mathcal{C}^*(X \times X)$$

and induces a cohomology map

$$\times\colon H^{q_1}(X) \otimes H^{q_2}(X) \to H^{q_1+q_2}(X \times X),$$

which is called the (cohomology) $\times$-*product*.

The composition of the $\times$-product with the cohomology map induced by the *diagonal map* $\varDelta\colon X \to X \times X$, $\varDelta(x) = (x,x)$, defines a product

$$\smile\colon H^{q_1}(X) \otimes H^{q_2}(X) \xrightarrow{\times} H^{q_1+q_2}(X \times X) \xrightarrow{\varDelta^*} H^{q_1+q_2}(X),$$

which is called the $\smile$-*product*, or *cohomology product*. It turns $H^*(X) = \bigoplus_{q\geqslant 0} H^q(X)$ into an associative and graded commutative ring. This ring structure on $H^*(X)$ is a homotopy invariant of X; it does not depend on the cell complex structure. It is also functorial, in the sense that a map $f\colon X \to Y$ induces a ring homomorphism $f^*\colon H^*(Y) \to H^*(X)$.

There is also a relative version of the cohomology product, given by the map

$$H^{q_1}(X;A) \otimes H^{q_2}(X,B) \longrightarrow H^{q_1+q_2}(X, A \cup B).$$

B.2. Elements of rational homotopy theory

We only review some basis notions and results here. For a detailed account of this much elaborated theory we refer to [**30**], [**140**] and [**89**].

Definition B.2.1. A map $f\colon X \to Y$ between spaces is called a *rational equivalence* if it induces isomorphisms in all rational homotopy groups, that is, $f_*\colon \pi_i(X) \otimes \mathbb{Q} \to \pi_i(Y) \otimes \mathbb{Q}$ is an isomorphism for all i.

The *rational homotopy type* of X is its equivalence class in the equivalence relation generated by rational equivalences.

Proposition B.2.2. *If both X and Y are simply connected, then $f\colon X \to Y$ is a rational equivalence if and only if $f_*\colon H_i(X;\mathbb{Q}) \to H_i(Y;\mathbb{Q})$ (or equivalently, $f^*\colon H^i(Y;\mathbb{Q}) \to H^i(X;\mathbb{Q})$) is an isomorphism for each i.*

A space X is *nilpotent* if its fundamental group $\pi_1(X)$ is nilpotent, and $\pi_1(X)$ acts nilpotently on higher homotopy groups $\pi_n(X)$. A simply connected space is nilpotent.

Rational homotopy theory, whose foundation was laid in the work of Quillen [**191**] and Sullivan [**213**], translates the study of the rational homotopy type of nilpotent spaces X into the algebraic language of dg-algebras and minimal models (see Section A.4). This translation is made via Sullivan's *algebra of piecewise polynomial differential forms* $A_{PL}(X)$, whose properties we briefly discuss below (for the precise definition we refer to the sources mentioned above).

The most basic dg-algebra model of a space X is its singular cochains $C^*(X;\mathbf{k})$. However, this dg-algebra is non-commutative, and therefore is difficult to handle. If X is a smooth manifold, and $\mathbf{k} = \mathbb{R}$, then we may consider the dg-algebra $\Omega^*(X)$ of de Rham differential forms instead of $C^*(X;\mathbb{R})$. It provides a functorial (with respect to smooth maps) and commutative dg-algebra model for X, with the same cohomology $H^*(X;\mathbb{R})$. It is therefore natural to ask whether a functorial commutative dg-algebra model exists for arbitrary cell complexes X over a field $\mathbf{k}$ of zero characteristic (over finite fields there are secondary cohomological operations obstructing such a construction). A first construction of a commutative dg-algebra model which worked for simplicial complexes was suggested by Thom in the end of the 1950s. Later, in mid-1970s, Sullivan provided a functorial construction of a dg-algebra model $A_{PL}(X)$ whose cohomology is $H^*(X;\mathbb{Q})$, using similar ideas as those of Thom (a combinatorial version of differential forms).

The algebra $A_{PL}(X)$ has the following two important properties.

THEOREM B.2.3.

(a) *$A_{PL}(X)$ is weakly equivalent to $C^*(X;\mathbb{Q})$ via a short zigzag of the form*

$$A_{PL}(X) \to D(X) \leftarrow C^*(X;\mathbb{Q}),$$

where $D(X)$ is another naturally defined dg-algebra;

(b) *there is a natural map of cochain complexes $A_{PL}(X) \to C^*(X;\mathbb{Q})$, the 'Stokes map', which induces an isomorphism in cohomology.*

(c) *if X is a smooth manifold, then the dg-algebra $\Omega^*(X)$ of de Rham forms is weakly equivalent to $A_{PL}(X) \otimes_{\mathbb{Q}} \mathbb{R}$.*

The proof of (a) can be found in [**140**, §III.3] or in [**89**, Corollary 10.10]. For (b), see [**30**, §2], and for (c), see [**140**, Theorem V.2] or [**89**, Theorem 11.4].

DEFINITION B.2.4. Given a cell complex X, we refer to a minimal model of the commutative dg-algebra $A_{PL}(X)$ as a *minimal model* of X, and denote it by M_X.

In the case of nilpotent spaces, the rational homotopy type of X is fully determined by the commutative dg-algebra $A_{PL}(X)$ or its minimal model M_X. More precisely, there is the following fundamental result.

THEOREM B.2.5. *There is a bijective correspondence between the set of rational homotopy types of nilpotent spaces and the set of classes of isomorphic minimal dg-algebras over $\mathbb{Q}$. Under this correspondence, there is a natural isomorphism*

$$\mathrm{Hom}\big(\pi_i(X),\mathbb{Q}\big) \cong M_X^i/(M_X^+ \cdot M_X^+),$$

i.e. the rank of the ith rational homotopy group of X equals the number of generators of degree i in the minimal model of X.

For the proof, see [**140**, Theorem IV.8].

DEFINITION B.2.6. A space X is called *formal* if $C^*(X;\mathbb{Q})$ is a formal dg-algebra (equivalently, if $A_{PL}(X)$ is a formal commutative dg-algebra).

If X in nilpotent, then it is formal if and only if there is a quasi-isomorphism $M_X \to H^*(X;\mathbb{Q})$. If X is a smooth manifold, then it is formal if and only if the de Rham algebra $\Omega^*(X)$ is formal; in this case we can define M_X as a minimal model of $\Omega^*(X)$ instead of $A_{PL}(X)$.

Example B.2.7.

1. Let $X = S^{2n+1}$ be an odd sphere, $n \geqslant 1$. Then

$$M_X = \Lambda[x], \quad \deg x = 2n+1, \quad dx = 0.$$

There is a quasi-isomorphism $M_X \to \Omega^*(X)$ sending x to the volume form of S^{2n+1}.

2. Let $X = S^{2n}$, $n \geqslant 1$. Then

$$M_X = \Lambda[y] \otimes \mathbb{R}[x], \quad \deg x = 2n,\ \deg y = 4n-1, \quad dx = 0,\ dy = x^2.$$

The quasi-isomorphism $M_X \to \Omega^*(X)$ sends x to the volume form of S^{2n} and y to zero.

3. Let $X = \mathbb{C}P^n$, $n \geqslant 1$. Then

$$M_X = \Lambda[y] \otimes \mathbb{R}[x], \quad \deg x = 2,\ \deg y = 2n+1, \quad dx = 0,\ dy = x^{n+1}.$$

There is a quasi-isomorphism $M_X \to \Omega^*(X)$ sending x to the Fubini–Study 2-form $\omega = \frac{i}{2\pi}\partial\overline{\partial}\log|\boldsymbol{z}|^2 \in \Omega^2(\mathbb{C}P^n)$ and y to zero.

4. Let $X = \mathbb{C}P^\infty$. Then

$$M_X = \mathbb{Q}[x] = H^*(X;\mathbb{Q}), \quad \deg x = 2, \quad dx = 0.$$

In all examples above the space X is formal (an exercise). Here is an example of a nonformal manifold.

Example B.2.8. Let G be the *Heisenberg group* consisting of matrices

$$\begin{pmatrix} 1 & x & z \\ 0 & 1 & y \\ 0 & 0 & 1 \end{pmatrix}, \quad x, y, z \in \mathbb{R},$$

and let $\Gamma = G_{\mathbb{Z}}$ be the subgroup consisting of matrices with $x, y, z \in \mathbb{Z}$. The quotient manifold $X = G/\Gamma$ is the classifying space for the nilpotent group $\Gamma = \pi_1(X)$, i.e. $X = K(\Gamma, 1)$. The minimal model M_X is generated by the left-invariant forms

$$\omega_1 = dx, \quad \omega_2 = dy, \quad \omega_3 = xdy - dz.$$

This dg-algebra is isomorphic to the dg-algebra of example A.4.7. Therefore, the manifold X is not formal.

Exercises.

B.2.9. Show that all spaces of Example B.2.7 are formal.

B.3. Group actions and equivariant cohomology

Here we present the background material related to the notion of equivariant cohomology. The basic reference is the monograph [**31**] by Bredon, where the details of the constructions and proofs can be found.

Let X be a Hausdorff topological space and G a topological group. It is said that G *acts* on X if for any element $g \in G$ there is a homeomorphism $\varphi_g \colon X \to X$, and the assignment $g \mapsto \varphi_g$ respects the algebraic and topological structure. In more precise terms, a (left) *action* of G on X is given by a continuous map

$$G \times X \to X, \quad (g, x) \mapsto gx$$

such that $g(hx) = (gh)x$ for any $g, h \in G,\ x \in X$, and $ex = x$, where e is the unit of G. The space X is called a (left) *G-space*. Right actions and right G-spaces are defined similarly. In the case when G is an abelian group, the notions of left and right action coincide.

A group action is *effective* if the set

$$\{g \in G\colon gx = x \text{ for all } x\}$$

consists of a single element $e \in G$. A continuous map $f\colon X \to Y$ of G-spaces is *equivariant* if it commutes with the group actions, i.e. $f(gx) = g(f(x))$ for all $g \in G$ and $x \in X$. A map f is *weakly equivariant* if there is an automorphism $\psi\colon G \to G$ such that $f(gx) = \psi(g)(f(x))$ for all $g \in G$ and $x \in X$. A weakly equivariant map corresponding to an automorphism ψ is also called *ψ-equivariant*.

Let $x \in X$. The set

$$G_x = \{g \in G\colon gx = x\}$$

of elements of G fixing the point x is a closed subgroup in G, called the *stationary subgroup*, or the *stabiliser* of x. A G-action is *free* if G_x is the trivial (unit) subgroup for all $x \in X$. The subspace

$$Gx = \{gx \in X : g \in G\} \subset X$$

is called the *orbit* of x with respect to the action of G. The set of all orbits is denoted by X/G, and we have a canonical projection $\pi\colon X \to X/G$. The set X/G is endowed with the standard quotient topology (a set $U \subset X/G$ is open if and only if $\pi^{-1}(U)$ is open), and is referred to as the *orbit space*, or the *quotient space*, and denoted by X/G. If G is a compact group, then the quotient X/G is Hausdorff, and the projection $\pi\colon X \to X/G$ is a closed and proper map (i.e. the image of a closed subset is closed, and the preimage of a compact subset is compact).

A point $x \in X$ is *fixed* if $Gx = x$, i.e. $G_x = G$. The set of all fixed points of a G-space X will be denoted by X^G.

A *principal G-bundle* is a locally trivial bundle $p\colon X \to B$ with fibre G and structure group G, such that G acts fibrewise on X by right translations (that is, an element $g \in G$ maps a point g' in the fibre to $g'g$). The standard action of G on itself by left translations commutes with the right translations. This implies that for any principal G-bundle $p\colon X \to B$ there is a canonical free left G-action on X covering the trivial action on B. The map $p\colon X \to B$ induces a homeomorphism $X/G \cong B$, and therefore p can be regarded as the projection onto the orbit space of a free G-action. If the group G is compact, under some mild topological assumptions on X the converse is also true, namely, if G acts on X freely, then X is the total space of a principal G-bundle (see [**31**, Chapter II]). Therefore, in this case the notions of a principal G-bundle and a free G-action are equivalent.

Now let G be a compact Lie group. Then there exists a principal G-bundle $EG \to BG$ whose total space EG is contractible. This bundle has the following universality property. Let $E \to B$ be another principal G-bundle. Then there is a unique up to homotopy map $f\colon B \to BG$ such that the pullback of the bundle $EG \to BG$ along f is the bundle $E \to B$. The space EG is referred to as the *universal G-space*, and the space BG is the *classifying space* for free G-actions (or simply the classifying space for G).

Let X be a left G-space. There is the following free left G-action on $EG \times X$:

$$g(e, x) = (eg^{-1}, gx), \qquad g \in G,\ e \in EG,\ x \in X.$$

The orbit space for this action is denoted by $EG \times_G X$ (we shall also use the notation $B_G X$) and is called the *Borel construction*, or the *homotopy quotient* of X by G. (The latter term is used since the free G-space $EG \times X$ is homotopy equivalent to the G-space X.) There are two canonical projections

$$\begin{array}{c} EG \times X \to EG \\ (e,x) \mapsto e \end{array} \quad \text{and} \quad \begin{array}{c} EG \times X \to X \\ (e,x) \mapsto x. \end{array}$$

After taking quotient by the G-action, the first projection gives rise to a bundle $EG \times_G X \to BG$ with fibre X and structure group G, called the bundle *associated* with the G-space X. The second projection induces a map $EG \times_G X \to X/G$. There is also a principal G-bundle $EG \times X \to EG \times_G X$.

For each fixed point $x \in X$ there is a section $BG \to EG \times_G X$ of the bundle $EG \times_G X \to BG$, and therefore we have an inclusion

$$BG \times X^G \to EG \times_G X.$$

The *equivariant cohomology* of X with coefficients in a ring $\mathbf{k}$ is defined by

$$H^*_G(X;\mathbf{k}) = H^*(EG \times_G X;\mathbf{k}).$$

Hence, $H^*_G(pt;\mathbf{k}) = H^*(BG;\mathbf{k})$, and the projection $EG \times_G X \to BG$ turns $H^*_G(X;\mathbf{k})$ into a $H^*_G(pt;\mathbf{k})$-module.

For a pair of G-spaces (X,A) (where the inclusion $A \subset X$ is an equivariant map), there is a long exact sequence

$$\cdots \to H^q_G(X,A) \xrightarrow{j^*} H^q_G(X) \xrightarrow{i^*} H^q_G(A) \xrightarrow{d} H^{q+1}_G(X,A) \to \cdots$$

Its existence follows from the exact sequence in ordinary cohomology and the equivariant homeomorphism $(EG \times X)/(EG \times A) \cong (EG \times (X/A))/(EG \times pt)$.

The map $EG \times_G X \to X/G$ between the homotopy and ordinary quotients of X by G induces a homomorphism $H^*(X/G) \to H^*_G(X)$, which is an isomorphism in the case of a free G-action. In general, the equivariant cohomology contains more information about the action and its algebraic invariants than the ordinary cohomology of the orbit space.

A bundle $\pi\colon E \to X$ with fibre F is called a *G-bundle* if π is an equivariant map of G-spaces. By applying the Borel construction to a G-bundle $\pi\colon E \to X$ we obtain a bundle $B_G E \to B_G X$ with the same fibre F. The *equivariant characteristic classes* of a G-vector bundle $\pi\colon E \to X$ are defined as the ordinary characteristic classes of the corresponding bundle $B_G E \to B_G X$. For example, the *equivariant Stiefel–Whitney classes* of a vector G-bundle $\pi\colon E \to X$ belong to $H^*_G(X;\mathbb{Z}/2)$ and are denoted by $w^G_i(E)$. If $E \to X$ is an oriented G-vector bundle, then the *equivariant Euler class* $e^G(E) \in H^*_G(X;\mathbb{Z})$ is defined. If $E \to X$ is a complex bundle and the G-action preserves the fibrewise complex structure, then the *equivariant Chern classes* $c^G_i(E) \in H^{2i}_G(X;\mathbb{Z})$ are defined.

Now let M be a smooth oriented G-manifold of dimension n, where G is a compact Lie group. Let $N \subset M$ be a G-invariant (e.g., fixed) oriented submanifold of codimension k. We can identify the normal G-bundle $\nu(N \subset M)$ with a G-invariant tubular neighbourhood U of N in M by means of a G-equivariant diffeomorphism. The same diffeomorphism identifies the Thom space $Th\,\nu$ (see Section E.2) of the bundle ν with the quotient space $\overline{U}/\partial\overline{U}$. We have the embedding $i\colon N \subset M$, the projection $\pi\colon U \to N$, and the *Pontrjagin–Thom map* $p\colon M \to Th\,\nu$ contracting the complement $M \setminus U$ to a point. In equivariant cohomology, the *Thom class* $\tau_N \in H^k_G(Th\,\nu)$ is uniquely determined by the identity

$$\big(\alpha \cdot p^*(\tau_N), \langle M\rangle\big) = \big(i^*\alpha, \langle N\rangle\big),$$

for any $\alpha \in H^{n-k}_G(M)$. Here $\langle M\rangle \in H^G_n(M)$ denotes the fundamental class of M in equivariant homology. The *Gysin homomorphism* in equivariant cohomology is defined by the composition

$$H^{*-k}_G(N) \xrightarrow{\pi^*} H^{*-k}_G(U) \xrightarrow{\cdot\tau_N} H^*_G(Th\,\nu) \xrightarrow{p^*} H^*_G(M),$$

and is denoted by i_*. Then $i^*(i_*(1)) \in H^k_G(N)$ is the equivariant Euler class of the normal bundle $\nu(N \subset M)$.

Exercises.

B.3.1. Let G be a compact group acting on a Hausdorff space X. Show that the G-orbits are closed, the orbit space X/G is Hausdorff, and the projection $\pi\colon X \to X/G$ is a closed and proper map.

B.3.2. If the quotient X/G is Hausdorff, then the G-orbits are closed.

B.3.3. Give an example of a Hausdorff space X with a G-action whose orbits are closed, but the quotient X/G is not Hausdorff.

B.4. Eilenberg–Moore spectral sequences

In their paper [**83**] of 1966, Eilenberg and Moore constructed a spectral sequence, which became one of the important computational tools of algebraic topology. It particular, it provides a method for calculation of cohomology of the fibre of a bundle $E \to B$ using the canonical $H^*(B)$-module structure on $H^*(E)$. This spectral sequence can be considered as an extension of Adams' approach to calculating cohomology of loop spaces [**1**]. In the 1960–70s applications of the Eilenberg–Moore spectral sequence led to many important results on cohomology of homogeneous spaces for Lie groups. More recently it has been used for different calculations with toric spaces. This appendix section contains the necessary information about the spectral sequence; we mainly follow L. Smith's paper [**202**] in this description. For a detailed account of differential homological algebra and the Eilenberg–Moore spectral sequence, as well as its applications which go beyond the scope of this book, we refer to McCleary's book [**153**].

Here we assume that $\mathbf{k}$ is a field. The following theorem provides an algebraic setup for the Eilenberg–Moore spectral sequence.

THEOREM B.4.1 (Eilenberg–Moore [**202**, Theorem 1.2]). *Let A be a differential graded $\mathbf{k}$-algebra, and let M, N be differential graded A-modules. Then there exists a spectral sequence $\{E_r, d_r\}$ converging to $\operatorname{Tor}_A(M,N)$ and whose E_2-term is*

$$E_2^{-i,j} = \operatorname{Tor}^{-i,j}_{H[A]}\big(H[M], H[N]\big), \quad i,j \geqslant 0,$$

where $H[\,\cdot\,]$ denotes the algebra or module of cohomology.

REMARK. The construction of Tor for differential graded objects requires some additional considerations (see e.g. [**202**] or [**145**, Chapter XII]).

The spectral sequence of Theorem B.4.1 lives in the second quadrant and its differentials d_r add $(r, 1-r)$ to the bidegree, for $r \geqslant 1$. We shall refer to it as the *algebraic Eilenberg–Moore spectral sequence*. Its E_∞-term is expressed via a certain decreasing filtration $\{F^{-p}\operatorname{Tor}_A(M,N)\}$ in $\operatorname{Tor}_A(M,N)$ by the formula

$$E_\infty^{-p,n+p} = F^{-p}\Bigl(\sum_{-i+j=n}\operatorname{Tor}_A^{-i,j}(M,N)\Bigr)\Big/F^{-p+1}\Bigl(\sum_{-i+j=n}\operatorname{Tor}_A^{-i,j}(M,N)\Bigr).$$

Topological applications of Theorem B.4.1 arise in the case when A, M, N are cochain algebras of topological spaces. The classical situation is described by the commutative diagram

$$\begin{tikzcd} E \arrow[r] \arrow[d] & E_0 \arrow[d] \\ B \arrow[r] & B_0, \end{tikzcd} \tag{B.1}$$

where $E_0 \to B_0$ is a Serre fibre bundle with fibre F over a simply connected base B_0, and $E \to B$ is the pullback along a continuous map $B \to B_0$. For any space X, let $C^*(X)$ denote the singular $\mathbf{k}$-cochain algebra of X. Then $C^*(E_0)$ and $C^*(B)$ are $C^*(B_0)$-modules. Under these assumptions the following statement holds.

Lemma B.4.2 ([**202**, Proposition 3.4]). *$\operatorname{Tor}_{C^*(B_0)}(C^*(E_0), C^*(B))$ is a $\mathbf{k}$-algebra in a natural way, and there is a canonical isomorphism of algebras*

$$\operatorname{Tor}_{C^*(B_0)}\bigl(C^*(E_0), C^*(B)\bigr) \to H^*(E).$$

Applying Theorem B.4.1 in the case $A = C^*(B_0)$, $M = C^*(E_0)$, $N = C^*(B)$ and taking into account Lemma B.4.2, we come to the following statement.

Theorem B.4.3 (Eilenberg–Moore). *There exists a spectral sequence $\{E_r, d_r\}$ of commutative algebras converging to $H^*(E)$ with*

$$E_2^{-i,j} = \operatorname{Tor}^{-i,j}_{H^*(B_0)}\bigl(H^*(E_0), H^*(B)\bigr).$$

The spectral sequence of Theorem B.4.3 is known as the (topological) *Eilenberg–Moore spectral sequence*. The case when B is a point is of particular importance, and we state the corresponding result separately.

Corollary B.4.4. *Let $E \to B$ be a fibration over a simply connected space B with fibre F. Then there exists a spectral sequence $\{E_r, d_r\}$ of commutative algebras converging to $H^*(F)$ with*

$$E_2 = \operatorname{Tor}_{H^*(B_0)}\bigl(H^*(E_0), \mathbf{k}\bigr).$$

We refer to the spectral sequence of Corollary B.4.4 as the *Eilenberg–Moore spectral sequence of the fibration $E \to B$*. In the case when E_0 is a contractible space we obtain a spectral sequence converging to the cohomology of the loop space ΩB_0.

Remark. In rational homotopy theory, the Sullivan–de Rham algebra $A^*(X)$ of piecewise polynomial forms is used as a (graded) commutative algebraic model for X, instead of the rational singular cochain algebra $C^*(X;\mathbb{Q})$, which is not commutative. It is proved in [**30**, §3] that the above results on the Eilenberg–Moore spectral sequence hold with C^* replaced by A^*. This result is not a direct corollary of algebraic properties of Tor, since the integration map $A^*(X) \to C^*(X,\mathbb{Q})$ is not multiplicative.

APPENDIX C

Categorical constructions

APPENDIX D

Regular T-manifolds and their orbit structures

In this appendix we present the results of Chapter 1 of Izmestiev's thesis [**127**], which summarise and generalise several earlier developments on equivariant topology, including those of Jänich [**128**], Bredon [**31**], Davis [**69**] and Davis–Januszkiewicz [**70**]. The reader show refer here for the background material on locally standard $\mathbb{T}^n$- and $\mathbb{Z}_2^n$-actions, their topological and smooth classification, and combinatorial structures in their orbit spaces. The latter include the notions of *manifolds with faces* and *manifolds with corners*, which are of independent interest in differential topology.

As usual, we denote by $\mathbb{Z}_2$ (respectively, $\mathbb{S} = \mathbb{T}$) the multiplicative group of real (respectively, complex) numbers of absolute value one.

Unless otherwise specified, in this appendix we denote by G one of the groups $\mathbb{Z}_2$ or $\mathbb{S} = \mathbb{T}$, and denote by $\mathbb{F}$ its ambient field $\mathbb{R}$ or $\mathbb{C}$ respectively. We refer to the coordinatewise action of G^n on $\mathbb{F}^n$ given by

$$(g_1, \ldots, g_n) \cdot (z_1, \ldots, z_n) = (g_1 z_1, \ldots, g_n z_n)$$

as the *standard action* or *standard representation.*

D.1. Locally standard actions, manifolds with faces, and manifolds with corners

Definition D.1.1. Let M be a with an action of G^n. A *standard chart* on M is a triple (U, f, ψ), where $U \subset M$ is a G^n-invariant open subset, ψ is an automorphism of the torus $\mathbb{T}^n$, and f is a ψ-equivariant homeomorphism $f\colon U \to W$ onto a G^n-invariant open subset $W \subset \mathbb{F}^n$. (Recall from Appendix B.3 that the latter means that $f(\boldsymbol{t} \cdot y) = \psi(\boldsymbol{t}) f(y)$ for all $\boldsymbol{t} \in \mathbb{T}^n$, $y \in U$.) A $\mathbb{T}^n$-action on M is said to be *locally standard* if M has a standard atlas, i.e. if any point of M belongs to a standard chart.

The dimension of a manifold with a locally standard G^n-action is n if $G = \mathbb{Z}_2$ and $2n$ if $G = \mathbb{T}$. The orbit space of the standard G^n-action on $\mathbb{F}^n$ is the orthant

$$\mathbb{R}^n_{\geqslant} = \bigl\{\boldsymbol{x} = (x_1, \ldots, x_n) \in \mathbb{R}^n \colon x_i \geqslant 0 \quad \text{for } i = 1, \ldots, n\bigr\}.$$

Therefore, the orbit space of a locally standard action is locally modelled by $\mathbb{R}^n_{\geqslant}$. There are two slightly different ways to formalise this property, depending on whether we work in the topological or smooth category.

Definition D.1.2. An *manifold with faces* (of dimension n) is a topological manifold Q with boundary ∂Q together with a covering of ∂Q by closed connected subsets $\{F_i\}_{i \in \mathcal{S}}$, called *facets*, satisfying the following properties:

(a) any facet F_i is an $(n-1)$-dimensional submanifold of ∂Q;

(b) for any finite subset $I \subset \mathcal{S}$, the intersection $\bigcap_{i\in I} F_i$ is either empty or a disjoint union of submanifolds of codimension $|I|$; in the latter case we refer to a connected component of the intersection $\bigcap_{i\in I} F_i$ as a *face* of Q;

(c) for any point $q \in Q$, there exist an open neighbourhood $U \ni q$ and a homeomorphism $\varphi\colon U \to W$ onto an open subset $W \subset \mathbb{R}^n_\geqslant$ such that

$$\varphi^{-1}(W \cap \{x_j = 0\}) = U \cap F_i$$

for some $i \in \mathcal{S}$.

Observe that set $\mathcal{S}$ has to be countable, hence the set of faces in a manifold with faces Q is also countable. If Q is compact, then both these sets are finite.

The orthant $\mathbb{R}^n_\geqslant$ itself has the canonical structure of a manifold with faces; each face has the form $\mathbb{R}^I_\geqslant$ for $I \subset [n]$, where $\mathbb{R}^I_\geqslant = \{\boldsymbol{x} \in \mathbb{R}^n_\geqslant \colon x_j = 0 \quad \text{for } j \notin I\}$. The *codimension* $c(\boldsymbol{x})$ of point $\boldsymbol{x} \in \mathbb{R}^n_\geqslant$ is the number of zero coordinates of $\boldsymbol{x}$. A point has *codimension* k if it belongs to a face of codimension k and does not belong to a face of codimension $k+1$.

DEFINITION D.1.3. An *manifold with corners* (of dimension n) is a topological manifold Q with boundary together with an atlas $\{U_i, \varphi_i\}$ consisting of homeomorphisms $\varphi_i\colon U_i \to W_i$ onto open subsets $W_i \subset \mathbb{R}^n_\geqslant$ such that $\varphi_i\varphi_j^{-1}\colon \varphi_j(U_i \cap U_j) \to \varphi_i(U_i \cap U_j)$ is a diffeomorphism for all i, j. (A homeomorphism between open subsets in $\mathbb{R}^n_\geqslant$ is called a *diffeomorphism* if it can be obtained by restriction of a diffeomorphism of open subsets in $\mathbb{R}^n$.)

For any $q \in Q$, its codimension $c(q)$ is well-defined. An *open face* of Q of codimension k is a connected component of $c^{-1}(k)$. A *closed face* (or simply *face*) of Q is the closure of an open face.

A manifold with corners Q is said to be *nice* if the covering of Q by its facets satisfies condition (c) of Definition D.1.2 (conditions (a) and (b) are satisfied automatically).

REMARK. Our definitions of faces differ from those of [**69**] and [**127**]; the reason is that we want our faces to be connected.

EXAMPLE D.1.4.

1. The 2-disc with a single 'corner point' on its boundary is a manifold with corners which is not nice. All other examples in this list will be nice.

2. A smooth manifold Q with boundary is a manifold with corners, whose facets are connected components of ∂Q, and there are no other faces.

3. A product of manifolds with corners is a manifold with corners. In particular, a product of smooth manifolds with boundary is a manifold with corners.

4. Let P be a simple polytope. For each vertex $v \in P$ we denote by U_v the open subset in P obtained by removing all faces of P that do not contain v. The subset U_v is affinely isomorphic to a neighbourhood of zero in $\mathbb{R}^n_\geqslant$. Therefore P is a compact manifold with corners, with atlas $\{U_v\}$.

As we shall see in the next sections, the orbit space $Q = M/\mathbb{T}^n$ of a locally standard torus action is a manifold with faces, and if the action is smooth, then the orbit space is a nice manifold with corners.

Exercises.

D.1.5. A manifold with corners Q is nice if an only if any point of codimension two is contained in exactly two facets.

D.1.6. Two simple polytopes are diffeomorphic as manifolds with corners if and only if they are combinatorially equivalent.

APPENDIX E

Stably complex structures and complex bordism

Here we summarise the required facts from the theory of bordism and cobordism, with the most attention given to complex (co)bordism. Cobordism theory is one of the deepest and most influential parts of algebraic topology, which experienced a spectacular development in the 1960s. Proofs of many results presented here would require a separate monograph and a substantial background in algebraic topology. There are a few exceptions where the proofs can be made concise, in which cases we include them here. For the rest an interested reader is referred to the works of Novikov [**171**], [**172**], and monographs of Conner–Floyd [**63**] and Stong [**212**].

We consider topological spaces which have the homotopy type of cell complexes. All manifolds are assumed to be smooth, compact and closed (without boundary), unless otherwise specified.

E.1. Bordism of manifolds

Given two n-dimensional manifolds M_0 and M_1, a *bordism* between them is an $(n+1)$-dimensional manifold W with boundary, whose boundary is the disjoint union of M_0 and M_1, that is, $\partial W = M_0 \sqcup M_1$. If such a W exists, M_0 and M_1 are called *bordant*. The bordism relation splits the set of manifolds into equivalence classes (see Fig. E.1.1), which are called *bordism classes*.

We denote the bordism class of M by $[M]$, and denote by $\varOmega_n^O$ the set of bordism classes of n-dimensional manifolds. Then $\varOmega_n^O$ is an Abelian group with respect to the disjoint union operation: $[M_1] + [M_2] = [M_1 \sqcup M_2]$. Zero is represented by the bordism class of the empty set (which is counted as a manifold in any dimension), or by the bordism class of any manifold which bounds. We also have $\partial(M \times I) = M \sqcup M$. Hence, $2[M] = 0$ and $\varOmega_n^O$ is a 2-torsion group.

Set $\varOmega_*^O = \bigoplus_{n\geqslant 0} \varOmega_n^O$. The direct product of manifolds induces a multiplication of bordism classes, namely $[M_1] \times [M_2] = [M_1 \times M_2]$. It makes $\varOmega_*^O$ a graded commutative ring, the *unoriented bordism ring.*

For any space X the bordism relation can be extended to maps of manifolds to X: two maps $M_1 \to X$ and $M_2 \to X$ are *bordant* if there is a bordism W between M_1 and M_2 and the map $M_1 \sqcup M_2 \to X$ extends to a map $W \to X$. The set of bordism classes of maps $M \to X$ with $\dim M = n$ forms an abelian group called the *n-dimensional unoriented bordism group of* X and denoted $O_n(X)$ (other notations: $N_n(X)$, $MO_n(X)$). We note that $O_n(pt) = \varOmega_n^O$, where *pt* is a point.

The bordism group $O_n(X, A)$ of a pair $A \subset X$ is defined as the set of bordism classes of maps of manifolds with boundary, $(M, \partial M) \to (X, A)$, where $\dim M = n$.

Figure E.1. Transitivity of the bordism relation.

(Two such maps $f_0\colon (M_0, \partial M_0) \to (X, A)$ and $f_1\colon (M_1, \partial M_1) \to (X, A)$ are *bordant* if there is W such that $\partial W = M_0 \cup M_1 \cup M$, where M is a bordism between ∂M_0 and ∂M_1, and a map $f\colon W \to X$ such that $f|_{M_0} = f_0$, $f|_{M_1} = f_1$ and $f(M) \subset A$.) We have $O_n(X, \varnothing) = O_n(X)$.

There is an obviously defined map $\varOmega^O_m \times O_n(X) \to O_{m+n}(X)$ turning $O_*(X) = \bigoplus_{n\geqslant 0} O_n(X)$ into a graded $\varOmega^O_*$-module. The assignment $X \mapsto O_*(X)$ defines a *generalised homology theory*, that is, it is functorial in X, homotopy invariant, has the excision property and exact sequences of pairs.

E.2. Thom spaces, bordism and cobordism

A remarkable geometric construction due to Pontrjagin and Thom reduces the calculation of the bordism groups $O_n(X)$ to a homotopical problem. Here we assume known basic facts from the theory of vector bundles.

Given an n-dimensional real Euclidean vector bundle ξ with total space $E = E\xi$ and Hausdorff compact base X, the *Thom space* of ξ is defined as the quotient

$$Th\,\xi = E/E_{\geqslant 1},$$

where $E_{\geqslant 1}$ is the subspace consisting of vectors of length $\geqslant 1$ in the fibres of ξ. Equivalently, $Th\,\xi = BE/SE$, where BE is the total space of the n-ball bundle associated with ξ and $SE = \partial BE$ is the $(n-1)$-sphere bundle. Also, $Th\,\xi$ is the one-point compactification of E.

Given two spaces X and Y with basepoints pt_X and pt_Y respectively, their *smash product* is defined as

$$X \wedge Y = X \times Y \big/ \big((X \times pt_Y) \cup (pt_X \times Y)\big).$$

The Thom space $Th\,\xi$ has a canonical basepoint, the image of $E_{\geqslant 1}$.

Proposition E.2.1. *If ξ and η are vector bundles over X and Y respectively, and $\xi \times \eta$ is the product vector bundle over $X \times Y$, then*

$$Th(\xi \times \eta) = Th\,\xi \wedge Th\,\eta.$$

The proof is left as an exercise.

Example E.2.2.

1. Regarding $\mathbb{R}^k$ as the total space of a k-plane bundle over a point, we obtain that the corresponding Thom space $Th(\mathbb{R}^k)$ is a k-sphere S^k.

2. If ξ is a 0-dimensional bundle over X, then $Th\,\xi = X_+ = X \sqcup pt$.

3. Let $\underline{\mathbb{R}}^k$ denote the trivial k-plane bundle over X. The Whitney sum $\xi \oplus \underline{\mathbb{R}}^k$ can be identified with the product bundle $\xi \times \mathbb{R}^k$, where $\mathbb{R}^k$ is the k-plane bundle over a point. Then Proposition E.2.1 implies that

$$Th(\xi \oplus \underline{\mathbb{R}}^k) = \Sigma^k\, Th\,\xi,$$

where Σ^k denote the k-fold suspension.

4. Combining the previous two examples, we obtain

$$Th(\underline{\mathbb{R}}^k) = \Sigma^k X \vee S^k.$$

Construction E.2.3 (Pontrjagin–Thom construction). Let M be a submanifold in $\mathbb{R}^m$ with normal bundle $\nu = \nu(M \subset \mathbb{R}^m)$. The *Pontrjagin–Thom collapse map*

$$S^m \to Th\,\nu$$

identifies the tubular neighbourhood of M in $\mathbb{R}^m \subset S^m$ with the set of vectors of length < 1 in the fibres of ν, and collapses the complement of the tubular neighbourhood to the basepoint of the Thom space $Th\,\nu$.

This construction can be generalised to submanifolds $M \subset E\xi$ in the total space of an arbitrary m-plane bundle ξ over a manifold, giving the collapse map

$$Th\,\xi \to Th\,\nu, \tag{E.1}$$

where $\nu = \nu(M \subset E\xi)$. Note that the Pontrjagin–Thom collapse map is a particular case of (E.1), as S^m is the Thom space of an m-plane bundle over a point.

Recall that a smooth map $f\colon W \to Z$ of manifolds is called *transverse* along a submanifold $Y \subset Z$ if, for every $w \in f^{-1}(Y)$, the image of the tangent space to W at w together with the tangent space to Y at $f(w)$ spans the tangent space to Z at $f(w)$:

$$f_*\mathcal{T}_w W + \mathcal{T}_{f(w)}Y = \mathcal{T}_{f(w)}Z.$$

If $f\colon W \to Z$ is transverse along $Y \subset Z$, then $f^{-1}(Y)$ is a submanifold in W of codimension equal to the codimension of Y in Z.

Construction E.2.4 (cobordism classes of η-submanifolds in $E\xi$). Let ξ be an m-plane bundle with total space $E\xi$ over a manifold X, and let η be an n-plane bundle over a manifold Y. An *η-submanifold* of $E\xi$ is a pair (M, f) consisting of a submanifold M in $E\xi$ and a map

$$f\colon \nu(M \subset E\xi) \to \eta$$

such that f is an isomorphism on each fibre (so that the codimension of M in $E\xi$ is n). Two η-submanifolds (M_0, f_0) and (M_1, f_1) are said to be *cobordant* if there is an η-submanifold with boundary (W, f) in the cylinder $E\xi \times I$ such that

$$\partial(W, f) = \bigl((M_0, f_0) \times 0\bigr) \cup \bigl((M_1, f_1) \times 1\bigr).$$

Theorem E.2.5. *The set of cobordism classes of η-submanifolds in $E\xi$ is in one-to-one correspondence with the set $[Th\xi, Th\eta]$ of homotopy classes of based maps of Thom spaces.*

Proof. Assume given a based map $g\colon\ Th\,\xi \to Th\,\eta$. By changing g within its homotopy class we may achieve that g is transverse along the zero section $Y \subset Th\,\eta$ (transversality is a local condition, and both $Th\,\xi$ and $Th\,\eta$ are manifolds outside the basepoints). Since η is an n-plane bundle, $M = g^{-1}(Y)$ is a submanifold of codimension n in $E\xi = Th\,\xi \setminus pt$ such that

$$\nu(M \subset E\xi) = g^*(\nu(Y \subset E\eta)) = g^*\eta.$$

That is, M is an η-submanifold in $E\xi$.

Conversely, assume given an η-submanifold $M \subset E\xi$. We therefore have the map of Thom spaces $Th\,\nu \to Th\,\eta$ (induced by the map of ν to η), whose composition with the collapse map (E.1) gives the required map $Th\,\xi \to Th\,\eta$.

The fact that homotopic based maps $Th\,\xi \to Th\,\eta$ correspond to cobordant η-submanifolds, and vice versa, is left as an exercise. □

Construction E.2.6 (cobordism groups). Let η_k be the universal vector k-plane bundle $EO(k) \to BO(k)$. Following the original notation of Thom, we denote $MO(k) = Th\,\eta_k$.

Every submanifold $M \subset \mathbb{R}^{n+k}$ of dimension n is an η_k-submanifold via the classifying map of the normal bundle $\nu(M \subset \mathbb{R}^{n+k})$. Denote by $\varOmega_O^{-n,k}$ the set of cobordism classes of $M \subset \mathbb{R}^{n+k}$. The base of η_k is not a manifold, but it is a direct limit of Grassmannians, and a simple limit argument shows that Theorem E.2.5 still holds for η_k-submanifolds. Hence,

$$\varOmega_O^{-n,k} = [S^{n+k}, \mathit{Th}\, \eta_k] = \pi_{n+k}(MO(k)).$$

There is the stabilisation map $\varOmega_O^{-n,k} \to \varOmega_O^{-n,k+1}$ obtained by composing the suspended map $S^{n+k+1} \to \Sigma\, MO(k)$ with the map $\Sigma\, MO(k) \to MO(k+1)$ induced by the bundle map $\eta_k \oplus \underline{\mathbb{R}} \to \eta_{k+1}$. We therefore may define the $(-n)$*th cobordism group* as

$$\varOmega_O^{-n} = \lim_{k\to\infty} \varOmega_O^{-n,k} = \lim_{k\to\infty} \pi_{n+k}(MO(k)). \tag{E.2}$$

PROPOSITION E.2.7. *We have a canonical isomorphism*

$$\varOmega_O^{-n} \cong \varOmega_n^O$$

between the cobordism and bordism groups, for $n \geqslant 0$. *In other words, two* n*-dimensional manifolds* M_0 *and* M_1 *are bordant if and only if there exist embeddings of* M_0 *and* M_1 *in the same* $\mathbb{R}^{n+k}$ *which are cobordant.*

PROOF. Forgetting the embedding $M \subset \mathbb{R}^{n+k}$ we obtain a map $\varOmega_O^{-n,k} \to \varOmega_n^O$, which may be shown to be a group homomorphism. It is compatible with the stabilisation maps, and therefore defines a homomorphism $\varOmega_O^{-n} \to \varOmega_n^O$. Since every manifold M may be embedded in some $\mathbb{R}^{n+k}$, it is an isomorphism. □

Together with (E.2), Proposition E.2.7 gives a homotopical interpretation for the (unoriented) bordism groups. This also implies that the notions of the 'bordism class' and 'cobordism class' of a manifold M are interchangeable. Theorem E.2.5 may be applied further to obtain a homotopical interpretation for the bordism groups $O_n(X)$ of a space X:

CONSTRUCTION E.2.8 (bordism and cobordism groups of a space). Let X be a space. We set $\xi = \mathbb{R}^{n+k}$ (an $(n+k)$-plane bundle over a point) and $\eta = X \times \eta_k$ (the product of a 0-plane bundle over X and the universal k-plane bundle η_k over $BO(k)$), and consider cobordism classes of η-submanifolds in $E\xi$. Such a submanifold is described by a pair (f, ι) consisting of a map $f\colon M \to X$ from an n-dimensional manifold to X and an embedding $\iota\colon M \to \mathbb{R}^{n+k}$ (the bundle map from $\nu(\iota)$ to η is the product of the map $E\nu(\iota) \to M \to X$ and the classifying map of $\nu(\iota)$). By Theorem E.2.5, the set of cobordism classes of η-submanifolds in $E\xi$ is given by

$$[S^{n+k}, \mathit{Th}(X \times \eta_k)] = \pi_{n+k}\bigl((X_+) \wedge MO(k)\bigr).$$

As in the proof of Proposition E.2.7, there is the map from the above set of cobordism classes to the bordism group $O_n(X)$, which forgets the embedding $M \subset \mathbb{R}^{n+k}$. A stabilisation argument shows that

$$O_n(X) = \lim_{k\to\infty} \pi_{k+n}\bigl((X_+) \wedge MO(k)\bigr), \tag{E.3}$$

providing a homotopical interpretation for the bordism groups of X.

We define the *cobordism groups* of X as

$$O^n(X) = \lim_{k\to\infty} \bigl[\Sigma^{k-n}(X_+), MO(k)\bigr]. \tag{E.4}$$

If X is a manifold, then the groups $O^n(X)$ may also be obtained by stabilising the set of cobordism classes of η-submanifolds in $E\xi$. Namely, we need to set $\xi = \underline{\mathbb{R}}^{k-n}$ (the trivial $(k-n)$-plane bundle over X), and $\eta = \eta_k$. In other words, a cobordism class in $O^n(X)$ is described by the composition

$$M \hookrightarrow X \times \mathbb{R}^{k-n} \longrightarrow X, \tag{E.5}$$

where the first map is an embedding of codimension k.

The maps of Thom spaces $MO(k) \wedge MO(l) \to MO(k+l)$ (induced by the classifying maps $\eta_k \oplus \eta_l \to \eta_{k+l}$ of the Whitney sums) turn $O^*(X) = \bigoplus_{n\in\mathbb{Z}} O^n(X)$ into a graded ring, called the *unoriented cobordism ring of* X.

Exercises.

E.2.9. Prove Proposition E.2.1.

E.2.10. If η is the tautological line bundle over $\mathbb{R}P^n$ (respectively, $\mathbb{C}P^n$), then $Th\,\eta$ can be identified with $\mathbb{R}P^{n+1}$ (respectively, $\mathbb{C}P^{n+1}$).

E.2.11. Prove that cobordism of η-submanifolds is an equivalence relation.

E.2.12. Complete the proof of Theorem E.2.5.

E.2.13. The forgetful map $\varOmega_O^{-n,k} \to \varOmega_n^O$ is a homomorphism of groups.

E.2.14. Given any $(k-n)$-plane bundle ξ over a manifold X and an embedding $M \hookrightarrow E\xi$ of codimension k, the composition

$$M \hookrightarrow E\xi \longrightarrow X$$

determines a cobordism class in $O^n(X)$. (Hint: reduce to (E.5) by embedding ξ into a trivial bundle over the same X.)

E.2.15 (Poincaré–Atiyah duality [**4**] in unoriented bordism). If X is an n-dimensional manifold, then

$$O^{n-k}(X) = O_k(X) \quad \text{for any } k.$$

In particular, for $X = pt$ we obtain the isomorphisms of Proposition E.2.7.

E.3. Oriented and complex bordism

The bordism relation may be extended to manifolds endowed with some additional structure, which leads to important bordism theories.

The simplest additional structure is an orientation. By definition, two oriented n-dimensional manifolds M_1 and M_2 are *oriented bordant* if there is an oriented $(n+1)$-dimensional manifold W with boundary such that $\partial W = M_1 \sqcup \overline{M}_2$, where $\overline{M}_2$ denotes M_2 with the orientation reversed. The *oriented bordism groups* $\varOmega_n^{SO}$ and the *oriented bordism ring* $\varOmega_*^{SO} = \bigoplus_{n\geqslant 0} \varOmega_n^{SO}$ are defined accordingly. Given an oriented manifold M, the manifold $M \times I$ has the canonical orientation such that $\partial(M \times I) = M \sqcup \overline{M}$. Hence, $-[M] = [\overline{M}]$ in $\varOmega_n^{SO}$. Unlike $\varOmega_n^O$, elements of $\varOmega_*^{SO}$ generally do not have order 2.

Complex structure gives another important example of an additional structure on manifolds. However, a direct attempt to define the bordism relation on complex manifolds fails because the manifold W must be odd-dimensional and therefore cannot be complex. This can be remedied by considering *stably complex* (also known as *stably almost complex* or *quasicomplex*) structures.

Let $\mathcal{T}M$ denote the tangent bundle of M. We say that M admits a *tangential stably complex structure* if there is an isomorphism of real vector bundles

$$c_{\mathcal{T}}\colon \mathcal{T}M \oplus \underline{\mathbb{R}}^k \to \xi \tag{E.6}$$

between the 'stable' tangent bundle and a complex vector bundle ξ over M. Some of the choices of such isomorphisms are deemed to be equivalent, that is, determine the same stably complex structures. This equivalence relation is generated by

(a) additions of trivial complex summands; that is, $c_{\mathcal{T}}$ is equivalent to

$$\mathcal{T}M \oplus \underline{\mathbb{R}}^k \oplus \underline{\mathbb{C}} \xrightarrow{c_{\mathcal{T}}\oplus\mathrm{id}} \xi \oplus \underline{\mathbb{C}},$$

where $\underline{\mathbb{C}}$ in the left hand side is canonically identified with $\underline{\mathbb{R}}^2$;

(b) compositions with isomorphisms of complex bundles; that is, $c_{\mathcal{T}}$ is equivalent to $\varphi \cdot c_{\mathcal{T}}$ for every $\mathbb{C}$-linear isomorphism $\varphi\colon \xi \to \zeta$.

The equivalence class of $c_{\mathcal{T}}$ may be described homotopically as the equivalence class of lifts of the map $M \to BO(2N)$ classifying the stable tangent bundle to a map $M \to BU(N)$ up to homotopy and stabilisation (see [**212**, Chapters II, VII]). A *tangential stably complex manifold* is a pair consisting of M and an equivalence class of isomorphisms $c_{\mathcal{T}}$; we shall use a simplified notation $(M, c_{\mathcal{T}})$ for such pairs. This notion is a generalisation of complex and *almost complex* manifolds (where the latter means a manifold with a choice of a complex structure on $\mathcal{T}M$, that is, a stably complex structure (E.6) with $k = 0$).

We say that M admits a *normal complex structure* if there is an embedding $i\colon M \hookrightarrow \mathbb{R}^N$ with the property that the normal bundle $\nu(i)$ admits a structure of a complex vector bundle. There is an appropriate notion of stable equivalence for such embeddings i, and a normal complex structure c_ν on M is defined as the corresponding equivalence class. Tangential and normal stably complex structures on M determine each other by means of the canonical isomorphism $\mathcal{T}M \oplus \nu(i) \cong \underline{\mathbb{R}}^N$.

EXAMPLE E.3.1. Let $M = \mathbb{C}P^1$. The standard complex structure on M is equivalent to the stably complex structure determined by the isomorphism

$$\mathcal{T}(\mathbb{C}P^1) \oplus \underline{\mathbb{R}}^2 \xrightarrow{\cong} \overline{\eta} \oplus \overline{\eta}$$

where η is the tautological complex line bundle. On the other hand, the isomorphism

$$\mathcal{T}(\mathbb{C}P^1) \oplus \underline{\mathbb{R}}^2 \xrightarrow{\cong} \eta \oplus \overline{\eta} \cong \underline{\mathbb{C}}^2$$

determines a trivial stably complex structure on $\mathbb{C}P^1$.

The bordism relation can be defined between stably complex manifolds by taking account of the stably complex structure in the bordism relation. As in the case of unoriented bordism, the set of bordism classes $[M, c_{\mathcal{T}}]$ of n-dimensional stably complex manifolds is an Abelian group with respect to disjoint union. This group is called the *n-dimensional complex bordism group* and denoted by Ω_n^U. The sphere S^n has the canonical normally complex structure determined by a complex structure on the trivial normal bundle of the embedding $S^n \hookrightarrow \mathbb{R}^{n+2}$. The corresponding bordism class represents the zero element in Ω_n^U. The opposite element to the bordism class $[M, c_{\mathcal{T}}]$ in the group Ω_n^U may be represented by the same manifold M with the stably complex structure determined by the isomorphism

$$\mathcal{T}M \oplus \underline{\mathbb{R}}^k \oplus \underline{\mathbb{C}} \xrightarrow{c_{\mathcal{T}}\oplus\tau} \xi \oplus \underline{\mathbb{C}}$$

where $\tau\colon \mathbb{C} \to \mathbb{C}$ is the complex conjugation. We shall use the abbreviated notations $[M]$ and $[\overline{M}]$ for the complex bordism class and its opposite whenever the stably complex structure $c_{\mathcal{T}}$ is clear from the context. There is a stably complex structure on $M \times I$ such that $\partial(M \times I) = M \sqcup \overline{M}$.

The direct product of stably complex manifolds turns $\varOmega^U_* = \bigoplus_{n\geqslant 0} \varOmega^U_n$ into a graded ring, called the *complex bordism ring*.

Construction E.3.2 (complex bordism and cobordism groups of a space). The *complex bordism groups* $U_n(X)$ and *cobordism groups* $U^n(X)$ may be defined homotopically similarly to (E.3) and (E.4):

$$\begin{aligned} U_n(X) &= \lim\nolimits_{k\to\infty} \pi_{2k+n}\bigl((X_+) \wedge MU(k)\bigr), \\ U^n(X) &= \lim\nolimits_{k\to\infty} \bigl[\Sigma^{2k-n}(X_+), MU(k)\bigr], \end{aligned} \tag{E.7}$$

where $MU(k)$ is the Thom space of the universal complex k-plane bundle $EU(k) \to BU(k)$. Here the direct limit uses the maps $\Sigma^2 MU(k) \to MU(k+1)$.

Both groups $U_n(X)$ and $U^n(X)$ may also be defined geometrically, in a way similar to the geometric construction of unoriented bordism and cobordism groups (Construction E.2.8). The complex bordism group $U_n(X)$ consists of bordism classes of maps $M \to X$ of stably complex n-dimensional manifolds M to X. Complex cobordism groups of a manifold X may be defined using the notion of *complex oriented maps*. A map $M \to X$ between manifolds is said to be *complex orientable of codimension* n if it can be decomposed as

$$M \hookrightarrow X \times \mathbb{R}^{2k-n} \longrightarrow X,$$

where the first map is an embedding whose normal bundle has a structure of a complex k-plane bundle, and the second map is the projection onto the first factor. A choice of this decomposition together with a complex bundle structure in the normal bundle is called a *complex orientation* of the map $M \to X$. The group $U^n(X)$ consists of cobordism classes of complex oriented maps $M \to X$ of codimension n.

If $X = pt$, then we obtain

$$U^{-n}(pt) = U_n(pt) = \varOmega^U_n$$

for $n \geqslant 0$, from either the homotopic or geometric description of the (co)bordism groups. We also set $\varOmega_U^{-n} = U^{-n}(pt)$.

The maps of Thom spaces $MU(k) \wedge MU(l) \to MU(k+l)$ (induced by the classifying maps of the Whitney sum) turn $U^*(X) = \bigoplus_{n\in\mathbb{Z}} U^n(X)$ into a graded ring, called the *complex cobordism ring of* X. It is also a module over $\varOmega_U^*$.

Example E.3.3. Let X be a manifold of dimension n. The inclusion of a point $pt \subset X$ defines the bordism class $1 \in U_0(X)$ and the *fundamental cobordism class of* X in $U^n(X)$. (The normal bundle of a point has a complex structure if n is even, otherwise the normal bundle of a point in $X \times \mathbb{R}$ has a complex structure.)

The identity map $X \to X$ defines the cobordism class $1 \in U^0(X)$. It defines the *fundamental bordism class of* X in $U_n(X)$ only when X is stably complex.

Here is another example of situation when certain cobordism classes can be represented very explicitly by maps of manifolds.

Construction E.3.4 (geometric cobordisms). For any cell complex X the cohomology group $H^2(X)$ can be identified with the set $[X, \mathbb{C}P^\infty]$ of homotopy classes of maps into $\mathbb{C}P^\infty$. Since $\mathbb{C}P^\infty = MU(1)$, it follows from (E.7) that every element $x \in H^2(X)$ determines a cobordism class $u_x \in U^2(X)$. The elements of $U^2(X)$ obtained in this way are called *geometric cobordisms* of X. We therefore may view $H^2(X)$ as a subset in $U^2(X)$, however the group operation in $H^2(X)$ is not obtained by restricting the group operation in $U^2(X)$ (the relationship between the two operations is discussed in Appendix F).

When X is a manifold, geometric cobordisms may be described by submanifolds $M \subset X$ of codimension 2 with a fixed complex structure on the normal bundle.

Indeed, every $x \in H^2(X)$ corresponds to a homotopy class of maps $f_x\colon X \to \mathbb{C}P^\infty$. The image $f_x(X)$ is contained in some $\mathbb{C}P^N \subset \mathbb{C}P^\infty$, and we may assume that $f_x(X)$ is transverse to a certain hyperplane $H \subset \mathbb{C}P^N$. Then $M_x = f_x^{-1}(H)$ is a codimension-2 submanifold in X whose normal bundle acquires a complex structure by restriction of the complex structure on the normal bundle of $H \subset \mathbb{C}P^N$. Changing the map f_x within its homotopy class does not affect the bordism class of the embedding $M_x \to X$.

Conversely, assume given a submanifold $M \subset X$ of codimension 2 whose normal bundle is endowed with a complex structure. Then the composition

$$X \to Th(\nu) \to MU(1) = \mathbb{C}P^\infty$$

of the Pontrjagin–Thom collapse map $X \to Th(\nu)$ and the map of Thom spaces corresponding to the classifying map $M \to BU(1)$ of ν defines an element $x_M \in H^2(X)$, and therefore a geometric cobordism.

If X is an oriented manifold, then a choice of complex structure on the normal bundle of a codimension-2 embedding $M \subset X$ is equivalent to orienting M. The image of the fundamental class of M in the homology of X is Poincaré dual to $x_M \in H^2(X)$.

Construction E.3.5 (connected sum). For manifolds of positive dimension the disjoint union $M_1 \sqcup M_2$ representing the sum of bordism classes $[M_1] + [M_2]$ may be replaced by their *connected sum*, which represents the same bordism class.

The connected sum $M_1 \# M_2$ of manifolds M_1 and M_2 of the same dimension n is constructed as follows. Choose points $v_1 \in M_1$ and $v_2 \in M_2$, and take closed ε-balls $B_\varepsilon(v_1)$ and $B_\varepsilon(v_2)$ around them (both manifolds may be assumed to be endowed with a Riemannian metric). Fix an isometric embedding f of a pair of standard ε-balls $D^n \times S^0$ (here $S^0 = \{0, 1\}$) into $M_1 \sqcup M_2$ which maps $D^n \times 0$ onto $B_\varepsilon(v_1)$ and $D^n \times 1$ onto $B_\varepsilon(v_2)$. If both M_1 and M_2 are oriented we additionally require the embedding f to preserve the orientation on the first ball and reverse in on the second. Now, using this embedding, replace in $M_1 \sqcup M_2$ the pair of balls $D^n \times S^0$ by a 'pipe' $S^{n-1} \times D^1$. After smoothing the angles in the standard way we obtain a smooth manifold $M_1 \# M_2$.

If both M_1 and M_2 are connected the smooth structure on $M_1 \# M_2$ does not depend on a choice of points v_1, v_2 and embedding $D^n \times S^0 \hookrightarrow M_1 \sqcup M_2$. It does however depend on the orientations; $M_1 \# M_2$ and $M_1 \# \overline{M_2}$ are not diffeomorphic in general. For example, the manifolds $\mathbb{C}P^2 \# \mathbb{C}P^2$ and $\mathbb{C}P^2 \# \overline{\mathbb{C}P^2}$ are not diffeomorphic (and not even homotopy equivalent because they have different signatures).

There are smooth contraction maps $p_1\colon M_1 \# M_2 \to M_1$ and $p_2\colon M_1 \# M_2 \to M_2$. In the oriented case the manifold $M_1 \# M_2$ can be oriented in such a way that both contraction maps preserve the orientations.

FIGURE E.2. Disjoint union and connected sum.

A bordism between $M_1 \sqcup M_2$ and $M_1 \# M_2$ may be constructed as follows. Consider a cylinder $M_1 \times I$, from which we remove an ε-neighbourhood $U_\varepsilon(v_1 \times 1)$ of the point $v_1 \times 1$. Similarly, remove the neighbourhood $U_\varepsilon(v_2 \times 1)$ from $M_2 \times I$ (each of these two neighbourhoods can be identified with the half of a standard open $(n+1)$-ball). Now connect the two remainders of cylinders by a 'half pipe' $S^n_{\leqslant} \times I$ in such a way that the half-sphere $S^n_{\leqslant} \times 0$ is identified with the half-sphere on the boundary of $U_\varepsilon(v_1 \times 1)$, and $S^n_{\leqslant} \times 1$ is identified with the half-sphere on the boundary of $U_\varepsilon(v_2 \times 1)$. Smoothing the angles we obtain a manifold with boundary $M_1 \sqcup M_2 \sqcup (M_1 \# M_2)$ (or $\overline{M_1} \sqcup \overline{M_2} \sqcup (M_1 \# M_2)$ in the oriented case), see Fig. E.2.

Finally, if M_1 and M_2 are stably complex manifolds, then there is a canonical stably complex structure on $M_1 \# M_2$, which is constructed as follows. Assume the stably complex structures on M_1 and M_2 are determined by isomorphisms

$$c_{\mathcal{T},1} \colon \mathcal{T}M_1 \oplus \underline{\mathbb{R}}^{k_1} \to \xi_1 \quad \text{and} \quad c_{\mathcal{T},2} \colon \mathcal{T}M_2 \oplus \underline{\mathbb{R}}^{k_2} \to \xi_2.$$

Using the isomorphism $\mathcal{T}(M_1 \# M_2) \oplus \underline{\mathbb{R}}^n \cong p_1^*\mathcal{T}M_1 \oplus p_2^*\mathcal{T}M_2$, we define a stably complex structure on $M_1 \# M_2$ by the isomorphism

$$\begin{aligned}&\mathcal{T}(M_1 \# M_2) \oplus \underline{\mathbb{R}}^{n+k_1+k_2}\\ &\qquad\qquad \cong p_1^*\mathcal{T}M_1 \oplus \underline{\mathbb{R}}^{k_1} \oplus p_2^*\mathcal{T}M_2 \oplus \underline{\mathbb{R}}^{k_2} \xrightarrow{c_{\mathcal{T},1} \oplus c_{\mathcal{T},2}} p_1^*\xi_1 \oplus p_2^*\xi_2.\end{aligned}$$

We shall refer to this stably complex structure as the *connected sum of stably complex structures* on M_1 and M_2. The corresponding complex bordism class is $[M_1] + [M_2]$.

Exercises.

E.3.6. Assume given a complex $(k-l)$-plane bundle ξ over a manifold X and an embedding $M \hookrightarrow E\xi$ whose normal bundle has a structure of a complex k-plane bundle. Then the composition

$$M \hookrightarrow E\xi \longrightarrow X$$

determines a complex orientation for the map $M \to X$ of codimension $2l$, and therefore a complex cobordism class in $U^{2l}(X)$. (Compare Exercise E.2.14.)

Similarly, an embedding $M \hookrightarrow E(\xi \oplus \underline{\mathbb{R}})$ whose normal bundle has a structure of a complex k-plane bundle determines a complex cobordism class in $U^{2l-1}(X)$, via the composition

$$M \hookrightarrow E(\xi \oplus \underline{\mathbb{R}}) \longrightarrow X.$$

This is how complex orientations were defined in [**193**].

E.3.7 (Poincaré–Atiyah duality [**4**] in complex bordism). If X is an n-dimensional stably complex manifold, then

$$U^{n-k}(X) = U_k(X) \quad \text{for any } k.$$

E.4. Structure results

The theory of unoriented (co)bordism was the first to be completed: the coefficient ring Ω^O_* was calculated by Thom, and the bordism groups $O_*(X)$ of cell complexes X were reduced to homology groups of X with coefficients in Ω^O_*. The corresponding results are summarised as follows.

Theorem E.4.1.

(a) *Two manifolds are unorientedly bordant if and only if they have identical sets of Stiefel–Whitney characteristic numbers.*
(b) *Ω^O_* is a polynomial ring over $\mathbb{Z}/2$ with one generator a_i in every positive dimension $i \neq 2^k - 1$.*
(c) *For every cell complex X the module $O_*(X)$ is a free graded Ω^O_*-module isomorphic to $H_*(X;\mathbb{Z}/2) \otimes_{\mathbb{Z}/2} \Omega^O_*$.*

Parts (a) and (b) were done by Thom [**216**]. Part (c) was first formulated by Conner and Floyd [**63**]; it also follows from the results of Thom.

Calculating the complex bordism ring Ω^U_* turned out to be a much more difficult problem:

Theorem E.4.2.

(a) *$\Omega^U_* \otimes \mathbb{Q}$ is a polynomial ring over $\mathbb{Q}$ generated by the bordism classes of complex projective spaces $\mathbb{C}P^i$, $i \geqslant 1$.*
(b) *Two stably complex manifolds are bordant if and only if they have identical sets of Chern characteristic numbers.*
(c) *Ω^U_* is a polynomial ring over $\mathbb{Z}$ with one generator a_i in every even dimension $2i$, where $i \geqslant 1$.*

Part (a) can be proved by the methods of Thom. Part (b) follows from the results of Milnor [**161**] and Novikov [**169**]. Part (c) is the most difficult one; it was done by Novikov [**169**] using the Adams spectral sequence and structure theory of Hopf algebras (see also [**170**] for a more detailed account) and Milnor (unpublished[1]). Another more geometric proof was given by Stong [**212**].

Note that part (c) of Theorem E.4.1 does not extend to complex bordism; $U_*(X)$ is not a free Ω^U_*-module in general (although it is a free Ω^U_*-module if $H_*(X;\mathbb{Z})$ is free abelian). Unlike the case of unoriented bordism, the calculation of complex bordism of a space X does not reduce to calculating the coefficient ring Ω^U_* and homology groups $H_*(X)$. The theory of complex (co)bordism is much richer than its unoriented analogue, and at the same time is not as complicated as oriented bordism or other bordism theories with additional structure, since the coefficient ring does not have torsion. Thanks to this richness and lack of torsion, complex cobordism theory found many striking and important applications in algebraic topology and beyond. Many of these applications were outlined in the pioneering work of Novikov [**171**].

The calculation of the oriented bordism ring was completed by Novikov [**169**] (ring structure modulo torsion and odd torsion) and Wall [**227**] (even torsion), with important earlier contributions made by Rokhlin, Averbuch, and Milnor. Unlike complex bordism, the ring Ω^{SO}_* has additive torsion. We give only a partial result here (which does not fully describe the torsion elements).

---

[1] Milnor's proof was announced in [**121**]; it was intended to be included in the second part of [**161**], but has never been published.

Theorem E.4.3.

(a) *$\Omega_*^{SO} \otimes \mathbb{Q}$ is a polynomial ring over $\mathbb{Q}$ generated by the bordism classes of complex projective spaces $\mathbb{C}P^{2i}$, $i \geqslant 1$.*

(b) *The subring* Tors $\subset \Omega_*^{SO}$ *of torsion elements contains only elements of order* 2. *The quotient $\Omega_*^{SO}/\mathrm{Tors}$ is a polynomial ring over $\mathbb{Z}$ with one generator a_i in every dimension $4i$, where $i \geqslant 1$.*

(c) *Two oriented manifolds are bordant if and only if they have identical sets of Pontrjagin and Stiefel–Whitney characteristic numbers.*

E.5. Multiplicative generators

To describe a set of multiplicative generators for the ring Ω_*^U we shall need a special characteristic class of complex vector bundles. Let ξ be a complex k-plane bundle over a manifold M. Write its total Chern class formally as follows:

$$c(\xi) = 1 + c_1(\xi) + \cdots + c_k(\xi) = (1 + x_1)\cdots(1 + x_k),$$

so that $c_i(\xi) = \sigma_i(x_1, \ldots, x_k)$ is the ith elementary symmetric function in formal indeterminates. These indeterminates acquire a geometric meaning if ξ is a sum $\xi_1 \oplus \ldots \oplus \xi_k$ of line bundles; then $x_j = c_1(\xi_j)$, $1 \leqslant j \leqslant k$. Consider the polynomial

$$P_n(x_1, \ldots x_k) = x_1^n + \cdots + x_k^n$$

and express it via the elementary symmetric functions:

$$P_n(x_1, \ldots, x_k) = s_n(\sigma_1, \ldots, \sigma_k).$$

Substituting the Chern classes for the elementary symmetric functions we obtain a certain characteristic class of ξ:

$$s_n(\xi) = s_n(c_1(\xi), \ldots, c_k(\xi)) \in H^{2n}(M).$$

This characteristic class plays an important role in detecting the polynomial generators of the complex bordism ring, because of the following properties (which follow immediately from the definition).

Proposition E.5.1.

1. $s_n(\xi) = 0$ *for* $2n > \dim M$.
2. $s_n(\xi \oplus \eta) = s_n(\xi) + s_n(\eta)$.

Given a stably complex manifold $(M, c_{\mathcal{T}})$ of dimension $2n$, define its characteristic number

$$s_n[M] = s_n(\xi)\langle M\rangle \in \mathbb{Z}$$

where ξ is the complex bundle from (E.6), and $\langle M\rangle \in H_{2n}(M)$ the fundamental homology class.

Corollary E.5.2. *If a bordism class $[M] \in \Omega_{2n}^U$ decomposes as $[M_1] \times [M_2]$ where $\dim M_1 > 0$ and $\dim M_2 > 0$, then $s_n[M] = 0$.*

It follows that the characteristic number s_n vanishes on decomposable elements of Ω_{2n}^U. It also detects indecomposables that may be chosen as polynomial generators. The following result featured in the proof of Theorem E.4.2.

THEOREM E.5.3. *A bordism class* $[M] \in \varOmega^U_{2n}$ *may be chosen as a polynomial generator* a_n *of the ring* $\varOmega^U_*$ *if and only if*

$$s_n[M] = \begin{cases} \pm 1, & \text{if } n \neq p^k - 1 \text{ for any prime } p; \\ \pm p, & \text{if } n = p^k - 1 \text{ for some prime } p. \end{cases}$$

There is no universal description of connected manifolds representing the polynomial generators $a_n \in \varOmega^U_*$. On the other hand, there is a particularly nice family of manifolds whose bordism classes generate the whole ring $\varOmega^U_*$. This family is redundant though, so there are algebraic relations between their bordism classes.

CONSTRUCTION E.5.4 (Milnor hypersurfaces). Fix a pair of integers $j \geqslant i \geqslant 0$ and consider the product $\mathbb{C}P^i \times \mathbb{C}P^j$. Its algebraic subvariety

$$H_{ij} = \{(z_0 : \cdots : z_i) \times (w_0 : \cdots : w_j) \in \mathbb{C}P^i \times \mathbb{C}P^j : \\ z_0 w_0 + \cdots + z_i w_i = 0\} \tag{E.8}$$

is called a *Milnor hypersurface*. Note that $H_{0j} \cong \mathbb{C}P^{j-1}$.

Denote by p_1 and p_2 the projections of $\mathbb{C}P^i \times \mathbb{C}P^j$ onto its factors, and by η the Hopf line bundle over a complex projective space; then $\bar{\eta}$ is the hyperplane section bundle (it is usually denoted by $\mathcal{O}(1)$ in the algebraic geometry literature). We have

$$H^*(\mathbb{C}P^i \times \mathbb{C}P^j) = \mathbb{Z}[x, y]/(x^{i+1} = 0,\ y^{j+1} = 0)$$

where $x = p_1^* c_1(\bar{\eta})$, $y = p_2^* c_1(\bar{\eta})$.

PROPOSITION E.5.5. *The geometric cobordism in* $\mathbb{C}P^i \times \mathbb{C}P^j$ *corresponding to the element* $x + y \in H^2(\mathbb{C}P^i \times \mathbb{C}P^j)$ *is represented by the submanifold* H_{ij}. *In particular, the image of the fundamental class* $\langle H_{ij} \rangle$ *in* $H_{2(i+j-1)}(\mathbb{C}P^i \times \mathbb{C}P^j)$ *is Poincaré dual to* $x + y$.

PROOF. We have $x + y = c_1(p_1^*(\bar{\eta}) \otimes p_2^*(\bar{\eta}))$. The classifying map $f_{x+y} \colon \mathbb{C}P^i \times \mathbb{C}P^j \to \mathbb{C}P^\infty$ is the composition of the *Segre embedding*

$$\sigma \colon \mathbb{C}P^i \times \mathbb{C}P^j \to \mathbb{C}P^{ij+i+j},$$
$$(z_0 : \cdots : z_i) \times (w_0 : \cdots : w_j) \mapsto (z_0 w_0 : z_0 w_1 : \cdots : z_k w_l : \cdots : z_i w_j),$$

and the embedding $\mathbb{C}P^{ij+i+j} \to \mathbb{C}P^\infty$. The codimension 2 submanifold in $\mathbb{C}P^i \times \mathbb{C}P^j$ corresponding to the cohomology class $x + y$ is obtained as the inverse image $\sigma^{-1}(H)$ of a generally positioned hyperplane in $\mathbb{C}P^{ij+i+j}$ (that is, a hyperplane H transverse to the image of the Segre embedding, see Construction E.3.4). By (E.8), the Milnor hypersurface is exactly $\sigma^{-1}(H)$ for one such hyperplane H. □

LEMMA E.5.6.

$$s_{i+j-1}[H_{ij}] = \begin{cases} j, & \text{if } \ i = 0, \text{ that is, } H_{0j} = \mathbb{C}P^{j-1}; \\ 2, & \text{if } \ i = j = 1; \\ 0, & \text{if } \ i = 1,\ j > 1; \\ -\binom{i+j}{i}, & \text{if } \ i > 1. \end{cases}$$

PROOF. Let $i = 0$. Since the stably complex structure on $H_{0j} = \mathbb{C}P^{j-1}$ is determined by the isomorphism $\mathcal{T}(\mathbb{C}P^{j-1}) \oplus \mathbb{C} \cong \bar{\eta} \oplus \cdots \oplus \bar{\eta}$ (j summands) and $x = c_1(\bar{\eta})$, we have

$$s_{j-1}[\mathbb{C}P^{j-1}] = j x^{j-1} \langle \mathbb{C}P^{j-1} \rangle = j.$$

Now let $i > 0$. Then

$$s_{i+j-1}(\mathcal{T}(\mathbb{C}P^i \times \mathbb{C}P^j)) = (i+1)x^{i+j-1} + (j+1)y^{i+j-1} = \begin{cases} 2x^j + (j+1)y^j, & \text{if } i = 1; \\ 0, & \text{if } i > 1. \end{cases}$$

Denote by ν the normal bundle of the embedding $\iota\colon H_{ij} \to \mathbb{C}P^i \times \mathbb{C}P^j$. Then

$$\mathcal{T}(H_{ij}) \oplus \nu = \iota^*(\mathcal{T}(\mathbb{C}P^i \times \mathbb{C}P^j)). \tag{E.9}$$

Since $c_1(\nu) = \iota^*(x+y)$, we obtain $s_{i+j-1}(\nu) = \iota^*(x+y)^{i+j-1}$.

Assume $i = 1$. Then (E.9) and Proposition E.5.1 imply that

$$\begin{aligned} s_j[H_{1j}] &= s_j\big(\mathcal{T}(H_{1j})\big)\langle H_{1j}\rangle = \iota^*\big(2x^j + (j+1)y^j - (x+y)^j\big)\langle H_{1j}\rangle \\ &= (2x^j + (j+1)y^j - (x+y)^j)(x+y)\langle \mathbb{C}P^1 \times \mathbb{C}P^j\rangle = \begin{cases} 2, & \text{if } j = 1; \\ 0, & \text{if } j > 1. \end{cases} \end{aligned}$$

Assume now that $i > 1$. Then $s_{i+j-1}(\mathcal{T}(\mathbb{C}P^i \times \mathbb{C}P^j)) = 0$, and we obtain from (E.9) and Proposition E.5.1 that

$$\begin{aligned} s_{i+j-1}[H_{ij}] &= -s_{i+j-1}(\nu)\langle H_{ij}\rangle = -\iota^*(x+y)^{i+j-1}\langle H_{ij}\rangle \\ &= -(x+y)^{i+j}\langle \mathbb{C}P^i \times \mathbb{C}P^j\rangle = -\tbinom{i+j}{i}. \end{aligned}$$

□

REMARK. Since $s_1[H_{11}] = 2 = s_1[\mathbb{C}P^1]$ the manifold H_{11} is bordant to $\mathbb{C}P^1$. In fact $H_{11} \cong \mathbb{C}P^1$ (an exercise).

THEOREM E.5.7. *The bordism classes* $\{[H_{ij}], 0 \leqslant i \leqslant j\}$ *multiplicatively generate the ring* Ω^U_*.

PROOF. A simple calculation shows that

$$\text{g.c.d.}\Big(\tbinom{n+1}{i},\ 1 \leqslant i \leqslant n\Big) = \begin{cases} p, & \text{if } n = p^k - 1, \\ 1, & \text{otherwise.} \end{cases}$$

Now Lemma E.5.6 implies that a certain integer linear combination of bordism classes $[H_{ij}]$ with $i + j = n + 1$ can be taken as the polynomial generator a_n of Ω^U_*, see Theorem E.5.3. □

REMARK. There is no universal description for a linear combination of bordism classes $[H_{ij}]$ with $i + j = n + 1$ giving the polynomial generator of Ω^U_*.

EXAMPLE E.5.8. Since $s_1[\mathbb{C}P^1] = 2$, $s_2[\mathbb{C}P^2] = 3$, the bordism classes $[\mathbb{C}P^1]$ and $[\mathbb{C}P^2]$ may be taken as polynomial generators a_1 and a_2 of Ω^U_*. However $[\mathbb{C}P^3]$ cannot be taken as a_3, since $s_3[\mathbb{C}P^3] = 4$, while $s_3(a_3) = \pm 2$. The bordism class $[H_{22}] + [\mathbb{C}P^3]$ may be taken as a_3.

Theorem E.5.7 admits the following important addendum, which is due to Milnor (see [**212**, Chapter 7] for the proof).

THEOREM E.5.9 (Milnor). *Every bordism class* $x \in \Omega^U_n$ *with* $n > 0$ *contains a nonsingular algebraic variety (not necessarily connected).*

The following question is still open.

Problem E.5.10 (Hirzebruch). Describe the set of bordism classes in $\varOmega_*^U$ containing connected nonsingular algebraic varieties.

Example E.5.11. The group $\varOmega_2^U$ is isomorphic to $\mathbb{Z}$ and is generated by $[\mathbb{C}P^1]$. Every class $k[\mathbb{C}P^1] \in \varOmega_2^U$ contains a nonsingular algebraic variety, namely, a disjoint union of k copies of $\mathbb{C}P^1$ for $k > 0$ and a Riemann surface of genus $(1-k)$ for $k \leqslant 0$. Connected algebraic varieties are contained only in the bordism classes $k[\mathbb{C}P^1]$ with $k \leqslant 1$.

Exercises.

E.5.12. Show that $H_{11} \cong \mathbb{C}P^1$.

E.6. Invariant stably complex structures

Let M be a $2n$-dimensional manifold with a stably complex structure determined by the isomorphism

$$c_{\mathcal{T}} \colon \mathcal{T}M \oplus \underline{\mathbb{R}}^{2(l-n)} \to \xi. \tag{E.10}$$

Assume that the torus $\mathbb{T}^k$ acts on M.

Definition E.6.1. A stably complex structure $c_{\mathcal{T}}$ is $\mathbb{T}^k$-*invariant* if for every $\boldsymbol{t} \in \mathbb{T}^k$ the composition

$$r(\boldsymbol{t}) \colon \xi \xrightarrow{c_{\mathcal{T}}^{-1}} \mathcal{T}M \oplus \underline{\mathbb{R}}^{2(l-n)} \xrightarrow{d\boldsymbol{t} \oplus \mathrm{id}} \mathcal{T}M \oplus \underline{\mathbb{R}}^{2(l-n)} \xrightarrow{c_{\mathcal{T}}} \xi \tag{E.11}$$

is a complex bundle map, where $d\boldsymbol{t}$ is the differential of the action by $\boldsymbol{t}$. In other words, (E.11) determines a representation $r \colon \mathbb{T}^k \to \mathrm{Hom}_{\mathbb{C}}(\xi, \xi)$.

Let $x \in M$ be an isolated fixed point of the $\mathbb{T}^k$-action on M. Then we have a representation $r_x \colon \mathbb{T}^k \to GL(l, \mathbb{C})$ in the fibre of ξ over x. This fibre $\xi_x \cong \mathbb{C}^l$ decomposes as $\mathbb{C}^n \oplus \mathbb{C}^{l-n}$, where r_x has no trivial summands on $\mathbb{C}^n$ and is trivial on $\mathbb{C}^{l-n}$. Also, the isomorphism $c_{\mathcal{T},x}$ of (E.10) induces an orientation of the tangent space $\mathcal{T}_x(M)$.

Definition E.6.2. For any fixed point $x \in M$, the *sign* $\sigma(x)$ is $+1$ if the isomorphism

$$\mathcal{T}_x(M) \xrightarrow{\mathrm{id} \oplus 0} \mathcal{T}_x(M) \oplus \mathbb{R}^{2(l-n)} \xrightarrow{c_{\mathcal{T},x}} \xi_x \cong \mathbb{C}^n \oplus \mathbb{C}^{l-n} \xrightarrow{p} \mathbb{C}^n,$$

respects the canonical orientations, and -1 if it does not; here p is the projection onto the first summand.

So $\sigma(x)$ compares the orientations induced by r_x and $c_{\mathcal{T},x}$ on $\mathcal{T}_x(M)$. If M is an almost complex $\mathbb{T}^k$-manifold (i.e., $l = n$) then $\sigma(x) = 1$ for every fixed point x.

Example E.6.3. We let the circle $\mathbb{T} = S^1$ act on $\mathbb{C}P^1$ in homogeneous coordinates by $t \cdot (z_0 : z_1) = (z_0 : tz_1)$. This action has two fixed points $(0 : 1)$ and $(1 : 0)$. In the standard stably complex structure (see Example E.3.1) the signs of both vertices are positive (this structure is complex and the map in Definition E.6.2 is a complex linear map $\mathbb{C} \to \mathbb{C}$).

On the other hand, the trivial stably complex structure on $\mathbb{C}P^1 \cong S^2$ may be thought of as induced from the embedding of S^2 into $\mathbb{R}^3 \oplus \mathbb{R}^1$ with trivial normal bundle, which can be made complex by identifying it with $\underline{\mathbb{C}}$. We therefore may think of the circle action as the rotation of the unit sphere in $\mathbb{R}^3$ around the vertical axis. The fixed points are the north and south poles. The rotations induced on the tangent planes to the two fixed points are in different directions. Therefore the signs of the two fixed points are different (which sign is positive depends on the convention one uses to induce an orientation on S^2 from the orientation of $\mathbb{R}^3$).

APPENDIX F

Formal group laws and Hirzebruch genera

The theory of *formal groups* originally appeared in algebraic geometry and plays an important role in number theory and cryptography. Formal groups laws were brought into bordism theory in the pioneering work of Novikov [**171**], and provided a very powerful tool for the theory of group actions on manifolds and generalised homology theories. Early applications of formal group laws in cobordism concerned finite group actions on manifolds, or 'differentiable periodic maps'. Subsequent developments included constructions of complex oriented cohomology theories and applications to *Hirzebruch genera*, one of the most important class of invariants of manifolds.

F.1. Elements of the theory of formal group laws

Let R be a commutative ring with unit.

A formal power series $F(u,v) \in R[[u,v]]$ is called a (commutative one-dimensional) *formal group law* over R if it satisfies the following equations:

(a) $F(u,0)=u$, $F(0,v)=v$;
(b) $F(F(u,v),w)=F(u,F(v,w))$;
(c) $F(u,v)=F(v,u)$.

The original example of a formal group law over a field $\mathbf{k}$ is provided by the expansion near the unit of the multiplication map $G\times G\to G$ in a one-dimensional algebraic group over $\mathbf{k}$. This also explains the terminology.

A formal group law F over R is called *linearisable* if there exists a coordinate change $u\mapsto g_F(u)=u+\sum_{i>1}g_iu^i\in R[[u]]$ such that

$$g_F(F(u,v))=g_F(u)+g_F(v). \tag{F.1}$$

Note that every formal group law over R determines a formal group law over $R\otimes\mathbb{Q}$.

THEOREM F.1.1. *Every formal group law F is linearisable over $R\otimes\mathbb{Q}$.*

PROOF. Consider the series $\omega(u)=\frac{\partial F(u,w)}{\partial w}\Big|_{w=0}$. Then

$$\omega(F(u,v))=\frac{\partial F(F(u,v),w)}{\partial w}\Big|_{w=0}=\frac{\partial F(F(u,w),v)}{\partial F(u,w)}\cdot\frac{\partial F(u,w)}{\partial w}\Big|_{w=0}=\frac{\partial F(u,v)}{\partial u}\omega(u).$$

We therefore have $\frac{du}{\omega(u)}=\frac{dF(u,v)}{\omega(F(u,v))}$. Set

$$g(u)=\int_0^u\frac{dv}{\omega(v)}; \tag{F.2}$$

then $dg(u)=dg(F(u,v))$. This implies that $g(F(u,v))=g(u)+C$. Since $F(0,v)=v$ and $g(0)=0$, we get $C=g(v)$. Thus, $g(F(u,v))=g(u)+g(v)$. □

A series $g_F(u) = u + \sum_{i>1} g_i u^i$ satisfying equation (F.1) is called a *logarithm* of the formal group law F; Theorem F.1.1 shows that a formal group law over $R \otimes \mathbb{Q}$ always has a logarithm. Its functional inverse series $f_F(t) \in R \otimes \mathbb{Q}[[t]]$ is called an *exponential* of the formal group law, so that we have $F(u,v) = f_F(g_F(u) + g_F(v))$ over $R \otimes \mathbb{Q}$. If R does not have torsion (i.e. $R \to R \otimes \mathbb{Q}$ is monomorphic), the latter formula shows that a formal group law (as a series with coefficients in R) is fully determined by its logarithm (which is a series with coefficients in $R \otimes \mathbb{Q}$).

EXAMPLE F.1.2. An example of a formal group law is given by the series

$$F(u,v) = (1+u)(1+v) - 1 = u + v + uv, \tag{F.3}$$

over $\mathbb{Z}$, called the *multiplicative formal group law.* Introducing a formal indeterminate β of degree -2, we may consider the 1-parameter extension of the multiplicative formal group law, given by $F_\beta = u + v - \beta uv$, with coefficients in $\mathbb{Z}[\beta]$. Its logarithm and exponential series are given by

$$g(u) = -\frac{\ln(1-\beta u)}{\beta}, \quad f(x) = \frac{1 - e^{-\beta x}}{\beta}.$$

Let $F = \sum_{k,l} a_{kl} u^k v^l$ be a formal group law over a ring R and $r\colon R \to R'$ a ring homomorphism. Denote by $r(F)$ the formal series $\sum_{k,l} r(a_{kl}) u^k v^l \in R'[[u,v]]$; then $r(F)$ is a formal group law over R'.

A formal group law $\mathcal{F}$ over a ring A is *universal* if for any formal group law F over any ring R there exists a unique homomorphism $r\colon A \to R$ such that $F = r(\mathcal{F})$.

PROPOSITION F.1.3. *Assume that a universal formal group law $\mathcal{F}$ over A exists. Then*

(a) *the ring A is multiplicatively generated by the coefficients of the series $\mathcal{F}$;*
(b) *the universal formal group law is unique: if $\mathcal{F}'$ is another universal formal group law over A', then there is an isomorphism $r\colon A \to A'$ such that $\mathcal{F}' = r(\mathcal{F})$.*

PROOF. To prove the first statement, denote by A' the subring in A generated by the coefficients of $\mathcal{F}$. Then there is a monomorphism $i\colon A' \to A$ satisfying $i(\mathcal{F}) = \mathcal{F}$. On the other hand, by universality there exists a homomorphism $r\colon A \to A'$ satisfying $r(\mathcal{F}) = \mathcal{F}$. It follows that $ir(\mathcal{F}) = \mathcal{F}$. This implies that $ir = \mathrm{id}\colon A \to A$ by the uniqueness requirement in the definition of $\mathcal{F}$. Thus $A' = A$. The second statement is proved similarly. □

THEOREM F.1.4 (Lazard [**139**]). *The universal formal group law $\mathcal{F}$ exists, and its coefficient ring A is isomorphic to the polynomial ring $\mathbb{Z}[a_1, a_2, \ldots]$ on an infinite number of generators.*

F.2. Formal group law of geometric cobordisms

The applications of formal group laws in cobordism theory build upon the following basic example.

CONSTRUCTION F.2.1 (Formal group law of geometric cobordisms). Let X be a cell complex and $u, v \in U^2(X)$ two geometric cobordisms (see Construction E.3.4) corresponding to elements $x, y \in H^2(X)$ respectively. Denote by $u +_{H} v$ the geometric cobordism corresponding to the cohomology class $x + y$.

PROPOSITION F.2.2. *The following relation holds in* $U^2(X)$*:*

$$u +_{H} v = F_U(u,v) = u + v + \sum_{k\geqslant 1,\, l\geqslant 1} \alpha_{kl}\, u^k v^l, \tag{F.4}$$

where the coefficients $\alpha_{kl} \in \varOmega_U^{-2(k+l-1)}$ *do not depend on* X*. The series* $F_U(u,v)$ *given by* (F.4) *is a formal group law over the ring* $\varOmega_U = \varOmega_U^*$*.*

PROOF. We first do calculations with the universal example $X = \mathbb{C}P^\infty \times \mathbb{C}P^\infty$. Then

$$U^*(\mathbb{C}P^\infty \times \mathbb{C}P^\infty) = \varOmega_U^*[[\underline{u}, \underline{v}]],$$

where $\underline{u}, \underline{v}$ are canonical geometric cobordisms given by the projections of $\mathbb{C}P^\infty \times \mathbb{C}P^\infty$ onto its factors. We therefore have the following relation in $U^2(\mathbb{C}P^\infty \times \mathbb{C}P^\infty)$:

$$\underline{u} +_{H} \underline{v} = \sum_{k,l\geqslant 0} \alpha_{kl}\, \underline{u}^k \underline{v}^l, \tag{F.5}$$

where $\alpha_{kl} \in \varOmega_U^{-2(k+l-1)}$.

Now let the geometric cobordisms $u, v \in U^2(X)$ be given by maps $f_u, f_v\colon X \to \mathbb{C}P^\infty$ respectively. Then $u = (f_u \times f_v)^*(\underline{u})$, $v = (f_u \times f_v)^*(\underline{v})$ and $u +_{H} v = (f_u \times f_v)^*(\underline{u} +_{H} \underline{v})$, where $f_u \times f_v\colon X \to \mathbb{C}P^\infty \times \mathbb{C}P^\infty$. Applying the $\varOmega_U^*$-module map $(f_u \times f_v)^*$ to (F.5) we obtain the required formula (F.4). The fact that $F_U(u,v)$ is a formal group law follows directly from the properties of the group multiplication $\mathbb{C}P^\infty \times \mathbb{C}P^\infty \to \mathbb{C}P^\infty$. □

Series (F.4) is called the *formal group law of geometric cobordisms*; nowadays it is also usually referred to as the *complex cobordism formal group law.*

By definition, the geometric cobordism $u \in U^2(X)$ is the first *Conner–Floyd Chern class* of the complex line bundle ξ over X obtained by pulling back the canonical bundle along the map $f_u\colon X \to \mathbb{C}P^\infty$. It follows that the formal group law of geometric cobordisms gives an expression of the first class $c_1^U(\xi \otimes \eta) \in U^2(X)$ of the tensor product of two complex line bundles over X in terms of the classes $u = c_1^U(\xi)$ and $v = c_1^U(\eta)$ of the factors:

$$c_1^U(\xi \otimes \eta) = F_U(u,v).$$

The coefficients of the formal group law of geometric cobordisms and its logarithms may be described geometrically by the following results.

THEOREM F.2.3 (Buchstaber [**37**, Theorem 4.8]).

$$F_U(u,v) = \frac{\sum_{i,j\geqslant 0}[H_{ij}]u^i v^j}{\bigl(\sum_{r\geqslant 0}[\mathbb{C}P^r]u^r\bigr)\bigl(\sum_{s\geqslant 0}[\mathbb{C}P^s]v^s\bigr)},$$

where H_{ij} $(0 \leqslant i \leqslant j)$ *are Milnor hypersurfaces* (E.8) *and* $H_{ji} = H_{ij}$*.*

PROOF. Set $X = \mathbb{C}P^i \times \mathbb{C}P^j$ in Proposition F.2.2. Consider the Poincaré–Atiyah duality map $D\colon U^2(\mathbb{C}P^i \times \mathbb{C}P^j) \to U_{2(i+j)-2}(\mathbb{C}P^i \times \mathbb{C}P^j)$ (see Exercise E.3.7) and the map $\varepsilon\colon U_*(\mathbb{C}P^i \times \mathbb{C}P^j) \to U_*(pt) = \varOmega_*^U$ induced by the projection $\mathbb{C}P^i \times \mathbb{C}P^j \to pt$. Then the composition

$$\varepsilon D\colon U^2(\mathbb{C}P^i \times \mathbb{C}P^j) \to \varOmega_{2(i+j)-2}^U$$

takes geometric cobordisms to the bordism classes of the corresponding submanifolds. In particular, $\varepsilon D(u +_{H} v) = [H_{ij}]$, $\varepsilon D(u^k v^l) = [\mathbb{C}P^{i-k}][\mathbb{C}P^{j-l}]$. Applying εD to (F.4) we obtain

$$[H_{ij}] = \sum_{k,\,l} \alpha_{kl} [\mathbb{C}P^{i-k}][\mathbb{C}P^{j-l}].$$

Therefore,

$$\sum_{i,j} [H_{ij}] u^i v^j = \Bigl(\sum_{k,\,l} \alpha_{kl} u^k v^l\Bigr)\Bigl(\sum_{i\geqslant k} [\mathbb{C}P^{i-k}] u^{i-k}\Bigr)\Bigl(\sum_{j\geqslant l} [\mathbb{C}P^{j-l}] v^{j-l}\Bigr),$$

which implies the required formula. □

THEOREM F.2.4 (Mishchenko [**171**, Appendix 1]). *The logarithm of the formal group law of geometric cobordisms is given by*

$$g_U(u) = u + \sum_{k\geqslant 1} \frac{[\mathbb{C}P^k]}{k+1} u^{k+1} \in \varOmega_U \otimes \mathbb{Q}[[u]].$$

PROOF. By (F.2),

$$dg_U(u) = \frac{du}{\frac{\partial F_U(u,v)}{\partial v}\Big|_{v=0}}.$$

Using the formula of Theorem F.2.3 and the identity $H_{i0} = \mathbb{C}P^{i-1}$, we calculate

$$\frac{dg_U(u)}{du} = \frac{1 + \sum_{k>0} [\mathbb{C}P^k] u^k}{1 + \sum_{i>0} ([H_{i1}] - [\mathbb{C}P^1][\mathbb{C}P^{i-1}]) u^i}.$$

Now $[H_{i1}] = [\mathbb{C}P^1][\mathbb{C}P^{i-1}]$ (see Exercise F.3.6; we have already seen that $H_{11} = \mathbb{C}P^1$ in the Remark preceding Theorem E.5.7). Therefore, $dg_U(u) = 1 + \sum_{k>0} [\mathbb{C}P^k] u^k$, which implies the required formula. □

Using these calculations the following most important property of the formal group law F_U can be easily established:

THEOREM F.2.5 (Quillen [**192**, Theorem 2]). *The formal group law F_U of geometric cobordisms is universal.*

PROOF. Let $\mathcal{F}$ be the universal formal group law over a ring A. Then there is a homomorphism $r\colon A \to \varOmega_U$ which takes $\mathcal{F}$ to F_U. The series $\mathcal{F}$, viewed as a formal group law over the ring $A \otimes \mathbb{Q}$, has the universality property for all formal group laws over $\mathbb{Q}$-algebras. By Theorem F.1.1, such a formal group law is determined by its logarithm, which is a series with leading term u. It follows that if we write the logarithm of $\mathcal{F}$ as $\sum b_k \frac{u^{k+1}}{k+1}$ then the ring $A \otimes \mathbb{Q}$ is the polynomial ring $\mathbb{Q}[b_1, b_2, \ldots]$. By Theorem F.2.4, $r(b_k) = [\mathbb{C}P^k] \in \varOmega_U$. Since $\varOmega_U \otimes \mathbb{Q} \cong \mathbb{Q}[[\mathbb{C}P^1], [\mathbb{C}P^2], \ldots]$, this implies that $r \otimes \mathbb{Q}$ is an isomorphism.

By Theorem F.1.4 the ring A does not have torsion, so r is a monomorphism. On the other hand, Theorem F.2.3 implies that the image $r(A)$ contains the bordism classes $[H_{ij}] \in \varOmega_U$, $0 \leqslant i \leqslant j$. Since these classes generate the whole ring $\varOmega_U$ (Theorem E.5.7), the map r is onto and thus an isomorphism. □

F.3. Hirzebruch genera

Every homomorphism $\varphi\colon \varOmega_U \to R$ from the complex cobordism ring to a commutative ring R with unit can be regarded as a multiplicative characteristic of manifolds which is an invariant of cobordism classes. Such a homomorphism is called a (complex) *R-genus*. (The term *'multiplicative genus'* is also used, to emphasise that such a genus is a ring homomorphism; in classical algebraic geometry, there are instances of genera which are not multiplicative.)

Assume that the ring R does not have additive torsion. Then every R-genus φ is fully determined by the corresponding homomorphism $\varOmega_U \otimes \mathbb{Q} \to R \otimes \mathbb{Q}$, which we shall also denote by φ. A famous construction due to Hirzebruch [**122**] allows us to describe homomorphisms $\varphi\colon \varOmega_U \otimes \mathbb{Q} \to R \otimes \mathbb{Q}$ by means of universal R-valued characteristic classes of special type.

CONSTRUCTION F.3.1 (Hirzebruch genera). Let $BU = \lim\limits_{n\to\infty} BU(n)$. Then $H^*(BU)$ is isomorphic to the graded ring of formal power series $\mathbb{Z}[[c_1, c_2, \ldots]]$ in universal Chern classes, $\deg c_k = 2k$. The set of Chern characteristic numbers of a manifold M defines an element in $\mathrm{Hom}(H^*(BU), \mathbb{Z})$, which in fact belongs to the subgroup $H_*(BU)$ in the latter group. We therefore obtain a group homomorphism

$$\varOmega_U \to H_*(BU). \tag{F.6}$$

Since the multiplication in the ring $H_*(BU)$ is obtained from the maps $BU_k \times BU_l \to BU_{k+l}$ corresponding to the Whitney sum of vector bundles, and the Chern classes have the appropriate multiplicative property, the map (F.6) is a ring homomorphism.

Part 2 of Theorem E.4.2 says that (F.6) is a monomorphism, and Part 1 of the same theorem says that the map $\varOmega_U \otimes \mathbb{Q} \to H_*(BU; \mathbb{Q})$ is an isomorphism. It follows that every homomorphism $\varphi\colon \varOmega_U \otimes \mathbb{Q} \to R \otimes \mathbb{Q}$ can be interpreted as an element of

$$\mathrm{Hom}_{\mathbb{Q}}(H_*(BU; \mathbb{Q}), R \otimes \mathbb{Q}) = H^*(BU; \mathbb{Q}) \otimes R,$$

or as a sequence of homogeneous polynomials $\{K_i(c_1, \ldots, c_i),\ i \geqslant 0\}$, $\deg K_i = 2i$. This sequence of polynomials cannot be chosen arbitrarily; the fact that φ is a ring homomorphism imposes certain conditions. These conditions may be described as follows: an identity

$$1 + c_1 + c_2 + \cdots = (1 + c_1' + c_2' + \cdots) \cdot (1 + c_1'' + c_2'' + \cdots)$$

implies the identity

$$\sum_{n\geqslant 0} K_n(c_1, \ldots, c_n) = \sum_{i\geqslant 0} K_i(c_1', \ldots, c_i') \cdot \sum_{j\geqslant 0} K_j(c_1'', \ldots, c_j''). \tag{F.7}$$

A sequence of homogeneous polynomials $\mathcal{K} = \{K_i(c_1, \ldots, c_i), i \geqslant 0\}$ with $K_0 = 1$ satisfying the identities (F.7) is called a *multiplicative Hirzebruch sequence.*

PROPOSITION F.3.2. *A multiplicative sequence $\mathcal{K}$ is completely determined by the series*

$$Q(x) = 1 + q_1 x + q_2 x^2 + \cdots \in R \otimes \mathbb{Q}[[x]],$$

where $x = c_1$, and $q_i = K_i(1, 0, \ldots, 0)$; moreover, every series $Q(x)$ as above determines a multiplicative sequence.

Proof. Indeed, by considering the identity

$$1 + c_1 + \cdots + c_n = (1 + x_1) \cdots (1 + x_n) \tag{F.8}$$

we obtain from (F.7) that

$$Q(x_1) \cdots Q(x_n) = 1 + K_1(c_1) + K_2(c_1, c_2) + \cdots \\ + K_n(c_1, \ldots, c_n) + K_{n+1}(c_1, \ldots, c_n, 0) + \cdots .$$

□

Along with the series $Q(x)$ it is convenient to consider the series $f(x) \in R \otimes \mathbb{Q}[[x]]$ with leading term x given by the identity

$$Q(x) = \frac{x}{f(x)}.$$

It follows that ring homomorphisms $\varphi\colon \varOmega_U \otimes \mathbb{Q} \to R \otimes \mathbb{Q}$ are in one-to-one correspondence with series $f(x) \in R \otimes \mathbb{Q}[[x]]$ with leading term x. Under this correspondence, the value of φ on an $2n$-dimensional bordism class $[M] \in \varOmega_U$ is given by

$$\varphi[M] = \Bigl(\prod_{i=1}^{n} \frac{x_i}{f(x_i)}, \langle M \rangle \Bigr), \tag{F.9}$$

where the Chern classes $c_1, \ldots, c_n$ are expressed via the indeterminates $x_1, \ldots, x_n$ by the relation (F.8). We shall also denote the characteristic class $\prod_{i=1}^{n} \frac{x_i}{f(x_i)}$ of a complex vector bundle ξ by $\varphi(\xi)$; so that $\varphi[M] = \varphi(\mathcal{T}M)\langle M \rangle$.

The homomorphism $\varphi\colon \varOmega_U \to R \otimes \mathbb{Q}$ given by formula (F.9) is called the *Hirzebruch genus* associated with the series $f(x) = x + \cdots \in R \otimes \mathbb{Q}[[x]]$. Thus, there is a one-two-one correspondence between series $f(x) \in R \otimes \mathbb{Q}[[x]]$ having leading term x and genera $\varphi\colon \varOmega_U \to R \otimes \mathbb{Q}$.

Every genus $\varphi\colon \varOmega_U \to R$ gives rise to a formal group law $\varphi(F_U)$ over R, where F_U is the formal group of geometric cobordisms (Construction F.2.1).

Theorem F.3.3. *For every genus $\varphi\colon \varOmega_U \to R \otimes \mathbb{Q}$, the exponential of the formal group law $\varphi(F_U)$ is given by the series $f(x) \in R \otimes \mathbb{Q}[[x]]$ corresponding to φ.*

This can be proved either directly, by appealing to the construction of geometric cobordisms, or indirectly, by calculating the values of the give Hirzebruch genus on projective spaces and comparing to the formula for the logarithm of the formal group law.

1st proof. Let X be a manifold and $u, v \in U^2(X)$ its two geometric cobordisms defined by the elements $x, y \in H^2(X)$ respectively. By definition (F.4) of the formal group law F_U we have the following relation between geometric cobordisms in $U^2(X)$:

$$[M_{x+y}] = \sum_{k,l \geqslant 0} \alpha_{kl} [M_{x^k y^l}]$$

in $\varOmega_U$, where $M_{x+y} \subset X$ is the codimension-2 submanifold dual to $x + y \in H^2(X)$, and $M_{x^k y^l} \subset X$ is a codimension-$2(k+l)$ submanifold dual to $x^k y^l \in H^2(X)$. Applying the genus φ we obtain

$$\varphi[M_{x+y}] = \sum \varphi(\alpha_{kl}) \varphi[M_{x^k y^l}]. \tag{F.10}$$

Let $\iota\colon M_{x+y} \subset X$ be the embedding. Considering the decomposition

$$\iota^*(\mathcal{T}X) = \mathcal{T}M_{x+y} \oplus \nu(\iota)$$

and using the multiplicativity of the characteristic class φ we obtain

$$\iota^*\varphi(\mathcal{T}X) = \varphi(\mathcal{T}M_{x+y}) \cdot \iota^*\left(\tfrac{x+y}{f(x+y)}\right).$$

Therefore,

$$\varphi[M_{x+y}] = \iota^*\big(\varphi(\mathcal{T}X) \cdot \tfrac{f(x+y)}{x+y}\big)\langle M_{x+y}\rangle = \big(\varphi(\mathcal{T}X) \cdot f(x+y)\big)\langle X\rangle. \tag{F.11}$$

Similarly, by considering the embedding $M_{x^k y^l} \to X$ we obtain

$$\varphi[M_{x^k y^l}] = \big(\varphi(\mathcal{T}X) \cdot f(x)^k f(y)^l)\big)\langle X\rangle. \tag{F.12}$$

Plugging (F.11) and (F.12) into (F.10) we finally obtain

$$f(x+y) = \sum_{k,l\geqslant 0} \varphi(\alpha_{kl}) f(x)^k f(y)^l.$$

This implies, by definition, that f is the exponential of $\varphi(F_U)$. □

2nd proof. The complex bundle isomorphism $\mathcal{T}(\mathbb{C}P^k) \oplus \underline{\mathbb{C}} = \bar\eta \oplus \ldots \oplus \bar\eta$ ($k+1$ summands) allows us to calculate the value of a genus on $\mathbb{C}P^k$ explicitly. Let $x = c_1(\bar\eta) \in H^2(\mathbb{C}P^k)$ and let g be the series functionally inverse to f; then

$$\begin{aligned}\varphi[\mathbb{C}P^k] &= \Big(\frac{x}{f(x)}\Big)^{k+1}\langle \mathbb{C}P^k\rangle\\ &= \text{coefficient of } x^k \text{ in } \Big(\frac{x}{f(x)}\Big)^{k+1} = \operatorname{res}_0\Big(\frac{1}{f(x)}\Big)^{k+1}\\ &= \frac{1}{2\pi i}\oint\Big(\frac{1}{f(x)}\Big)^{k+1}dx = \frac{1}{2\pi i}\oint \frac{1}{u^{k+1}}g'(u)du\\ &= \operatorname{res}_0\Big(\frac{g'(u)}{u^{k+1}}\Big) = \text{coefficient of } u^k \text{ in } g'(u).\end{aligned}$$

(Integrating over a closed path around zero makes sense only for convergent power series with coefficients in $\mathbb{C}$, however the result holds for all power series with coefficients in $R \otimes \mathbb{Q}$.) Therefore,

$$g'(u) = \sum_{k\geqslant 0} \varphi[\mathbb{C}P^k]u^k.$$

Theorem F.2.4 then implies that g is the logarithm of the formal group law $\varphi(F_U)$, and thus f is its exponential. □

A parallel theory of genera exists for oriented manifolds. These genera are homomorphisms $\varOmega_{SO} \to R$ from the oriented cobordism ring, and the Hirzebruch construction expresses genera over $\mathbb{Q}$-algebras via certain Pontrjagin characteristic classes (which replace the Chern classes).

Example F.3.4. We take $R = \mathbb{Z}$ in these examples.

1. The top Chern number $c_n(\xi)[M]$ is a Hirzebruch genus, and its corresponding f-series is $f(x) = \frac{x}{1+x}$. The value of this genus on a stably complex manifold $(M, c_\mathcal{T})$ equals the Euler characteristic of M if $c_\mathcal{T}$ is an *almost* complex structure.

2. The *L-genus* $L[M]$ corresponds to the series $f(x) = \tanh(x)$ (the hyperbolic tangent). It is equal to the *signature* $\operatorname{sign}(M)$ by the classical Hirzebruch formula **[122]**.

3. The *Todd genus* $\mathrm{td}[M]$ corresponds to the series $f(x) = 1 - e^{-x}$. The corresponding formal group law is given by (F.3). The Todd genus takes value 1 on every complex projective space $\mathbb{C}P^k$.

The 'trivial' genus $\varepsilon\colon \varOmega_U \to \mathbb{Z}$ corresponding to the series $f(x) = x$ gives rise to the *augmentation transformation* $U^* \to H^*$ from complex cobordism to ordinary cohomology (also known as the *Thom homomorphism*). More generally, for every genus $\varphi\colon \varOmega_U \to R$ and a space X we may set $h^*_\varphi(X) = U^*(X) \otimes_{\varOmega_U} R$. Under certain conditions guaranteeing the exactness of the sequences of pairs (known as the *Landweber exact functor theorem* [**138**]) the functor $h^*_\varphi(\cdot)$ gives rise to a complex-oriented cohomology theory with the coefficient ring R.

As an example of this procedure, consider the genus corresponding to the 1-parameter extension of the multiplicative formal group law, see Example F.1.2. It is also usually called the *Todd genus*, and takes values in the ring $\mathbb{Z}[\beta]$, $\deg\beta = -2$. By interpreting β as the *Bott element* in the complex K-group $\widetilde{K}^0(S^2) = K^{-2}(pt)$ we obtain a homomorphism $\mathrm{td}\colon \varOmega^*_U \to K^*(pt)$. It gives rise to a multiplicative transformation $U^* \to K^*$ from complex cobordism to complex K-theory introduced by Conner and Floyd [**64**]. In this paper Conner and Floyd proved that complex cobordism determines complex K-theory by means of the isomorphism $K^*(X) = U^*(X) \otimes_{\varOmega_U} \mathbb{Z}[\beta]$, where the $\varOmega_U$-module structure on $\mathbb{Z}[\beta]$ is given by the Todd genus. Their proof makes use of the Conner–Floyd Chern classes; several proofs were given subsequently, including one which follows directly from the Landweber exact functor theorem.

Example F.3.5. Another important example from the original work of Hirzebruch is given by the χ_y-*genus*. It corresponds to the series

$$f(x) = \frac{1 - e^{-x(1+y)}}{1 + ye^{-x(1+y)}},$$

where $y \in \mathbb{R}$ is a parameter. Setting $y = -1$, $y = 0$ and $y = 1$ we get the top Chern number $c_n[M]$, the Todd genus $\mathrm{td}[M]$ and the L-genus $L[M] = \mathrm{sign}(M)$ respectively.

If M is a complex manifold then the value $\chi_y[M]$ can be calculated in terms of the Euler characteristics of *Dolbeault complexes* on M, see [**122**].

Exercises.

F.3.6. Show that $[H_{1i}] = [\mathbb{C}P^1][\mathbb{C}P^{i-1}]$ (as complex bordism classes) for $i \geqslant 1$ by calculating the Chern numbers. (The authors do not know of an explicit construction realising this cobordism.)

Bibliography

[1] J. Frank Adams. *On the cobar construction.* Proc. Nat. Acad. Sci. USA **42** (1956), 409–412.

[2] Ron M. Adin. *A new cubical h-vector.* Discrete Math. **157** (1996), no. 1–3, 3–14.

[3] Selman Akbulut and John D. McCarthy. *Casson's Invariant for Oriented Homology* 3-*spheres.* Math. Notes **36**. Princeton Univ. Press, Princeton, N.J., 1990.

[4] Michael F. Atiyah. *Bordism and cobordism.* Proc. Cambridge Philos. Soc. **57** (1961), 200–208.

[5] Michael F. Atiyah. *Convexity and commuting Hamiltonians.* Bull. London Math. Soc. **14** (1982), no. 1, 1–15.

[6] Michael F. Atiyah and Raoul Bott. *A Lefschetz fixed point formula for elliptic complexes. I.* Ann. of Math. **86** (1967), no. 2, 374–407.

[7] Michael F. Atiyah and Friedrich Hirzebruch. *Spin-manifolds and group actions.* In: *Essays in Topology and Related Topics.* Springer-Verlag, Berlin-Heidelberg-New York, 1970, pp. 18–28.

[8] Michèle Audin. *The Topology of Torus Actions on Symplectic Manifolds.* Progress in Mathematics **93**. Birkhäuser, Basel, 1991.

[9] Anton Ayzenberg and Victor Buchstaber. *Moment-angle complexes and polyhedral products for convex polytopes.* Preprint (2010); arXiv:1010.1922.

[10] Eric K. Babson, Louis J. Billera and Clara S. Chan. *Neighborly cubical spheres and a cubical lower bound conjecture.* Israel J. Math. **102** (1997), 297–315.

[11] Anthony Bahri, Martin Bendersky, Frederic R. Cohen, and Samuel Gitler. *The polyhedral product functor: a method of computation for moment-angle complexes, arrangements and related spaces.* Adv. Math. **225** (2010), no. 3, 1634–1668.

[12] Anthony Bahri, Martin Bendersky, Frederic R. Cohen, and Samuel Gitler. *A new topological construction of infinite families of toric manifolds implying fan reduction.* Preprint (2010); arXiv:1011.0094.

[13] Anthony Bahri, Matthias Franz, Dietrich Notbohm, and Nigel Ray. *Classifying weighted projective spaces.* Preprint (2011); arXiv:1108.1938v3 [math.AT].

[14] David Barnette. *Diagrams and Schlegel diagrams.* In: *Combinatorial Structures and Their Applications.* Proc. Calgary Internat. Conference, Calgary 1969. Gordon and Breach, New York, 1970, pp. 1–4.

[15] David Barnette. *The minimum number of vertices of a simple polytope.* Israel J. Math. **10** (1971), 121–125.

[16] David Barnette. *A proof of the lower bound conjecture for convex polytopes.* Pacific J. Math. **46** (1973), 349–354.

[17] David Barnette. *The triangulations of the* 3-*sphere with up to* 8 *vertices.* J. Combinatorial Theory. Ser. A **14** (1973), 37–52.

[18] Ilia V. Baskakov. *Cohomology of K-powers of spaces and the combinatorics of simplicial subdivisions.* Uspekhi Mat. Nauk **57** (2002), no. 5, 147–148 (Russian); Russian Math. Surveys **57** (2002), no. 5, 989–990 (English translation).

[19] Ilia V. Baskakov. *Massey triple products in the cohomology of moment-angle complexes.* Uspekhi Mat. Nauk **58** (2003), no. 5, 199–200 (Russian); Russian Math. Surveys **58** (2003), no. 5, 1039–1041 (English translation).

[20] Ilia V. Baskakov, Victor M. Buchstaber and Taras E. Panov. *Cellular cochain algebras and torus actions.* Uspekhi Mat. Nauk **59** (2004), no. 3, 159–160 (Russian); Russian Math. Surveys **59** (2004), no. 3, 562–563 (English translation).

[21] Victor V. Batyrev. *Quantum cohomology rings of toric manifolds.* In: *Journées de Géométrie Algébrique d'Orsay (Juillet 1992).* Astérisque **218**. Sociéte Mathématique de France, Paris, 1993, pp. 9–34; arXiv:alg-geom/9310004.

[22] Louis J. Billera and Carl W. Lee. *Sufficiency of McMullen's conditions for f-vectors of simplicial polytopes.* Bull. Amer. Math. Soc. **2** (1980), 181–185.

[23] Louis J. Billera and Carl W. Lee. *A proof of the sufficiency of McMullen's conditions for f-vectors of simplicial polytopes.* J. Comb. Theory, Ser. A **31** (1981), 237–255.

[24] Anders Björner and Frank Lutz. *Simplicial manifolds, bistellar flips and 16-vertex triangulation of the Poincaré homology 3-sphere.* Experiment. Math. **9** (2000), no. 2, 275–289.

[25] Anders Björner and Martin Tancer. *Note: Combinatorial Alexander duality—a short and elementary proof.* Discrete Comput. Geom. **42** (2009), no. 4, 586–593; arXiv:0710.1172.

[26] Gerd Blind and Roswitha Blind. *Triangle-free polytopes with few facets.* Arch. Math. (Basel) **58** (1992), no. 6, 599–605.

[27] Frédéric Bosio and Laurent Meersseman. *Real quadrics in $\mathbb{C}^n$, complex manifolds and convex polytopes.* Acta Math. **197** (2006), no. 1, 53–127.

[28] Raoul Bott and Hans Samelson. *Application of the theory of Morse to symmetric spaces.* Amer. J. of Math. **80** (1958), 964–1029.

[29] Raoul Bott and Clifford Taubes. *On the self-linking of knots. Topology and physics.* J. Math. Phys. **35** (1994), no. 10, 5247–5287.

[30] A. K. Bousfield and Victor K. Gugenheim. *On PL de Rham Theory and Rational Homotopy Type.* Memoirs of the Amer. Math. Soc. **9**, no. 179. Amer. Math. Soc., Providence, R.I., 1976.

[31] Glen E. Bredon. *Introduction to Compact Transformation Groups.* Academic Press, New York–London, 1972.

[32] Arne Brønsted. *An Introduction to Convex Polytopes.* Graduate Texts in Math. **90**, Springer-Verlag, New-York, 1983.

[33] Edward M. Brown. *The Hauptvermutung for 3-complexes.* Trans. Amer. Math. Soc. **140** (1969), 173–196.

[34] Winfried Bruns and Joseph Gubeladze. *Combinatorial invariance of Stanley–Reisner rings.* Georgian Math. J. **3** (1996), no. 4, 315–318.

[35] Winfried Bruns and Jürgen Herzog. *Cohen–Macaulay Rings*, revised edition. Cambridge Studies in Adv. Math., vol. **39**, Cambridge Univ. Press, Cambridge, 1998.

[36] David A. Buchsbaum and David Eisenbud. *Algebra structures for finite free resolutions, and some structure theorems for ideals of codimension 3.* Amer. J. Math. **99** (1977), no. 3, 447–485.

[37] Victor M. Buchstaber [Bukhshtaber]. *The Chern–Dold character in cobordisms I.* Mat. Sbornik **83** (1970), no. 4, 575–595 (Russian); Math. USSR Sbornik **12** (1970), no. , 573–594 (English translation).

[38] Victor M. Buchstaber. *Ring of simple polytopes and differential equations.* Trudy Mat. Inst. Steklova **263** (2008), 18–43 (Russian); Proc. Steklov Inst. Math. **263** (2008), 1–25 (English translation).

[39] Victor M. Buchstaber. *Lectures on toric topology.* In *Proceedings of Toric Topology Workshop KAIST 2008.* Trends in Math. **10**, no. 1. Information Center for Mathematical Sciences, KAIST, 2008, pp. 1–64.

[40] Victor M. Buchstaber and Elena V. Koritskaya. *Quasilinear Burgers–Hopf equation and Stasheff polytopes.* Funktsional. Anal. i Prilozhen. **41** (2007), no. 3, 34–47 (Russian); Funct. Anal. and Appl. **41** (2007), no. 3, 196–207 (English translation).

[41] Victor M. Buchstaber and Nickolai Erokhovets. *Polytopes, Hopf algebras and quasi-symmetric functions.* Preprint (2010), arXiv:1011.1536.

[42] Victor M. Buchstaber and Taras E. Panov. *Algebraic topology of manifolds defined by simple polyhedra.* Uspehi Mat. Nauk **53** (1998), no.3, 195–196 (Russian); Russian Math. Surveys **53** (1998), no. 3, 623–625 (English translation).

[43] Victor M. Buchstaber and Taras E. Panov. *Torus actions and the combinatorics of polytopes.* Trudy Mat. Inst. Steklova **225** (1999), 96–131 (Russian); Proc. Steklov Inst. Math. **225** (1999), 87–120 (English translation).

[44] Victor M. Buchstaber and Taras E. Panov. *Torus actions, combinatorial topology and homological algebra.* Uspekhi Mat. Nauk **55** (2000), no. 5, 3–106 (Russian). Russian Math. Surveys **55** (2000), no. 5, 825–921 (English translation).

[45] Victor M. Buchstaber and Taras E. Panov. *Torus Actions and Their Applications in Topology and Combinatorics*. University Lecture Series **24**. Amer. Math. Soc., Providence, R.I., 2002.

[46] Victor M. Buchstaber and Taras E. Panov. *Combinatorics of simplicial cell complexes and torus action*. Trudy Mat. Inst. Steklova **247** (2004), pp. 41–58 (Russian); Proc. Steklov Inst. Math., vol. **247** (2004), pp. 33–49 (English translation).

[47] Victor M. Buchstaber and Taras E. Panov. *Torus Actions in Topology and Combinatorics* (in Russian). MCCME, Moscow, 2004.

[48] Victor M. Buchstaber, Taras E. Panov and Nigel Ray. *Spaces of polytopes and cobordism of quasitoric manifolds*. Moscow Math. J. **7** (2007), no. 2, 219–242; arXiv:math.AT/0609346.

[49] Victor M. Buchstaber, Taras E. Panov and Nigel Ray. *Toric genera*. Internat. Math. Research Notices **16** (2010), 3207–3262; arXiv:0908.3298.

[50] Victor M. Buchstaber [Bukhshtaber] and Nigel Ray. *Toric manifolds and complex cobordisms*. Uspekhi Mat. Nauk **53** (1998), no. 2, 139–140 (Russian); Russian Math. Surveys **53** (1998), no. 2, 371–373 (English translation).

[51] Victor M. Buchstaber and Nigel Ray. *Flag manifolds and the Landweber–Novikov algebra*. Geom. Topol. **2** (1998), 79–101; arXiv:math.AT/9806168.

[52] Victor M. Buchstaber and Nigel Ray. *Tangential structures on toric manifolds, and connected sums of polytopes*. Internat. Math. Res. Notices **4** (2001), 193–219; arXiv:math.AT/0010025.

[53] Victor M. Buchstaber and Nigel Ray. *An invitation to toric topology: vertex four of a remarkable tetrahedron*. In: *Toric Topology* (M. Harada *et al.*, eds.). Contemp. Math., vol. **460**, Amer. Math. Soc., Providence, RI, 2008, pp. 1–27.

[54] Victor M. Buchstaber, Vadim Volodin. *Precise upper and lower bounds for nestohedra*. Izv. Ross. Akad. Nauk, Ser. Mat. **75** (2011), no. 6, 17–46 (Russian); Izv. Math. **75** (2011), no. 6 (English translation).

[55] James W. Cannon. *Shrinking cell-like decompositions of manifolds. Codimension three.* Ann. of Math. (2) **110** (1979), no. 1, 83–112.

[56] Xiangyu Cao, Zhi Lü. *Möbius transform, moment-angle complexes and Halperin–Carlsson conjecture*. Preprint (2009); arXiv:0908.3174.

[57] Michael P. Carr and Satyan L. Devadoss. *Coxeter complexes and graph-associahedra*. Topology Appl. **153** (2006), no. 12, 2155–2168; arXiv:math.QA/0407229.

[58] Ruth Charney and Michael Davis. *The Euler characteristic of a nonpositively curved, piecewise Euclidean manifold*. Pacific J. Math. **171** (1995), no. 1, 117–137.

[59] Suyoung Choi and Jang Soo Kim. *A combinatorial proof of a formula for Betti numbers of a stacked polytope*. Electron. J. Combin. **17** (2010), no. 1, Research Paper 9, 8 pp.; arXiv:math.CO/0902.2444.

[60] Suyoung Choi, Mikiya Masuda and Dong Youp Suh. *Quasi-toric manifolds over a product of simplices*. Preprint (2008); arXiv:0803.2749.

[61] Yusuf Civan and Nigel Ray. *Homotopy decompositions and K-theory of Bott towers*. K-Theory **34** (2005), no. 1, 1–33.

[62] George F. Clements and and Bernt Lindström. *A generalization of a combinatorial theorem of Macaulay*. J. Combinatorial Theory **7** (1969), 230–238.

[63] Pierre E. Conner and Edwin E. Floyd. *Differentiable Periodic Maps*. Academic Press Inc., New York, 1964.

[64] Pierre E. Conner and Edwin E. Floyd. *The Relation of Cobordism to K-theories*. Lecture Notes in Math. **28**, Springer-Verlag, Berlin, 1966.

[65] David A. Cox. *The homogeneous coordinate ring of a toric variety*. J. Algebraic Geom. **4** (1995), no. 1, 17–50; arXiv:alg-geom/9210008.

[66] David A. Cox, John B. Little and Henry K. Schenck. *Toric varieties*. Graduate Studies in Math. **124**. Amer. Math. Soc., Providence, RI, 2011.

[67] H. S. M. Coxeter. *Regular Polytopes*. 3rd edition. Dover Publications, 1973.

[68] Vladimir I. Danilov *The geometry of toric varieties*. Uspekhi Mat. Nauk **33** (1978), no. 2, 85–134 (Russian); Russian Math. Surveys **33** (1978), no. 2, 97–154 (English translation).

[69] Michael W. Davis. *Groups generated by reflections and aspherical manifolds not covered by Euclidean space*. Ann. of Math. **117** (1983), no. 2, 293–324.

[70] Michael W. Davis and Tadeusz Januszkiewicz. *Convex polytopes, Coxeter orbifolds and torus actions*. Duke Math. J. **62** (1991), no. 2, 417–451.

[71] Corrado De Concini and C. Procesi. *Wonderful models of subspace arrangements.* Selecta Math. (N.S.) **1** (1995), no. 3, 459–494.

[72] Mark de Longueville. *The ring structure on the cohomology of coordinate subspace arrangements.* Math. Z. **233** (2000), no. 3, 553–577.

[73] Thomas Delzant. *Hamiltoniens périodiques et images convexes de l'application moment.* Bull. Soc. Math. France **116** (1988), no. 3, 315–339 (French).

[74] Michel Demazure. *Sous-groupes algébriques de rang maximum du groupe de Cremona.* Ann. Sci. École Norm. Sup. (4) **3** (1970), 507–588 (French).

[75] Graham Denham and Alexander I. Suciu. *Moment-angle complexes, monomial ideals, and Massey products.* Pure Appl. Math. Q. **3** (2007), no. 1, 25–60; arXiv:math.AT/0512497.

[76] Natalia Dobrinskaya. *The classification problem for quasitoric manifolds over a given simple polytope.* Funktsional. Anal. i Prilozhen. **35** (2001), no. 2, 3–11 (Russian); Funct. Anal. and Appl. **35** (2001), no. 2, 83–89 (English translation).

[77] Nikolai P. Dolbilin, Mikhail A. Shtan'ko and Mikhail I. Shtogin. *Cubic manifolds in lattices.* Izv. Ross. Akad. Nauk Ser. Mat. **58** (1994), no. 2, 93–107 (Russian); Russian Acad. Sci. Izv. Math. **44** (1995), no. 2, 301–313 (English translation).

[78] Albrecht Dold. *Geometric cobordism and the fixed point transfer.* In: *Algebraic Topology, Proceedings Vancouver 1977.* Lecture Notes in Math. **673**. Springer Verlag, Berlin, 1978, pp. 32–87.

[79] Johannes J. Duistermaat and G. J. Heckman. *On the variation in the cohomology of the symplectic form of the reduced phase space.* Invent. Math. **69** (1982), no. 2, 259–268.

[80] Art M. Duval. *Free resolutions of simplicial posets.* J. Algebra **188** (1997), 363–399.

[81] Robert D. Edwards. *The double suspension of a certain homology 3-sphere is* S^5. Notices AMS **22**, A-334 (1975).

[82] Robert D. Edwards. *The topology of manifolds and cell-like maps.* In: Proceedings of the International Congress of Mathematicians (Helsinki, 1978). Acad. Sci. Fennica, Helsinki, 1980, pp. 111–127.

[83] Samuel Eilenberg and John C. Moore. *Homology and fibrations. I.* Comment. Math. Helv. **40** (1966), 199–236.

[84] Davin Eisenbud. *Commutative Algebra with a View Toward Algebraic Geometry.* Graduate Texts in Math. **150**, Springer-Verlag, New-York, 1995.

[85] Nickolay Yu. Erokhovets. *The Bukhshtaber invariant of simple polytopes.* Uspekhi Mat. Nauk **63** (2008), no. 5, 187–188 (Russian); Russian Math. Surveys 63 (2008), no. 5, 962–964 (English translation); arXiv:0908.3407.

[86] Günter Ewald. *Combinatorial Convexity and Algebraic Geometry.* Graduate Texts in Math. **168**, Springer-Verlag, New-York, 1996.

[87] Günter Ewald and Christoph Schulz. *Nonstarshaped spheres.* Arch. Math. (Basel) **59** (1992), no. 4, 412–416.

[88] Eva Maria Feichtner and Bernd Sturmfels. *Matroid polytopes, nested sets and Bergman fans.* Port. Math. (N.S.) **62** (2005), no. 4, 437–468; arXiv:math.CO/0411260.

[89] Yves Félix, Stephen Halperin and Jean-Claude Thomas. *Rational Homotopy Theory.* Springer-Verlag, 2001.

[90] John Fogarty, Frances Kirwan and David Mumford. *Geometric Invariant Theory.* Third Edition. Springer-Verlag, Berlin, 1994.

[91] Anatoliy T. Fomenko and Dmitriy B. Fuks. *A Course in Homotopic Topology* (in Russian). Nauka, Moscow, 1989.

[92] Sergey Fomin and Andrei Zelevinsky. *Cluster algebras. I. Foundations.* J. Amer. Math. Soc. **15** (2002), no. 2, 497–529.

[93] Matthias Franz. *On the integral cohomology of smooth toric varieties.* Proc. Steklov Inst. Math. **252** (2006), 53–62; arXiv:math.AT/0308253.

[94] Michael H. Freedman and Frank Quinn. *Topology of 4-manifolds.* Princeton Mathematical Series **39**. Princeton Univ. Press, Princeton, N.J., 1990.

[95] Yukiko Fukukawa and Mikiya Masuda. *Buchstaber invariants of skeleta of a simplex.* Osaka J. Math. **48** (2011), 549–582; arXiv:0908.3448.

[96] William Fulton. *Introduction to Toric Varieties.* Ann. of Math. Studies **131**. Princeton Univ. Press, Princeton, N.J., 1993.

[97] Światosław R. Gal. *Real root conjecture fails for five and higher dimensional spheres.* Discrete Comput. Geom. **34** (2005), no. 2, 260–284; arXiv:math/0501046.

[98] Israel M. Gelfand, Mikhail M. Kapranov, and Andrei V. Zelevinsky. *Discriminants, Resultants, and Multidimensional Determinants.* Birkhäuser, Boston, 1994.

[99] Samuel Gitler and Santiago López de Medrano. *Intersections of quadrics, moment-angle manifolds and connected sums.* Preprint (2009); arXiv:0901.2580.

[100] Evgeniy S. Golod. *On the cohomology of some local rings* (Russian); *Soviet Math. Dokl.* **3** (1962), 745–749.

[101] Mark Goresky, R. Kottwitz and Robert MacPherson. *Equivariant cohomology, Koszul duality and the localisation theorem.* Invent. Math. **131** (1998), no. 1, 25–83.

[102] Mark Goresky and Robert MacPherson. *Stratified Morse Theory.* Springer-Verlag, Berlin–New York, 1988.

[103] Jelena Grbić and Stephen Theriault. *The homotopy type of the complement of the codimension-two coordinate subspace arrangement. Uspekhi Mat. Nauk* **59** (2004), no. 6, 203–204 (Russian); *Russian Math. Surveys* **59** (2004), no. 6, 1207–1209 (English translation).

[104] Jelena Grbić and Stephen Theriault. *The homotopy type of the complement of a coordinate subspace arrangement.* Topology **46** (2007), no. 4, 357–396; arXiv:math/0601279.

[105] Mikhail Gromov. *Hyperbolic groups.* In: *Essays in Group Theory*, S. M. Gersten, ed. MSRI Publ. **8**. Springer-Verlag, New York, 1987, pp. 75–263.

[106] Michael Grossberg and Yael Karshon. *Bott towers, complete integrability, and the extended character of representations.* Duke Math. J. **76** (1994), 23–58.

[107] Branko Grünbaum. *Convex Polytopes.* Second edition. Graduate Texts in Math. **221**. Springer-Verlag, New York, 2003.

[108] Branko Grünbaum and V. P. Sreedharan. *An enumeration of simplicial 4-polytopes with 8 vertices.* J. Combinatorial Theory **2** (1967), 437–465.

[109] Victor Guillemin. *Moment Maps and Combinatorial Invariants of Hamiltonian T^n-spaces.* Progress in Math. **122**. Birkhaüser, Boston, 1994.

[110] Victor Guillemin and Shlomo Sternberg. *Convexity properties of the moment mapping.* Inv. Math. **67** (1982), no. 3, 491–513.

[111] Victor W. Guillemin and Catalin Zara. *Equivariant de Rham theory and graphs.* Asian J. Math. **3** (1999), no. 1, 49–76; arXiv:math.DG/9808135.

[112] Victor W. Guillemin and Catalin Zara. *One-skeleta, Betti numbers and equivariant cohomology.* Duke Math. J. **107** (2001), no. 2, 283–349; arXiv:math/9903051.

[113] Tor Gulliksen and Gerson Levin. *Homology of Local Rings.* Queen's Papers in Pure and Applied Mathematics, vol. **20**. Queen's University, Kingston, Ontario, 1969.

[114] Stephen Halperin. *Rational homotopy and torus actions.* In: *Aspects of Topology.* London Math. Soc. Lecture Note Series, vol. **93**, Cambridge Univ. Press, Cambridge, 1985, pp. 293–306.

[115] Allen Hatcher. *Algebraic Topology.* Cambridge Univ. Press, Cambridge, 2002.

[116] Akio Hattori. *Symplectic manifolds with semi-free Hamiltonian S^1-action.* Tokyo J. Math. **15** (1992), no. 2, 281–296.

[117] Akio Hattori and Mikiya Masuda, *Theory of multi-fans.* Osaka J. Math. **40** (2003), 1–68; arXiv:math/0106229.

[118] Akio Hattori and Mikiya Masuda, *Elliptic genera, torus manifolds and multi-fans.* Internat. J. of Math. **16** (2005), no. 9, 957–998; arXiv:math/0107014.

[119] Akio Hattori and Tomoyoshi Yoshida. *Lifting compact group actions in fiber bundles.* Japan. J. Math. **2** (1976), 13–25.

[120] Jürgen Herzog, Victor Reiner and Volkmar Welker. *Componentwise linear ideals and Golod rings. Michigan Math. J.* **46** (1999), no. 2), 211–223.

[121] Friedrich Hirzebruch. *Komplexe Mannigfaltigkeiten.* (German) Proc. Internat. Congress Math, 1958. Cambridge Univ. Press, New York, 1960, pp. 119–136.

[122] Friedrich Hirzebruch. *Topological Methods in Algebraic Geometry*, 3rd edition. Springer-Verlag, Berlin–Heidelberg, 1966.

[123] Friedrich Hirzebruch, Thomas Berger and Rainer Jung. *Manifolds and Modular Forms*, Second Edition. A Publication from the Max-Planck-Institut für Mathematik, Bonn. Friedr. Vieweg & Sohn, Braunschweig, 1994.

[124] Melvin Hochster. *Cohen–Macaulay rings, combinatorics, and simplicial complexes.* In: *Ring Theory II (Proc. Second Oklahoma Conference).* B. R. McDonald and R. Morris, eds. Dekker, New York, 1977, pp. 171–223.

[125] J. F P. Hudson. *Piecewise Linear Topology.* W. A. Benjamin, Inc., New York–Amsterdam, 1969.

[126] Dmitri Ilinskii. *On the almost free action of one-dimensional torus on a toric variety.* Mat. Sbornik **197** (2006), no. 5, 51–74 (Russian); Sbornik Math. **197** (2006), no. 5, 681–703 (English translation).

[127] Ivan Izmestiev. *Geometry of manifolds with a regular action of the torus* (in Russian). Ph.D. thesis, Moscow State University, 2000.

[128] Klaus Jänich. *Differenzierbare Mannigfaltigkeiten mit Rand als Orbiträume differenzierbarer G-Mannigfaltigkeiten ohne Rand.* (in German) Topology **5** (1966), 301–320.

[129] Tetsuro Kawasaki. *Cohomology of twisted projective spaces and lens complexes.* Math. Ann. **206** (1973), 243–248.

[130] Gil Kalai. *Rigidity and the lower bound Theorem. I.* Inv. Math. **88** (1987), no. 1, 125–151.

[131] Kalle Karu. *Hard Lefschetz theorem for nonrational polytopes.* Invent. Math. **157** (2004), no. 2, 419–447.

[132] Askold G. Khovanskii [Hovanskii]. *Newton polyhedra and toric varieties.* Funk. Anal. i Priloz **11** (1977), no. 4, 56–64 (Russian); Funct. Anal. Appl. **11** (1977), no. 4, 289–296 (English translation).

[133] Askold G. Khovanskii. *Hyperplane sections of polyhedra, toric varieties and discrete groups in Lobachevskii space.* Funk. Anal. i Prilozhen. **20** (1986), no. 1, 50–61 (Russian); Funct. Anal. Appl. **20** (1986), no. 1, 41–50 (English translation).

[134] Robion C. Kirby and Laurent C. Siebenmann. *Foundational Essays on Topological Manifolds, Smoothings, and Triangulations.* Ann. of Math. Stud. **88**, Princeton Univ. Press, Princeton, N.J., 1977.

[135] Victor Klee. *A combinatorial analogue of Poincaré's duality theorem.* Canad. J. Math. **16** (1964), 517–531.

[136] Victor Klee and Peter Kleinschmidt, *Convex polytopes and related complexes.* In *Handbook of Combinatorics, Vol. 1,2.* R. L. Graham et al, eds. Elsevier Science B.V., Amsterdam, 1995, pp. 875–917.

[137] Igor M. Krichever. *Formal groups and the Atiyah-Hirzebruch formula.* Izv. Akad. Nauk SSSR **38** (1974), no. 6, 1289–1304 (Russian); Math. USSR Izv. **8** (1974), no. 6, 1271–1285 (English translation).

[138] Peter S. Landweber. *Homological properties of comodules over $MU_*(MU)$ and $BP_*(BP)$.* Amer. J. Math. **98** (1976), no. 3, 591–610.

[139] Michel Lazard. *Sur le groupes de Lie formels à un paramètre.* Bull. Soc. Math. France **83** (1955), no. 3, 251–274.

[140] Daniel Lehmann. *Théorie homotopique des forms différentielles (d'aprés D. Sullivan).* Astérisque **45**, Sociéte Mathématique de France, Paris, 1977.

[141] Ivan Limonchenko. *Bigraded Betti numbers of some simple polytopes.* Math. Notes (2012), to appear; arXiv:1101.0066.

[142] Zhi Lü and Taras Panov. *Moment-angle complexes from simplicial posets.* Cent. Eur. J. Math. **9** (2011), no. 4, 715–730; arXiv:0912.2219.

[143] Frank H. Lutz. *Triangulated manifolds with few vertices and vertex-transitive group actions.* Dissertation, Technischen Universität Berlin, Berlin, 1999. Berichte aus der Mathematik, Shaker Verlag, Aachen, 1999.

[144] F. S. Macaulay. *Some properties of enumeration in the theory of modular systems.* Proc. London Math. Soc. **26** (1927), 531–555.

[145] Saunders Mac Lane. *Homology.* Springer-Verlag, Berlin, 1963.

[146] Hiroshi Maeda, Mikiya Masuda and Taras Panov. *Torus graphs and simplicial posets.* Advances in Math. **212** (2007), no. 2, 458–483; arXiv:math/0511582.

[147] Peter Mani. *Spheres with few vertices.* J. Combinatorial Theory. Ser. A **13** (1972), 346–352.

[148] Mikiya Masuda. *Unitary toric manifolds, multi-fans and equivariant index.* Tohoku Math. J. **51** (1999), no. 2, 237–265.

[149] Mikiya Masuda. *h-vectors of Gorenstein* simplicial posets.* Adv. Math. **194** (2005), no. 2, 332–344; arXiv:math/0305203.

[150] Mikiya Masuda and Taras Panov. *On the cohomology of torus manifolds.* Osaka J. Math. **43** (2006), 711–746; arXiv:math/0306100.

[151] Mikiya Masuda and Taras Panov. *Semifree circle actions, Bott towers, and quasitoric manifolds.* Mat. Sbornik **199** (2008), no. 8, 95–122 (Russian); Sbornik Math. **199** (2008), no. 7–8, 1201–1223 (English translation); arXiv:math.AT/0607094.

[152] Sergei V. Matveev. *Algorithmic Topology and Classification of* 3*-manifolds.* Algorithms and Computation in Mathematics, **9**. Springer-Verlag, Berlin, 2003.

[153] John McCleary. *A User's Guide to Spectral Sequences,* 2nd edition. Cambdridge Univ. Press, Cambridge, 2001.

[154] Peter McMullen. *The maximum numbers of faces of a convex polytope.* Mathematika **17** (1970), 179–184.

[155] Peter McMullen. *The numbers of faces of simplicial polytopes.* Israel J. Math. **9** (1971), 559–570.

[156] Peter McMullen. *On simple polytopes.* Invent. Math. **113** (1993), no. 2, 419–444.

[157] Peter McMullen. *Weights on polytopes.* Discrete Comput. Geom. **15** (1996), no. 4, 363–388.

[158] Peter McMullen and G. Shephard. *Convex Polytopes and the Upper Bound Conjecture.* London Math. Soc. Lecture Notes Series **3**, Cambridge Univ. Press, Cambridge, 1971.

[159] Peter McMullen and David Walkup. *A generalized lower bound conjecture for simplicial polytopes.* Mathematika **18** (1971), 264–273.

[160] Ezra Miller and Bernd Sturmfels. *Combinatorial Commutative Algebra.* Graduate Texts in Math. **227**. Springer-Verlag, New York, 2005.

[161] John Milnor. *On the cobordism ring* Ω^* *and a complex analogue. I.* Amer. J. Math. **82** (1960), 505–521.

[162] John W. Milnor and James D. Stasheff. *Characteristic classes.* Princeton Univ. Press, Princeton, N.J., 1974.

[163] Andrey Mironov and Taras Panov. *Intersections of quadrics, moment-angle manifolds, and Hamiltonian-minimal Lagrangian embeddings.* Preprint (2011); arXiv:1103.4970.

[164] Edwin E. Moise. *Geometric Topology in Dimensions* 2 *and* 3. Graduate Texts in Math. **47**. Springer-Verlag, New York-Heidelberg, 1977.

[165] James R. Munkres. *Elementary Differential Topology.* Revised edition. Annals of Math. Studies **54**. Princeton Univ. Press, Princeton, N.J., 1966.

[166] Oleg R. Musin. *Actions of the circle on homotopy complex projective spaces.* Mat. Zametki **28** (1980), no. 1, 139–152 (Russian). Math. Notes **28** (1980), no. 1–2, 533–539 (English translation).

[167] A. Nabutovsky. *Geometry of the space of triangulations of a compact manifold.* Comm. Math. Phys. **181** (1996), no. 2, 303–330.

[168] Dietrich Notbohm and Nigel Ray. *On Davis–Januszkiewicz homotopy types I; formality and rationalisation.* Algebraic & Geometric Topology **5** (2005), 31–51; arXiv:math/0311167.

[169] Sergei P. Novikov. *Some problems in the topology of manifolds connected with the theory of Thom spaces.* Dokl. Akad. Nauk SSSR **132** (1960), 1031–1034 (Russian). Soviet Math. Dokl. **1** (1960), 717–720 (English translation).

[170] Sergei P. Novikov. *Homotopy properties of Thom complexes.* Mat. Sbornik **57(99)** (1962), 407–442 (Russian).

[171] Sergei P. Novikov. *Methods of algebraic topology from the point of view of cobordism theory.* Izv. Akad. Nauk SSSR, Ser. Mat. **31** (1967), no. 4, 855–951 (Russian). Math. USSR – Izv. **1** (1967), no. 4, 827–913 (English translation).

[172] Sergei P. Novikov. *Topology I.* Encyclopaedia Math. Sci. **12**. Springer-Verlag, Berlin, 1996.

[173] Tadao Oda. *Convex Bodies and Algebraic Geometry. An introduction to the theory of toric varieties.* Springer-Verlag, New York, 1988.

[174] Udo Pachner. *Konstruktionsmethoden und das kombinatorische Homöomorphieproblem für Triangulationen kompakter semilinearer Mannigfaltigkeiten.* Abh. Math. Sem. Univ. Hamburg **57** (1987), 69–86.

[175] Udo Pachner. *P.L. homeomorphic manifolds are equivalent by elementary shellings.* European J. Combin. **12** (1991), no. 2, 129–145.

[176] Taras E. Panov. *Classification up to cobordism of manifolds that carry a simple action of the group* $\mathbb{Z}/p$. Mat. Zametki **63** (1998), no. 2, 260–268 (Russian). Math. Notes **63** (1998), no. 1–2, 225–232 (English translation); arXiv:math.AT/9908166.

[177] Taras E. Panov. *Calculation of Hirzebruch genera of manifolds with* $\mathbb{Z}/p$ *group action in terms of the invariants of the action.* Izv. Ross. Acad. Nauk, Ser. Mat. **62** (1998),

no. 3, 87–120 (Russian); Izv. Math. **62** (1998), no. 3, 515–548 (English translation); arXiv:math.AT/9909081.

[178] Taras E. Panov. *Hirzebruch genera of manifolds with torus actions.* Izv. Ross. Akad. Nauk, Ser. Mat. **65** (2001), no. 3, 123–138 (Russian); Izv. Math. **65** (2001), no. 3, 543–556 (English translation); arXiv: math.AT/9910083.

[179] Taras Panov. *Cohomology of face rings, and torus actions.* In: *Surveys in Contemporary Mathematics.* London Math. Soc. Lecture Note Series, vol. **347**, Cambridge, U.K., 2008, pp. 165–201; arXiv:math.AT/0506526.

[180] Taras Panov. *Toric Kempf–Ness sets.* Trudy Mat. Inst. Steklova **263** (2008), 159–172 (Russian); Proc. Steklov Inst. Math. **263** (2008), pp. 150–162 (English translation); arXiv:math.AG/0603556.

[181] Taras Panov. *Moment-angle manifolds and complexes.* Trends in Mathematics - New Series. Information Center for Mathematical Sciences, KAIST. Vol. 12 (2010), no. 1, pp. 43–69; arXiv:1008.5047.

[182] Taras Panov and Nigel Ray. *Categorical aspects of toric topology.* In: *Toric Topology*, M. Harada *et al.*, eds. Contemp. Math., vol. **460**. Amer. Math. Soc., Providence, RI, 2008, pp. 293–322; arXiv:0707.0300.

[183] Taras Panov, Nigel Ray and Rainer Vogt. *Colimits, Stanley–Reiner algebras, and loop spaces.* In: *Categorical decomposition techniques in algebraic topology (Isle of Skye, 2001).* Progress in Math., vol. **215**, Birkhäuser, Basel, 2004, pp. 261–291; arXiv:math/0202081.

[184] Taras Panov and Yuri Ustinovsky. *Complex-analytic structures on moment-angle manifolds.* Moscow Math. J. **12** (2012), no. 1, 149–172; arXiv:1008.4764.

[185] Lev S. Pontryagin. *Foundations of Combinatorial Topology.* Graylock Press, Rochester, N.Y., 1952.

[186] Gerald J. Porter. *The homotopy groups of wedges of suspensions.* Amer. J. Math. **88** (1966) 655—663.

[187] Alexander Postnikov. *Permutohedra, associahedra, and beyond.* Int. Math. Res. Not. **2009**, no. 6, 1026–1106; arXiv:math.CO/0507163.

[188] Alexander Postnikov, Victor Reiner, L. Williams. *Faces of generalized permutohedra.* Doc. Math. **13** (2008), 207–273; arXiv:math.CO/0609184.

[189] J. Scott Provan and Louis J. Billera. *Decompositions of simplicial complexes related to diameters of convex polyhedra.* Math. Oper. Res. **5** (1980), no. 4, 576–594.

[190] Volker Puppe. *Multiplicative aspects of the Halperin–Carlsson conjecture.* Georgian Math. J. **16** (2009), no. 2, 369–379; arXiv:0811.3517.

[191] Daniel Quillen. *Rational homotopy theory.* Ann. of Math. (2) **90** (1969), no. 2, 205–295.

[192] Daniel G. Quillen. *On the formal group laws of unoriented and complex cobordism theory.* Bull. Amer. Math. Soc. **75** (1969), no. 6, 1293–1298.

[193] Daniel G. Quillen. *Elementary proofs of some results of cobordism theory using steenrod operations. Adv. Math.* **7** (1971), 29–56.

[194] Andrew Ranicki. *On the Hauptvermutung.* In: *The Hauptvermutung Book. K*-Monogr. Math. **1**, Kluwer Acad. Publ., Dordrecht, 1996, pp. 3–31.

[195] Nigel Ray. *On a construction in bordism theory. Proc. Edin. Math. Soc.* **29** (1986), 413–422.

[196] Gerald A. Reisner. *Cohen–Macaulay quotients of polynomial rings*, Adv. Math. **21** (1976), no. 1, 30–49.

[197] Colin P. Rourke and Brian J. Sanderson. *Introduction to Piecewise-Linear Topology.* Springer-Verlag, Berlin, 1972.

[198] J. Hyam Rubinstein. *An algorithm to recognise the* 3*-sphere.* Proceedings of the International Congress of Mathematicians, Vol 1, 2 (Zürich, 1994), 601-–611; Birkhäuser, Basel, 1995.

[199] Yuli B. Rudyak. *Piecewise linear structures on topological manifolds.* arXiv:math.AT/0105047 (2001).

[200] Gerald W. Schwarz. *The topology of algebraic quotients.* In: *Topological Methods in Algebraic Transformation Groups.* Progr. in Math **80**. Birkhäuser, Boston, 1989, pp. 135–151.

[201] Igor R. Shafarevich. *Basic Algebraic Geometry. 2. Schemes and complex manifolds.* Second edition. Translated from the 1988 Russian edition by Miles Reid. Springer-Verlag, Berlin, 1994.

[202] Larry Smith. *Homological algebra and the Eilenberg–Moore spectral sequence.* Trans. Amer. Math. Soc. **129** (1967), 58–93.

[203] Richard P. Stanley. *The upper bound conjecture and Cohen-Macaulay rings.* Studies in Appl. Math. **54** (1975), no. 2, 135–142.

[204] Richard P. Stanley. *The number of faces of simplicial convex polytope.* Adv. Math. **35** (1980), no. 3, 236–238.

[205] Richard P. Stanley. *Enumerative Combinatorics*, Vol. 1. Wadsworth and Brooks/Cole, Monterey, California, 1986.

[206] Richard P. Stanley. *Generalized H-vectors, intersection cohomology of toric varieties, and related results.* In: *Commutative algebra and combinatorics (Kyoto, 1985)*, M. Nagata and H. Matsumure, eds. Adv. Stud. Pure Math. **11**, North-Holland, Amsterdam, 1987, pp. 187–213.

[207] Richard P. Stanley. *f-vectors and h-vectors of simplicial posets.* J. Pure Appl. Algebra. **71** (1991), 319–331.

[208] Richard P. Stanley. *Combinatorics and Commutative Algebra*, second edition. Progr. in Math. **41**. Birkhäuser, Boston, 1996.

[209] Richard P. Stanley. *Positivity problems and conjectures in algebraic combinatorics.* In: *Mathematics: Frontiers and Perspectives*, V. I. Arnold *et al*, eds. Amer. Math. Soc., Providence, R.I., 2000, pp. 295–319.

[210] James D. Stasheff. *Homotopy associativity of H-spaces. I.* Transactions Amer. Math. Soc. **108** (1963), 275–292.

[211] James D. Stasheff. *From operads to "physically" inspired theories.* In: *Operads: Proceedings of Renaissance Conferences (Hartford, CT/Luminy, 1995).* Contemp. Math. **202** (1997), 53–81.

[212] Robert E. Stong. *Notes on Cobordism Theory.* Princeton Univ. Press, Princeton, 1968.

[213] Dennis Sullivan. *Infinitesimal computations in topology.* Publ. I.H.E.S. **47** (1977), 269–331.

[214] R. H. Szczarba. *On tangent bundles of fibre spaces and quotient spaces.* Amer. J. Math. **86** (1964), 685–697.

[215] Naoki Terai and Takayuki Hibi. *Computation of Betti numbers of monomial ideals associated with stacked polytopes.* Manuscripta Math. **92** (1997), no. 4, 447–453.

[216] René Thom. *Quelques propriétés globales des variétés différentiables.* Comment. Math. Helv. **28** (1954), 17–86.

[217] Emery Thomas. *Complex structures on real vector bundles.* Amer. J. Math. **89** (1967), 887–908.

[218] Vladlen Timorin. *An analogue of the Hodge–Riemann relations for simple convex polytopes.* Uspekhi Mat. Nauk **54** (1999), no. 2, 113–162 (Russian); Russian Math. Surveys **54** (1999), no. 2, 381–426 (English translation).

[219] Valerio Toledano Laredo. *Quasi-Coxeter algebras, Dynkin diagram cohomology, and quantum Weyl groups.* Int. Math. Res. Pap. IMRP **2008**, Article ID rpn009, 167 pp.

[220] Susan Tolman and Jonathan Weitsman. *On semifree symplectic circle actions with isolated fixed points.* Topology **39** (2000), no. 2, 299–309.

[221] Yury Ustinovsky. *Toral rank conjecture for moment-angle complexes.* Mat. Zametrki **90** (2011), no. 2, 300–305 (Russian); Math. Notes **90** (2011), no. 1–2, 279–283 (English translation); arXiv:0909.1053.

[222] Yury Ustinovsky. *On almost free torus actions and Horrocks conjecture.* Far Eastern Mathematical Journal **12** (2012), no. 1 (Russian); arXiv:.

[223] Ernest B. Vinberg. *Discrete linear groups generated by reflections.* Izv. Akad. Nauk SSSR Ser. Mat. **35** (1971), 1072–1112 (Russian); Math. USSR Izvestija **5** (1971), no. 5, 1083–1119 (English translation).

[224] Ernest B. Vinberg and Vladimir L. Popov. *Invariant theory.* In: *Algebraic Geometry 4.* VINITI, Moscow, 1989, pp. 137–314 (Russian); Encyclopaedia Math. Sci. **55** Springer, Heidelberg, 1994, pp. 123–284 (English translation).

[225] I. A. Volodin, V. E. Kuznecov, A. T. Fomenko. *The problem of the algorithmic discrimination of the standard three-dimensional sphere.* Appendix by S. P. Novikov. Uspekhi Mat. Nauk **29** (1974), no. 5, 71–168 (Russian); Russian Math. Surveys **29** (1974), no. 5, 71–172 (English translation).

[226] V. A. Yemelichev, M. M. Kovalev and M. K. Kravtsov. *Polytopes, Graphs and Optimisation.* Translated from the Russian by G. H. Lawden. Cambridge Univ. Press, Cambridge, 1984.

[227] C. T. C. Wall. *Determination of the cobordism ring.* Ann. of Math. (2) **72** (1960), 292-–311.

[228] Günter M. Ziegler. *Lectures on Polytopes*. Graduate Texts in Math., vol. **152**, Springer-Verlag, New York, 1995.